


# Periodic Status Review for the Peregrine Falcon



**Mark S. Vekasy**  
*Washington Department of*  
**FISH AND WILDLIFE**  
*Wildlife Program*

The Washington Department of Fish and Wildlife maintains a list of endangered, threatened, and sensitive species (Washington Administrative Codes 232-12-014 and 232-12-011, Appendix A). In 1990, the Washington Wildlife Commission adopted listing procedures developed by a group of citizens, interest groups, and state and federal agencies (Washington Administrative Code 232-12-297). The procedures include how species listings will be initiated, criteria for listing and delisting, a requirement for public review, the development of recovery or management plans, and the periodic review of listed species.

The Washington Department of Fish and Wildlife is directed to conduct reviews of each endangered, threatened, or sensitive wildlife species at least every five years after the date of its listing by the Washington Fish and Wildlife Commission. The periodic status reviews are designed to include an update of the species status report to determine whether the status of the species warrants its current listing status or deserves reclassification. The agency notifies the general public and specific parties who have expressed their interest to the Department of the periodic status review at least one year prior to the five-year period so that they may submit new scientific data to be included in the review. The agency notifies the public of its recommendation at least 30 days prior to presenting the findings to the Fish and Wildlife Commission. In addition, if the agency determines that new information suggests that the classification of a species should be changed from its present state, the agency prepares documents to determine the environmental consequences of adopting the recommendations pursuant to requirements of the State Environmental Policy Act.

This document is the Draft Periodic Status Review for the Peregrine Falcon. It contains a review of information pertaining to the status of the Peregrine Falcon in Washington. It was reviewed by species experts and will be available for a 90-day public comment period. All comments received will be considered during the preparation of the final periodic status review. The Department intends to present the results of this periodic status review to the Fish and Wildlife Commission at a meeting in November.

**Submit written comments on this report by e-mail by 10 October 2016 to:**  
T&Epubliccom@dfw.wa.gov

**Or by mail to:**

**Listing and Recovery Section Manager, Wildlife Program  
Washington Department of Fish and Wildlife  
600 Capitol Way North, Olympia, Washington 98501-1091**

**This report should be cited as:**

Vekasy, M. S. 2016. Draft Periodic status review for the Peregrine Falcon. Washington Department of Fish and Wildlife, Olympia, Washington. 18+iii pp.

*Cover background photo of Sandridge eyrie by Glenn Phillips;  
Peregrine photographer unknown*



*This work was supported in part by  
personalized and endangered species  
license plates*



The Washington Department of Fish and Wildlife maintains a list of endangered, threatened, and sensitive species (Washington Administrative Codes 232-12-014 and 232-12-011). In 1990, the Washington Wildlife Commission adopted listing procedures developed by a group of citizens, interest groups, and state and federal agencies (Washington Administrative Code 232-12-297). The procedures include how species listings will be initiated, criteria for listing and delisting, a requirement for public review, the development of recovery or management plans, and the periodic review of listed species.

The Washington Department of Fish and Wildlife is directed to conduct reviews of each endangered, threatened, or sensitive wildlife species at least every five years after the date of its listing by the Washington Fish and Wildlife Commission. The periodic status reviews are designed to include an update of the species status report to determine whether the status of the species warrants its current listing status or deserves reclassification. The agency notifies the general public and specific parties who have expressed their interest to the Department of the periodic status review at least one year prior to the five-year period so that they may submit new scientific data to be included in the review. The agency notifies the public of its findings at least 30 days prior to presenting the findings to the Fish and Wildlife Commission. In addition, if the agency determines that new information suggests that the classification of a species should be changed from its present state, the agency prepares documents to determine the environmental consequences of adopting the recommendations pursuant to requirements of the State Environmental Policy Act.

This document is the Draft Periodic Status Review for the Peregrine Falcon. It contains a review of information pertaining to the status of the peregrine falcon in Washington. It was reviewed by species experts and will be available for a 90-day public comment period. All comments received will be considered during the preparation of the final periodic status review. The Department will present the results of this periodic status review to the Fish and Wildlife Commission for action at the November 2016 meeting.

**Submit written comments on this report by e-mail to:**

T&Epubliccom@dfw.wa.gov

**Or by mail to:**

Listing and Recovery Section Manager, Wildlife Program  
Washington Department of Fish and Wildlife  
600 Capitol Way North, Olympia, Washington 98501-1091

This report should be cited as:

Vekasy, M. S. 2016. Draft periodic status review for the peregrine falcon in Washington. Washington Department of Fish and Wildlife, Olympia, Washington. 19pp.

**Draft**  
**Periodic Status Review for the Peregrine Falcon**  
**in Washington**



Prepared by

Mark S. Vekasy

Washington Department of Fish and Wildlife  
Wildlife Program  
1340 N 13<sup>th</sup> Ave  
Walla Walla, WA 99362

July 2016

## TABLE OF CONTENTS

ACKNOWLEDGEMENTS.....	ii
EXECUTIVE SUMMARY .....	iii
INTRODUCTION .....	1
SPECIES BACKGROUND.....	1
POPULATIONS AND HABITAT STATUS .....	4
FACTORS AFFECTING CONTINUED EXISTENCE.....	8
MANAGEMENT ACTIVITIES.....	11
CONCLUSIONS AND RECOMMENDATION.....	11
LITERATURE CITED .....	13

## LIST OF FIGURES AND APPENDICES

Figure 1. Peregrine falcon.....	1
Figure 2. Distribution of occupied peregrine falcon territories in Washington, 2016.....	1
Figure 3. Trend in occupied territories and nesting success in Washington, 1990-2009. ....	7
Appendix A. Occupancy and reproductive success at American peregrine falcon nesting territories included in the federal post-delisting monitoring, Washington, 1999-2012.....	18

## ACKNOWLEDGEMENTS

Funding for the preparation of this periodic status review came from Washington State background license plates for endangered wildlife. Washington State personalized license plates, and a State Wildlife Grant from the U.S. Fish and Wildlife Service. This draft was improved by reviews by Clifford Anderson, David Anderson, Joe Buchanan, Myke Chutter, Paul DeBruyn, Fred Dobler, Gerald Hayes, Kelly McAllister, Joel Pagel, Doug Pineo, Sue Thomas, Patricia Thompson, Dan Varland, and Jim Watson.

## EXECUTIVE SUMMARY

Peregrine falcons (*Falco peregrinus*) exhibited well-documented population declines across North America and much of their global range following the widespread use of DDT shortly after the Second World War. The peregrine falcon was listed nationally as an endangered species by the U.S. Fish and Wildlife Service (USFWS) in 1970 and by the Washington Fish and Wildlife Commission in 1980 when only five pairs were found to be nesting statewide. With the restriction placed on the use of DDT, the peregrine population has recovered and was removed from the federal endangered species list in 1999. In 2002 the species was reclassified as a state sensitive species after >70 territories were found occupied.

In 2004, the USFWS and Washington Department of Fish and Wildlife began allowing small numbers of peregrine falcon nestlings to be taken for falconry, and in 2010 the regulations were modified to include trapping of first-year Washington falcons in the vicinity of nest sites. WDFW last completed comprehensive surveys of peregrine falcon territories in 2009. In that year, the Department identified 108 occupied territories, an increase from 91 occupied territories in 2006, and a continued linear increase in the number of occupied territories since 1990. In 2012 as a response to state down-listing of the peregrine, the Washington Forest Practices Board approved the removal of peregrine falcon critical habitat from Forest Practice Rules (WAC 222-16-080). WDFW will continue to recommend site specific management plans associated with nesting peregrines when appropriate to avoid or reduce disturbance.

The species no longer meets the definition of a state sensitive species under Washington law, which is described as “*vulnerable or declining and is likely to become endangered or threatened in a significant portion of its range within the state without cooperative management or removal of threats*” (WAC 232-12-297). WDFW therefore recommends that peregrine falcon be delisted at the state level in Washington. The species will remain classified as “protected wildlife” under state law (WAC 232-12-011) and will continue to be protected under the federal Migratory Bird Treaty Act.

## INTRODUCTION

The peregrine falcon (*Falco peregrinus*) (Figure 1) is divided taxonomically into three subspecies in North America, two of which breed in Washington State, the Peale's peregrine (*F. p. pealei*) and American peregrine (*F. p. anatum*) and the third, the Arctic peregrine (*F. p. tundrius*) occurring as a migrant or rare winter resident (Varland et al. 2012). The Peale's peregrine falcon occurs in coastal regions of the state, primarily along the outer coast, northern coast of the Olympic Peninsula and the San Juan Islands but always within a half mile of salt water. The American peregrine falcon breeds in the Cascade Mountains, the San Juan Islands and the major cities of the Puget Sound basin. They also occur across eastern Washington. Some peregrines breeding along the outer coast and islands of Puget Sound south to central Oregon may be intergrades between the two subspecies (Sheppard 1983, Brown et al. 2007; J. Pagel, pers. comm.). The peregrine was federally listed as endangered in 1970 after dramatic declines following the widespread use of DDT in the 1940's and 50's. In 1980, the peregrine was listed as endangered in Washington when only five pairs could be found nesting. Nationally, restrictions on DDT use combined with releases of young American peregrines to the wild facilitated population recovery (Enderson et al. 1995, White et al. 2002) and the Arctic peregrine falcon and American peregrine falcon were removed from the federal endangered species list in 1994 and 1999, respectively (Mesta 1999). The restriction on DDT use was the primary factor in the eventual recovery of peregrines in Washington. Releases of 145 young peregrines from 1982-1997 in the Cascade Mountains, Columbia Gorge, and Columbia Basin may have contributed to the eventual establishment of some nesting pairs in these areas (Hayes and Buchanan 2002). In Washington, the population increased to 72 occupied territories by 2001, and in 2002 the peregrine was down-listed to a state sensitive species.



Figure 1. Peregrine falcon.

## SPECIES BACKGROUND

**Distribution-breeding.** Peregrines can now be found nesting throughout much of the state (Figure 2). Peregrine falcon nesting is dependent upon availability of abundant prey in proximity to adequate nesting sites, usually near large water bodies (Ratcliffe 1993, White et al. 2002).




Figure 2. Distribution of peregrine falcon nesting territories in Washington, 2016.

The greatest numbers of nesting sites in the state occur in the San Juan Islands, the lowlands of northern Puget Sound, particularly in the cities, and along the outer northern coast. In these regions, peregrines usually nest on islands, “sea stacks”, or shoreline cliffs associated with seabird colonies, waterfowl concentrations, and other prey species. In this region, peregrines also nest in the urban areas of Seattle and Tacoma of central Puget Sound. Lower numbers occur along the forested slopes of the Cascade Mountains and in the Columbia River Basin, where peregrines nest on cliffs that are typically in close proximity to large lakes, or overlook river valleys such as Columbia and Snake Rivers.

***Distribution-winter.*** Western Washington is noted for its high density of wintering peregrines (Anderson and Herman 2005). The mild maritime climate and extensive habitat supports high densities of prey, including shorebirds and waterfowl, rock pigeons (*Columba livia*), and European starlings (*Sturnus vulgaris*). Peregrine wintering areas in western Washington include Grays Harbor, Willapa Bay, the estuaries of Puget Sound, the Columbia River estuary, the outer coastal beaches, low-lying agricultural and pasture lands, the Columbia Gorge, and many urban areas (Anderson and Herman 2005). Peale’s and American peregrines both are found in these habitats throughout spring and fall migration as well as in winter (Anderson and Herman 2005).

***Migration.*** Evidence of a west coast peregrine migration was first described by Anderson et al. (1988). The arctic subspecies was formerly considered an uncommon migrant in the region, but several recent records from British Columbia, Washington, Oregon, and California, demonstrated that they do migrate and winter in the region. In addition, Varland et al. (2012) later described an immature female overwintering on the Long Beach Peninsula in 2000-2001 and another at Ocean Shores in 2010 confirming earlier reports ranging from British Columbia to California that Arctic peregrines are a rare winter resident in Washington (Varland et al. 2008a, Varland et al. 2012). In eastern Washington, the peregrine is now also found in widely scattered localities in open habitats (the channeled scablands, agricultural areas, etc.), but had been considered rare in this area during winter in the late 1990s (Anderson and Herman 2005). Banding data from Washington falcons indicate that at least some resident adult American peregrines generally remain near their nests throughout the year. However, locally produced juveniles may wander widely (C.M. Anderson, unpubl. data). First year banded birds from western Washington have been observed as far south as Los Angeles, California north to Vancouver, British Columbia, and east to Alberta, Canada.

***Breeding habitat requirements.*** The presence of a prominent cliff, tall building, or steel bridge is the most common physiographic characteristic of peregrine nesting territories. Cliffs and tall buildings function as both nesting and perching sites and provide unobstructed views of the surrounding landscape. A successful nest site also requires the presence of ledges or potholes that are essentially inaccessible to mammalian predators, provide protection from the elements, and are protected from heavy rain (Campbell et al. 1990, Johnsgard 1990). A source of open water, such as a river, lake, marsh, or marine waters, is typically found in close proximity to the nest site. The primary advantage of an open body of water is that it provides a featureless hunting area where small terrestrial birds have no cover and are thereby more easily captured. However, peregrines will nest at locations other than cliff sites, such as at the apex of steep,

grass-covered slopes, rock quarries, trees, on the ground, and man-made structures such as bridges, tall buildings, smoke stacks, and cooling towers in urban areas (White et al. 2002).

**Winter habitat requirements.** Habitats used by peregrines during the non-breeding season usually support high densities of shorebirds, waterfowl and other small- to medium-sized birds (Anderson and Herman 2005). Coastal and estuarine that are used include beaches, tidal flats, islands, and marshes. Human-altered habitats and environs include agricultural fields (particularly when flooded), airports, and cities where rock pigeons and European starlings are abundant (White et al. 2002). Roost sites are also an important element of wintering habitat. The first radio telemetry study on peregrines conducted in Washington (Anderson and DeBruyn 1979) discovered that adult female peregrines wintering on the Samish Flats (Skagit County) showed strong fidelity to their roost sites and used them continuously during the winter. Two of these tagged peregrines used different nearby offshore islands as their roost sites. During another wintering peregrine telemetry study on the Lummi Flats in Whatcom County (Anderson et al. 1984) a tagged adult female flew from Sandy Point to Orcas Island each night, a distance of 8.4 miles over open-ocean. Clearly, islands offshore of mainland foraging areas are important as winter roost sites. Dobler (1993) later reported another peregrine flying 15 miles to a roost.

**Diet and foraging.** Peregrine falcons prey on a variety of birds found near cliffs and aquatic features in the vicinity of eyries. During migration and at wintering sites their prey is usually captured near large bodies of water. Studies of Peale's peregrine food habits on Tatoosh Island (Paine et al. 1990), located off the northwestern tip of the Olympic Peninsula, indicated that many species were taken, but Cassin's (*Ptychoramphus aleuticus*) and rhinoceros auklets (*Cerorhinca monocerata*), were by far the most common prey encountered. This was despite higher densities of common murre (*Uria aalge*), and glaucous-winged gulls (*Larus glaucescens*), presumably less preferred because of their larger size. Smaller gull species, such as ring-billed (*L. delawarensis*) and mew gulls (*L. canus*), are frequently taken by peregrines on the outer coast and at inland sites in central Washington along the Columbia and Snake Rivers (Dobler 1993). Other species commonly taken by peregrines in Washington include various species of waterfowl, shorebirds, swallows, and swifts. American robins (*Turdus migratorius*), European starlings, rock pigeons, and cedar waxwings (*Bombycilla cedorum*) were the most common prey of peregrines breeding in the San Juan Islands (Anderson 1995, 1996, 1997). These same species constitute the main diet of urban nesting peregrines in Seattle, Tacoma, and Everett sites as well (Ed Deal, pers. comm.). Cade et al. (1996) also lists these species as common prey in cities across North America, and starlings and rock pigeons are common near peregrine nest sites in the Columbia Basin.

**Home range and movements.** Home ranges of peregrines during the breeding season can be expansive and large size seems closely dependent on distant foraging sorties from eyriess. In Colorado, the largest home ranges averaged 450 mi<sup>2</sup> (1,251 km<sup>2</sup>) for three females while those of two males averaged 405 mi<sup>2</sup> (1,126 km<sup>2</sup>) (Enderson and Craig 1997); hunting flights within these home ranges extended as far as 12-26 mi (20-43 km) from the eyrie. The home ranges of two females in the United Kingdom were 8.3 mi<sup>2</sup> (23 km<sup>2</sup>) and 42.1 mi<sup>2</sup> (117 km<sup>2</sup>) and a hunting female was observed 11 mi (18 km) from the nest (Mearns 1985). On Cape Peninsula, South

Africa, two female and two male peregrines had an average home range size of 44.3 mi<sup>2</sup> (123.0 km<sup>2</sup>) (Jenkins and Benn 1998); hunting excursions from nest sites averaged 10 mi (16.7 km) per flight. Two other studies reported average hunting excursions of 12 and 16 mi (20 and 27 km), a potential indication of substantial home range size, but they did not determine home range size (Porter et al. 1973, Kumari 1974 cited in Mearns 1985).

During winter, peregrines can range over extensive areas when hunting prey. In Washington, in the vicinity of Sequim, where three birds were monitored for most of a single winter, home range size was 23.7 mi<sup>2</sup> (65.8 km<sup>2</sup>) for an immature female and 30.9 mi<sup>2</sup> (85.7 km<sup>2</sup>) for an immature male (Dobler 1993). Core areas (areas of concentrated activity) were 4.9 mi<sup>2</sup> (13.5 km<sup>2</sup>) and 9.1 mi<sup>2</sup> (25.3 km<sup>2</sup>) for the female and male, respectively. At Grays Harbor, an immature male peregrine had a home range of 28 mi<sup>2</sup> (78 km<sup>2</sup>) and core area of 7.1 mi<sup>2</sup> (19.8 km<sup>2</sup>) during a single winter (Dobler and Spencer 1989).

**Mortality.** Mortality factors represented for peregrines in Washington include: unhatched eggs, adults accidentally displacing young off the breeding ledge (1), nestling dying on nest ledge during the first week of life (multiple), nestling brain abscess (1), hitting windows in urban environments (common), collisions with vehicles (multiple), electrocution (1), aircraft bird strikes (2), striking wires (2), collision with powerlines (1), broken wing (multiple), ensnared in fishing line (1), nestling mycoplasma meningitis (1), unknown mammalian predation (raccoon/coyote) at nest sites (multiple), killed by golden eagle while defending eggs (1), territorial battle with other female (1), run over by vehicles, shot (1), on ground after fledging in cities (common), infected by pigeon sourced *Trichomonas gallinae* (multiple), trapped in hollow bridge beam (1), trapped in window well (1), eggs killed by flooded substrates from rain (multiple), premature fledging (2), eggs broken by improper substrate (multiple), drowning after fledging (multiple), and avian influenza (1).

**Juvenile and adult survival.** Peregrines may live up to 20 years (White et al. 2002). In Colorado, Craig et al. (2004) estimated survival rates for three age classes of peregrines: 54% for 0-1 year olds, 67% for 1-2 year olds, and 80% for birds older than 2 years of age. In California, first year survival was estimated as 38%, second year survival as 86% and adult survival (>2 years old) as 85% (Kauffman et al. 2003). For peregrines using coastal beach habitat along the Washington coast, Varland et al. (2008b) estimated an annual apparent survival rate of 59.7%, which included both juvenile (<1 yr old) and adult (≥1 yr old) peregrines. Wootton and Bell (1992) modeled the peregrine falcon population in California and determined that adult survivorship was the most important factor affecting population growth. With the continued growth of the peregrine population in Washington, we surmise that adult survivorship is not limiting the population, and survivorship and productivity are sufficient to support an increasing population.

## POPULATIONS AND HABITAT STATUS

### North America

The peregrine falcon is found throughout North America from the Canadian arctic to Mexico. It occurs wherever suitable nest sites and prey populations are found. While many historical sites in the eastern United States have yet to be recolonized, the eastern population has adapted to the use of urban areas and artificial structures and the numbers of nesting birds in most states are believed to rival historical estimates (Carter et al. 2003, Katzner et al 2012, Faccio et al. 2013, Gahbauer et al. 2015, Watts et al. 2015), although the estimates may be biased low. In the western United States, most historical (pre-1970-1975) peregrine eyries have been reoccupied across their range, and the population now is recovered beyond documented historical levels. This is in part due to the addition of urban and artificial nest sites, artificial habitat alterations (e.g., reservoirs), and increased availability of alternative prey species (Bond 1946, Enderson et al. 2012, Sharpe 2014, Barnes et al. 2015). Certainly the introduction of the rock pigeon and European starling has also had a major effect on increasing the population. Historical baseline estimates of peregrine populations for North America are poorly known due to lack of systematic surveys (Enderson et al. 1995), however, based on more complete records and new surveys, the pre-decline population has been estimated as: 7,000-10,000 nesting territories with an 80-90% occupancy rate (Kiff 1988), 7,300 pairs (Enderson et al. 1995), and 10,600-12,000 pairs (Cade 2003). By the mid-1990s an estimated 7,169 pairs were estimated breeding in North America (Enderson et al. 1995), an estimated 8,000-10,000 pairs by the late 1990s (White et al. 2002:32) and the most recent upper population estimate is 10,368 breeding pairs based on an analysis by the USFWS to determine a harvest quota for falconry (USFWS 2008).

Territory occupancy, nest success and productivity are indices of the overall health of peregrine populations (USFWS 2003). The breeding “territory,” or “breeding site,” refers to an area containing, or historically contained, one or more nest ledges where a peregrine falcon pair have been observed, at least once, in reproductive activity, including nest defense, courtship flights, nest building and repair, copulation, incubation, egg-laying, and/or successful breeding and fledging of young (Postupalsky 1974, Steenhof and Newton 2007). The rate of occupancy is defined as the percentage of the total known territories where activity patterns indicate the presence of a mated, territorial pair of potential breeders. These activity patterns include: two adults or an adult/subadult pair associated with a nest, incubation behavior, presence of eggs, or presence of young (Postupalsky 1974). Nest success is defined as the percentage of occupied territories (for which the outcome of nesting is known) which produce one or more young to an advanced stage of development (Postupalsky 1974, USFWS 2003, Steenhof and Newton 2007). Productivity is another measure of reproductive success and is defined as the number of young (fledging or advanced age of development) per occupied nest (Postupalsky 1974, Steenhof and Newton 2007). These three indices of population health were low between 1950 and 1980 when populations declined severely but rebounded during population recovery (Cade et al. 1988, Enderson et al. 1995, White et al. 2002).

The federal post-delisting Monitoring Plan for the American peregrine falcon populations was designed to detect a significant decline in territory occupancy, nest success, or productivity in six recovery regions across the U.S. Data was to be collected at a random sample of peregrine territories for five sampling period, at three year intervals, beginning in 2003 and ending in 2015.

Therefore, to meet the mandate of the USFWS to monitor peregrines for not less than five years after delisting, the plan called for continued monitoring in 2003, 2006, 2009, 2012, and 2015 (USFWS 2003). Results of federal post-delisting monitoring have been made available for the 2003 survey year (Green et al. 2006). In 2003, territory occupancy across six recovery regions in the United States ranged from 67% to 98% and averaged 87% for all regions combined. Nest success across the regions ranged from 56% to 90% and averaged 71%, and productivity ranged from 1.27 to 2.32 fledglings/occupied territory with an average productivity of 1.64 fledglings/occupied territory (Green et al. 2006). These values are consistent with stable or expanding populations (Craig et al. 2004, Enderson et al. 2012).

## Washington

Following the state down-listing of the peregrine falcon in February of 2002, WDFW conducted a comprehensive survey of peregrines in the 2002 breeding season that included both the Peale's peregrine and American peregrine nest sites. In 2003, the USFWS implemented a federal post delisting Monitoring Plan (USFWS 2003) nationwide to monitor the status of the American peregrine following federal delisting. Twenty five nesting territories were randomly selected in Washington by USFWS for the national monitoring project (hereafter the "USFWS" sample) in 2003, 2006, 2009, and 2012. The minimum USFWS sample in Washington (n=25) was designed to detect declines in regional peregrine populations. To provide more complete information on the status of Washington's American peregrine population, the Department surveyed, in addition to the USFWS sample sites, as many additional nest sites as possible in 2003, 2006, and 2009. In 2012, WDFW only surveyed the USFWS sample sites.

**Occupancy rate.** Overall, the rate of occupancy of eyries in Washington has been high. Over the 10-year period from 1992-2001, the occupancy rate averaged 79%, and even higher for the five-year period from 1997-2001 (82%), with some regional variation: Outer Coast (84%), Puget Sound (89%), Upland Forested (74%), and Arid (72%) (Hayes and Buchanan 2002). Similarly, high occupancy rates continued to be observed during comprehensive surveys in 2006 (79%) and 2009 (82%), and also during the federal post-delisting monitoring of the species that began in 2003 at a random sample of nest sites. These sites were monitored at three year intervals (Appendix A). The overall high occupancy rates compare well with that of stable populations (Herbert and Herbert 1969, Rice 1969, Craig et al. 2004, Enderson et al. 2012).

**Nest success.** A nesting pair is considered successful if it raises at least one young to the minimum acceptable age for the species, which for peregrines is 28 days (USFWS 2003). However, in Washington, a nesting pair was considered successful if young were observed in the nest, regardless of age. This produces inflated success rates when compared to studies using the standard definition. Even taking this into account, we believe actual nest success has been high. Over the 10-year period from 1992-2001, nest success averaged 62%. During the five-year period from 1997-2001, nest success averaged 64% with some regional variation: Outer Coast (57%), Puget Sound (65%), Upland Forested (76%), and Arid (69%) (Hayes and Buchanan 2002). Comparable nest success rates were observed during recent comprehensive surveys in 2006 (68%) but nest success was lower in 2009 (37%). During the federal post-delisting

monitoring of the species that began in 2003, nest success was  $\geq 50\%$  at a random sample of nest sites (Appendix A). Nest success rates observed for other recovering populations include an average of 73% (1984-1996) for a cliff-nesting population in northern New England and New York (Corser et al. 1999), 62% (1991-1995) for a population in the Midwest (Tordoff and Redig 1997), and 70-83% (2005-2009) for populations in Colorado, Montana, and Wyoming (Enderson et al. 2012).

**Productivity.** In Washington over the five-year period from 1997-2001, productivity averaged 1.53 young/occupied territory. Productivity at six urban nest sites averaged 1.65 young/occupied territory over that same time period (Hayes and Buchanan 2002). More recently, during comprehensive surveys in 2006 and 2009, productivity was 2.09 and 1.79 young/occupied territory, respectively. In the federal post-delisting monitoring of the species that began in 2003 productivity was  $>1.00$  young/occupied territory at a random sample of nest sites (Appendix A). Overall productivity rates compare well with increasing peregrine populations in the eastern United States (Corser et al. 1999), the Midwest (Tordoff and Redig 1997), and Colorado, Montanan, and Wyoming (Enderson et al. 2012).

WDFW last completed comprehensive surveys of peregrine falcon territories in 2009. In that year, the Department identified 108 occupied territories, an increase from 91 occupied territories in 2006, and a continued

linear increase in the number of occupied territories since 1990 (WDFW, WSDM database; Figure 3). Nest success in 2009 was lower than the long-term average (60%), with less than 40% of occupied territories producing at least one nestling to fledging age in 2009. However, surveys in 2012 at a random sample of sites (n=25) found an 84% occupancy rate, 76%




Figure 3. Trend in occupied territories and nesting success in Washington, 1990-2009.

nest success rate and 1.81 fledglings/occupied territory (Appendix A). The long-term trends of increasing numbers of nesting territories, high occupancy rates, and moderate fledging rates are consistent with a stable or increasing population. Historical estimates of territories in Washington (Bond 1946) vary and likely underestimate the actual number of territories, but only nine territories were identified from a 1980 survey, although Herman (*in* Porter and White 1977) reviewed the existing literature and other sources and estimated as many as 25 historical territories. Although surveys were not conducted in 2016, the Department has 181 nesting territories documented in its peregrine falcon nesting territory database (WSDM database); applying the 2009 occupancy rate of 82% to the 181 known territories, we estimate 148

territories currently occupied statewide. The estimated 148 occupied territories in 2016 far exceeds the 25 historical sites documented during the pre-DDT era and the minimum of 30 pairs established for Washington as part of the federal delisting criteria for the Pacific Coast American peregrine population (USFWS 1982).

***Habitat-nesting.*** The number of available natural nesting sites has likely changed little from the number available historically. There have been some habitat changes at a few sites that have made cliffs unattractive or unavailable to nesting pairs, while a few may have been created by rock quarries, logging, and fire (Bell 2001, Hayes and Buchanan 2002). Many pairs have become established on human-made structures, such as buildings, and bridges (Cade and Bird 1990).

***Habitat-foraging.*** The net effect of human modification on peregrine foraging habitat in Washington is difficult to determine because the peregrine is a generalist predator. As human populations have increased, peregrine foraging habitats (e.g., wetlands, marine waters, coastal barrier islands, and river valleys) have been destroyed or degraded. During the same period, humans have created opportunities in urban areas where there are alternative artificial nest sites and abundant pigeon and starling populations. The wide variety of habitat types and prey species used by the peregrine and the increasing population trend suggest that foraging habitat and prey availability are not limiting the population in Washington.

## **FACTORS AFFECTING CONTINUED EXISTENCE**

***Adequacy of regulatory mechanisms.*** Peregrines are protected by the Federal Migratory Bird Treaty Act which prohibits take, possession, import, export, transport, selling, purchase, barter, or offering for sale, purchase or barter, any migratory bird, their eggs, parts, nests, except as authorized under a valid permit. There are no federal laws that specifically protect the habitat of this species. However, loss of habitat was not identified as a limiting factor in peregrine recovery (Mesta 1999) and was not a factor identified as contributing to the species' listing.

In 2002, the Washington Fish and Wildlife Commission reclassified the peregrine falcon as a state sensitive species. A sensitive species is defined in the Washington Administrative Code (WAC 232-12-297, Section 2.6) as a species “*native to the state of Washington that is vulnerable or declining and is likely to become endangered or threatened in a significant portion of its range within the state without cooperative management or removal of threats.*” Regulatory protection of peregrine falcons as a state sensitive species is the same as that afforded by federal law under the Migratory Bird Treaty Act. In addition, the peregrine is designated a “priority species” and cliff-nesting habitat is identified as a “priority habitat feature” under the Priority Habitat and Species (PHS) Program. The PHS Program provides important wildlife and habitat information and management recommendations to agencies, landowners, municipalities, and consultants for land use planning (WDFW 2008, Larsen et al. 2004). State Forest Practices Rules identify critical habitat for endangered and threatened species, and in 2012 the Washington Forest Practices Board approved the removal of peregrine falcon Critical Habitat from State forest practices rules (WAC 222-16-080).

***Falconry.*** Washington State has received approval by the USFWS to regulate and issue permits for the take of nestlings and Washington fledglings while adhering to the Code of Federal Regulations (50 CFR Section 21.29). The USFWS has set a maximum allowable harvest of up to 5% of the annual fledgling production. States are under no obligation to issue permits up to the maximum sustainable harvest. In 2008, the USFWS published the Final Environmental Assessment and Management Plan on Take of Migrant Peregrine Falcons from the Wild for Use in Falconry, and Reallocation of Nestling/Fledgling Take (USFWS 2008). The Service allocated “take” of up to 116 wild first-year peregrine falcons, including 41 in Alaska and 75 apportioned among states west of 100° west longitude, with capture period limited to between May 1<sup>st</sup> and August 31<sup>st</sup>. In 2009, the Pacific Flyway Council approved the Pacific Flyway Nongame Migratory Bird Technical Committee’s recommendation to follow authorizations of the USFWS. Between 2009 and 2014, nine of the eleven Pacific Flyway states (excluding Alaska) authorized an average take of 63 peregrines per year (California and Nevada do not allow any take). An average of 19 peregrines were removed from the wild per year, well below the authorized limit of 75. The Pacific Flyway allocation has not been re-evaluated since 2008, and therefore the allocation remains conservative, and the actual take of first-year peregrines has never achieved the allocation and likely has no population impact. In 2015, WDFW allocated 11 nestlings or fledglings for harvest, and falconers filled seven permits consisting of six nestlings and one fledgling. An average of 4.6 nestlings and fledglings were taken in Washington per year from 2011 through 2015.

***Contaminants.*** An important regulatory mechanism protecting peregrine falcons is the requirement that pesticides be registered with the Environmental Protection Agency (EPA). Under the authority of the Federal Insecticide, Fungicide and Rodenticide Act, the EPA requires environmental testing of all new pesticides. Testing the effects of pesticides on representative wildlife species prior to registration is required, although this testing does not include evaluation of the combined effects of multiple legal pesticides which may have detrimental effects.

DDT and its metabolites were the primary cause of peregrine falcon decline across North America. Despite restriction on the use of DDT in the U.S. and Canada in 1972, peregrine falcon populations were slow to recover. DDT and its metabolites are persistent in the environment with a half-life up to 57 years (Cooke and Stringer 1982). As peregrines are known to accumulate contaminants in wintering areas (Henny et al. 1982), or by consumption of prey that overwinter in those areas (Fyfe et al. 1990), the continued use of DDT south of the U.S. border was an ongoing concern and was addressed in Mexico with the implementation of the North American Agreement for Environmental Cooperation, signed in 1997 by the United States, Canada and Mexico. Mexico met the obligations of the agreement by the year 2000, and no longer produces or permits use of DDT. However, DDT use south of U.S. was much less significant to peregrines breeding in the Pacific Northwest that don’t migrate that far south. In addition, prey species returning from DDT contaminated areas south of the U.S. border arrive on the breeding grounds in the Pacific Northwest after peregrines have already started nesting (J. Pagel, pers. comm.). In Washington, a surface water monitoring program conducted by the Washington State Department of Agriculture found elevated DDT or its metabolites in two

streams within seven monitored watersheds (Tuttle and Castro 2015). River, stream, and estuary sediments in many areas are likely still contaminated and may continue to pose an unknown level of environmental risk, although bald eagles, osprey, and peregrine falcons, all impacted by DDT contaminants, continue to show population increases and healthy productivity.

A new environmental contaminant issue that arose in recent years was the detection of widely used flame retardant chemicals, polybrominated diphenyl ethers (PBDEs), in peregrine eggs in Spain, Canada, and California (Park et al. 2009, Guerra et al. 2012). The compounds are classified as persistent, bioaccumulative, and toxic (PBT), and may have neurological and endocrine effects that at high levels could affect reproduction. To date, it has not been demonstrated that these compounds are impacting reproduction of peregrine falcons in the wild. In addition, since 2000, the European Union, the U.S. Environmental Protection Agency, and many states, including Washington, have placed restrictions and bans on the use of PBDEs. In Washington, manufacturers in 2014 voluntarily discontinued production of two of the three most commonly used forms of PBDEs, and the major manufacturers of the third form of PBDE agreed to voluntarily discontinue its use by the end of 2013 (Washington State Department of Health; <http://www.doh.wa.gov/CommunityandEnvironment/Contaminants/PBDEs>). In California, detection of the compounds declined by half between 2002-2012 in sport fish, and by 65-95% in bivalves, bird eggs, and pregnant women (Sutton et al. 2015, Zota et al. 2013).

Organophosphates and neonicotinoid insecticides, PCBs, heavy metals, avicides and oil are other chemicals that have the potential to impact peregrines. In addition, it has long been recognized that combinations of various compounds may have far more deleterious effects on wildlife than the individual chemicals themselves. The significance of these synergistic effects is impossible to quantify at present because they are species-specific and also vary as a function of the types and amounts of chemicals present in animal tissues. Needless to say, widespread presence of harmful chemicals or an oil spill off the Washington coast that decimates prey populations could have significant local or regional impacts on the peregrine population, but current regulatory mechanisms are in place to limit environmental exposure to chemical pollutants. Broad scale use of neonicotinoid insecticides in agricultural landscapes could affect local peregrine populations by reducing their local prey base through direct toxic effects by consumption of treated seeds or depletion of insect food resources (Goulson 2013, Mineau and Palmer 2013, Hallmann et al. 2014).

***Climate change.*** Models of climate change indicate changes in precipitation levels and temperature throughout the Pacific Northwest. Although the models vary in their specific predictions, all of them indicate that substantial changes will occur. As a consequence, it appears likely that such changes will alter conditions in the marine and other aquatic environments important to peregrine falcons. In the marine environment, future climate projections for sea surface temperature and upwelling intensity, based on a regional climate model for the California Current Ecosystem, have forecast accelerated declines of some Cassin's auklet populations (Wolf et al. 2010). Declines in seabird productivity or abundance may impact coastal nesting peregrines, but to what degree is difficult to predict. Inland, late season storms and increased

drought and fire could contribute to habitat loss of prey could negatively affect reproductive success of peregrines.

**Other factors.** A highly pathogenic avian influenza (HPAI) outbreak in the winter of 2014-2015 was responsible for the deaths of three captive gyrfalcons (*Falco rusticolus*) (Ip et al. 2015) and a wild peregrine falcon ([https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/animal-disease-information/avian-influenza-disease/ct\\_avian\\_influenza\\_disease](https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/animal-disease-information/avian-influenza-disease/ct_avian_influenza_disease)) in Washington and three captive peregrine falcons used for falconry and two privately owned gyrfalcons in Idaho (IDHW 2015). It is believed these birds became infected after being feeding on infected wild-caught waterfowl. Captive falcons are highly susceptible to the virus (Lierz et al. 2007), however, large-scale die-offs of wild raptors testing positive for the virus have not been detected, either due to low densities, difficulty in finding carcasses, or some degree of immunity in wild populations. At this time, it does not appear that HPAI is having a population-level effect on peregrines or other raptors.

Avian trichomoniasis (*Trichomonas gallinae*) is an emerging issue with golden eagles in the Snake River Birds of Prey Conservation Area (J. Watson pers. comm.) and is known to have killed several wild peregrines in Washington, particularly nestlings (C. M. Anderson, pers. comm.). However, the northern goshawk (*Accipiter gentilis*), a cosmopolitan species similar in distribution to the peregrine, was shown to have a high incidence of exposure but a low rate of pathological affects, perhaps due to evolutionary adaptations to the parasite (Krone et al. 2005), and the same may apply for the peregrine falcon. Other diseases, human disturbance through recreational (rock climbing, hiking, beach walking, etc.) or industrial (blasting, logging, etc.) activities, illegal shooting, habitat loss, and inbreeding depression are some of the other factors with the potential to impact peregrine falcons, although currently none of these issues are known to be limiting North American peregrine populations.

## MANAGEMENT ACTIVITIES

WDFW has developed site specific management recommendations that mainly involve variable buffers to activities around peregrine nest sites, but also include wetland protection and pesticide application limitations around nest sites and wintering areas. In addition, WDFW is consulted on site-specific plans to avoid or reduce disturbance of nesting peregrines, primarily related to recreational activities such as rock climbing and hiking cliffs above eyries. WDFW permits a small number of peregrine falcons (nestlings or fledglings) to be taken each year for falconry purposes. The level of allowed take is currently at 12 individuals per year and is based on federal regulations for the take of this species. Washington state does not allow non-resident take of wild peregrines.

## CONCLUSIONS AND RECOMMENDATION

When the peregrine falcon was federally listed in 1970, the primary factor contributing to its status under the Endangered Species Act was dramatic population declines due to low productivity caused by the accumulation of PBT compounds, specifically DDT and its

metabolites. DDT use was restricted in the U.S. in 1972 (37 FR 13369), with a single exception made in 1974 of use for Douglas-fir tussock moth (*Orgyia pseudotsugata*) control in the Blue Mountains of Washington, Oregon, and Idaho (Henny 1977). Following the banning of DDT, peregrine populations slowly recovered in response to 1) regulatory actions on DDT and other PBT compounds, 2) natural productivity of known and unknown nest sites, and 3) nest site protection. PBT compounds probably remain the greatest potential threat to peregrine populations worldwide, but regulatory mechanisms like those through the EPA and the Stockholm Convention are likely to mitigate that threat. Climate change may be the next major threat and could have the greatest impact on coastal peregrines (23%-42% of the Washington population) that depend on seabirds, which in turn depend on forage fish and ocean conditions. There is no way to predict how peregrine populations would respond to declines in seabird densities, as there are likely to be both positive and negative climate change effects on alternate prey species. HPAI is another emerging issue that has the potential to impact peregrine populations. Falconry birds have been especially sensitive to infection, which indicates the potential for infection in the wild, but so far it does not appear to have had a detrimental effect on wild populations of peregrines. Although a number of threats remain, peregrine falcon numbers in Washington State have been increasing in a linear fashion for over two decades with no indications of leveling off. Peregrine falcons breeding in Washington have likely recovered well beyond to pre-DDT levels, and the population continues to increase. WDFW will continue to recommend site specific management plans associated with nesting peregrines when appropriate to avoid or reduce disturbance.

The species no longer meets the definition of a state sensitive species under Washington law, which is described as “*..vulnerable or declining and is likely to become endangered or threatened in a significant portion of its range within the state without cooperative management or removal of threats*” (WAC 232-12-297). WDFW therefore recommends that the peregrine falcon be delisted at the state level in Washington. The species will remain classified as “protected wildlife” under state law (WAC 232-12-011) and will continue to be protected under the federal Migratory Bird Treaty Act.

## LITERATURE CITED

- Anderson, C. M. 1995. The San Juan peregrine project: a report on the 1994 and 1995 breeding seasons. Unpubl. Rep. Falcon Research Group, Bow, Washington.
- Anderson, C. M. 1996. The San Juan peregrine project: a report on the 1996 breeding season. Unpubl. Rep. Falcon Research Group, Bow, Washington.
- Anderson, C. M. 1997. The San Juan Islands peregrine project: a report on the 1997 breeding season. Unpubl. Rep. Falcon Research Group, Bow, Washington.
- Anderson, C. M. and P. DeBruyn. 1979. Behavior and ecology of peregrine falcons wintering on the Samish Flats, Washington. A report on the 1979 field season. Washington Department of Game, Threatened Species and Management, Olympia, Washington.
- Anderson, C. M. and S. G. Herman. 2005. Peregrine falcon species account in T. R. Wahl, B. Tweit, and S. G. Mlodinow, editors. The Birds of Washington. Oregon State University Press, Corvallis.
- Anderson, C. M., R. K. Knapp, and J. K. Fackler. 1984. The behavior and ecology of fall peregrine falcons at Lummi Bay and vicinity, Whatcom county, Washington. Unpubl. Rep.
- Anderson, C. M., D.G. Roseneau, B.J. Walton, and P. J. Bente. 1998. New evidence of a peregrine migration on the West Coast of North America. Pages 507-516 in Cade, T.J., J. H. Enderson, C. J. Thelander, and C. M. White, editors. 1988. Peregrine Falcon Populations: their management and recovery. The Peregrine Fund, Inc., Boise, Idaho.
- Barnes, J.G., R.D. Haley, D.B. Thompson, and J.R. Jaeger. 2015. Attributes of a breeding population of peregrine falcons associated with reservoirs on the Colorado River. Journal of Raptor Research 49(3):269-280.
- Bell, S.W. 2001. Reclaiming abandoned rock quarries for peregrine falcon aeries in the lowlands of western Washington. M.S. problem statement. Washington State University, Pullman.
- Bond, R.M. 1946. The peregrine population of western North America. The Condor 48(3):101-116.
- Brown, J. W., P. J. Van Coeverden de Groot, T. P. Birt, G. Seutin, P. T. Boag, and V. L. Friesen. 2007. Appraisal of the consequences of the DDT-induced bottleneck on the level and geographic distribution of neutral genetic variation in Canadian peregrine falcons, *Falco peregrinus*. Molecular Ecology 16:27-343.
- Cade, T.J. 2003. Life history traits of the peregrine in relation to recovery. Pages 2-11 in T.J. Cade, W. Burnham, and P. Burnham, editors. Return of the Peregrine: a North American Saga of Tenacity and Teamwork The Peregrine Fund, Boise, Idaho.
- Cade, T. J. and D. M. Bird. 1990. Peregrine falcons, *Falco peregrinus*, nesting in an urban environment: a review. Canadian Field-Naturalist 104:209-218.
- Cade, T.J., J. H. Enderson, C. J. Thelander, and C. M. White, editors. 1988. Peregrine Falcon Populations: their management and recovery. The Peregrine Fund, Inc., Boise, Idaho.
- Cade T. J., A. M. Martell, P. Redig, G. Septon, and H. Tordoff. 1996. Peregrine falcons in urban North America. Pages 3-13 in D. Bird, D. Varland, and J.J. Negro. Raptors in human landscapes: adaptations to built and cultivated environments.
- Campbell, R. W., N. K. Dawe, I. McTaggart-Cowan, J. M. Cooper, G. W. Kaiser, and M. C. E. McNall. 1990. The Birds of British Columbia. University of British Columbia Press, Vancouver, British Columbia, Canada.
- Carter, K.M., M.J. Lacki, and M.R.Dzialak. 2003. Food habits of peregrine falcons in Kentucky. Journal of Raptor Research 37(4):344-349.
- Cooke, B.K. and A. Stringer. 1982. Distribution and breakdown of DDT in orchard soil. Pesticide Science 13:545-551.

- Corser, J. D., M. Amaral, C. J. Martin, and C. C. Rimmer. 1999. Recovery of a cliff-nesting peregrine falcon, *Falco peregrinus*, population in northern New York and New England, 1984-1996. *Canadian Field-Naturalist* 113:472-480.
- Craig, G.R., G.C. White, and J.H. Enderson. 2004. Survival, recruitment, and rate of population change of the peregrine falcon population in Colorado. *Journal of Wildlife Management*, 68(4):1032-1038.
- Dobler, F. C. 1993. Wintering peregrine falcon (*Falco peregrinus*) habitat utilization near Sequim, Washington. *Northwest Science* 67: 231-237.
- Dobler, F. C. and R. D. Spencer. 1989. Wintering peregrine falcon (*Falco peregrinus*) habitat utilization in Grays Harbor, Washington. Pages 71-78 in B-U. Meyburg and R. D. Chancellor, editors. *Raptors in the modern world: proceedings of the III world conference on birds of prey and owls*. WWGBP, Berlin, Germany.
- Enderson, J. H. and G.R. Craig. 1997. Wide ranging by nesting peregrine falcons (*Falco peregrinus*) determined by radiotelemetry. *Journal of Raptor Research* 31:333-338.
- Enderson, J. H., W. Heinrich, L. Kiff, and C. M. White. 1995. Population changes in North American peregrines. *Transactions of the 60<sup>th</sup> North American Wildlife and Natural Resources Conference* 142-161.
- Enderson, J.H., R.J. Oakleaf, R.R. Rogers, and J.S.Sumner. 2012. Nesting performance of peregrine falcons in Colorado, Montana, and Wyoming, 2005-2009. *The Wilson Journal of Ornithology* 124(1):127-132.
- Faccio, S.D., M. Amaral, C.J. Martin, J.D. Lloyd, T.W. French, and A. Tur. 2013. Movement patterns, natal dispersal, and survival of peregrine falcons banded in New England. *Journal of Raptor Research* 47(3):246-261
- Fyfe, R. W., U. Banasch, V. Benavides, N. Hilgert De Benavides, A. Luscombe, and J. Sanchez. 1990. Organochlorine residues in potential prey of peregrine falcons, *Falco peregrinus*, in Latin America. *Canadian Field-Naturalist* 104:285-292.
- Gahbauer, M.A., D.M. Bird, K.E. Clark, T. French, D.W. Brauning, and F.A. McMorris. 2015. Productivity, mortality, and management of urban peregrine falcons in northeastern North America. *Journal of Wildlife Management* 79(1):10-19.
- Goulson, D. 2013. An overview of the environmental risks posed by neonicotinoid insecticides. *Journal of Applied Ecology* 50:977-987.
- Green, M., T. Swem, M. Morin, R. Mesta, M. Klee, K. Hollar, R. Hazelwood, P. Delphey, R. Currie, and M. Aramal. 2006. Monitoring results for breeding American peregrine falcons (*Falco peregrinus anatum*), 2003. *Biological Technical Publication BTP-R1005-2006*. U.S. Department of Interior, Washington, D.C.
- Guerra, P., M. Alae, B. Jimenez, G. Pacepavicius, C. Marvin, G. MacInnis, E. Eljarrat, D. Barcelo, L. Champoux, and K. Fernie. 2012. Emerging and historical brominated flame retardants in peregrine falcon (*Falco peregrinus*) eggs from Canada and Spain. *Environment International* 40:179-186.
- Hallmann, C.A., R.P.B. Foppen, C.A.M. van Turnhout, H.De Kroon, and E. Jongejans. 2014. Declines in insectivorous birds are associated with high neonicotinoid concentrations. *Nature Letter* doi:10.1038/nature13531.
- Hayes, G.E. and J.B. Buchanan. 2002. Washington state status report for the peregrine falcon. Washington Department of Fish and Wildlife, Olympia, WA.
- Henny, C. J. 1977. Birds of prey, DDT, and tussock moths in the Pacific Northwest. *Transactions of the North American Wildlife and Natural Resources Conference* 42:397-411.
- Henny, C. J., L. J. Blus, and R. M. Prouty. 1982. Organochlorine residues and eggshell thinning in Oregon seabird eggs. *Murrelet* 63:15-21.

- Herbert, R. A. and K. G. S. Herbert. 1969. The extirpation of the Hudson River peregrine falcon population. Pages 133-154 in J. J. Hickey, editor. Peregrine falcon populations: their biology and decline. University of Wisconsin Press, Madison, Wisconsin.
- (IDHW) Idaho Department of Health and Welfare. 2015. Avian influenza viruses found in Pacific NW birds: focus on Idaho, public health concerns, and interventions. Disease Bulletin 22(1).
- Ip, H. S., M. K. Torchetti, R. Crespo, P. Kohrs, P. DeBruyn, K. G. Mansfield, T. Bazler, L. Badcoe, B. Bodenstein, V. Shearn-Bochsler, M. L. Killian, J. C. Pedersen, N. Hines, T. Gidlewski, T. Deliberto, and J. M. Sleeman. 2015. Novel Eurasian highly pathogenic avian influenza A H5 viruses in wild birds, Washington, USA, 2014. USDA National Wildlife Research Center – Staff Publications Paper 1736.
- Jenkins, A. R. and G. A. Benn. 1998. Home range size and habitat requirements of peregrine falcons on the Cape Peninsula, South Africa. *Journal of Raptor Research* 32:90-97.
- Johnsgard, P. A. 1990. Hawks, eagles, and falcons of North America: biology and natural history. Smithsonian Institution. Washington, D.C.
- Kauffman, M. J., W. F. Frick, and J. Linthicum. 2003. Estimation of habitat-specific demography and population growth for peregrine falcons in California. *Ecological Applications* 13:1802-1816.
- Katzner, T., J.D. Winton, F.A. McMorris, and D. Brauning. 2012. Dispersal, band encounters, and causes of death in a reintroduced and rapidly growing population of peregrine falcons. *Journal of Raptor Research* 46(1):75-83.
- Krone, O., R. Altenkamp, and N. Kenntner. 2005. Prevalence of *Trichomonas gallinea* in northern goshawks from the Berlin area of northeastern Germany. *Journal of Wildlife Diseases* 41(2):304-309.
- Larsen, E., J. M. Azerrad, N. Nordstrom, editors. 2004. Management recommendations for Washington's priority species, Volume IV: Birds. Washington Department of Fish and Wildlife, Olympia, Washington, USA.
- Lierz, M., H.M. Hafez, R. Klopffleisch, D. Luschow, C. Prusas, J.P. Teifke, M. Rudolf, C. Grund, D. Kalthoff, T. Mettenleiter, M. Beer, and T. Harder. 2007. Protection and virus shedding of falcons vaccinated against highly pathogenic avian influenza A virus (H5N1). *Emerging Infectious Diseases* 13(11):1667-1674.
- Mearns, R. 1985. The hunting ranges of two female peregrines towards the end of a breeding season. *Raptor Research* 19:20-26.
- Mesta, R. 1999. Final rule to remove the American peregrine falcon from the federal list of endangered and threatened wildlife, and to remove the similarity of appearance provision for free-flying peregrines in the conterminous United States. *Federal Register* 64 (164): 46541-46558.
- Mineau, P. and C. Palmer. 2013. Neonicotinoid Insecticides and Birds: the impact of the Nation's most widely used insecticides on birds. American Bird Conservancy.
- Paine, R. T., J. T. Wootton, and P. D. Boersma. 1990. Direct and indirect effects of peregrine falcon predation on seabird abundance. *Auk* 107:1-9.
- Park J.S., A. Holden, V. Chu, M. Kim, A. Rhee, and P. Patel. 2009. Time-trends and congener profiles of PBDEs and PCBs in California peregrine falcons (*Falco peregrinus*). *Environ. Sci. Technol.* 43:8744–51.
- Porter, R. D. and C. M. White. 1977. Status of some rare and lesser known hawks in western United States. Pages 39-45 in R.D. Chancellor, editor. Proceedings of the world conference on birds of prey. International Council for Bird Preservation.
- Porter, R. D. and C. M. White, and R. J. Erwin. 1973. The peregrine falcon in Utah, emphasizing ecology and competition with the prairie falcon. *Brigham Young University Science Bulletin* 18(1).

- Postupalsky, S. 1974. Raptor reproductive success: some problems with methods, criteria, and terminology. Pages 21-31 in F. N. Hamerstrom, B. E. Harrell, and R. R. Olendorff, editors. Proceedings of the conference on raptor conservation techniques. Fort Collins, Colorado.
- Ratcliffe, D. A. 1993. The Peregrine Falcon. Second edition. T& A D Poyser, London, England.
- Rice, J. N. 1969. The decline of the peregrine population in Pennsylvania. Pages 155-163 in J. J. Hickey, editor. Peregrine falcon populations: their biology and decline. University of Wisconsin Press, Madison, Wisconsin.
- Sharpe, P. B. 2014. Peregrine falcon monitoring on the California Channel Islands, California, 2013. Unpublished report prepared by the Institute for Wildlife Studies, Arcata, California for Montrose Settlements Restoration Program.
- Sheppard, J.M. 1983. Proposed reclassification of the peregrine falcons in North America. Federal Register 48:8796-8801.
- Steenhof, K. and I. Newton. 2007. Assessing nesting success and productivity. In: D.M. Bird, and K.L. Bildstein (Eds.), Raptor research and management techniques. Hancock House, Surrey, Canada. pp. 181–192.
- Sutton, R., M.D. Sedlak, D. Yee, J.A. Davis, D. Crane, R. Grace, and N. Arsem. 2015. Declines in polybrominated diphenyl ether contamination of San Francisco Bay following production phase-outs and bans. Environ. Sci. technol. 49(2):777-784.
- Tordoff, H. B. and P. T. Redig. 1997. Midwest peregrine falcon demography, 1982-1995. Journal of Raptor Research 31:339-346.
- Tuttle, G. and H. Castro. 2015. Surface water monitoring program for pesticides in salmonid-bearing streams, 2014 data summary. AGR PUB 104-494. Washington Department of Agriculture, Olympia, WA.
- (USFWS) U.S. Fish and Wildlife Service. 1982. Pacific Coast recovery plan for the American peregrine falcon (*Falco peregrinus anatum*).
- (USFWS) U.S. Fish and Wildlife Service. 2003. Monitoring plan for the American peregrine falcon: a species recovered under the Endangered Species Act. U.S. Fish and Wildlife Service, Divisions of Endangered Species and Migratory Birds and State Programs, Pacific Region, Portland, Oregon.
- (USFWS) U.S. Fish and Wildlife Service. 2008. Final revised environmental assessment and management plan: take of migrant peregrine falcons from the wild for use in falconry and reallocation of nestling/fledgling take. U.S. Fish and Wildlife Service, Arlington, Virginia.
- Varland, D.E., J.B. Buchanan, T.L. Fleming, M.K. Kenney, and T.M. Loughin. 2012. Peregrine falcons on coastal beaches of Washington: Fifteen years of banding and surveys. Journal of Raptor Research 46:1.
- Varland, D.E., T. L. Fleming, J. B. Buchanan. 2008a. Tundra peregrine falcon (*Falco peregrinus tundrius*) occurrence in Washington. Washington Birds 10:48-57.
- Varland, D. E., L. A. Powell, M. K. Kenney, and T. L. Fleming. 2008b. Peregrine falcon survival and resighting frequencies on the Washington coast, 1995-2003. Journal of Raptor Research 42:161-171.
- Watts, B.D., K.E. Clark, C.A. Koppie, G.D. Therres, M.A. Byrd, and K.A. Bennett. 2015. Establishment and growth of the peregrine falcon breeding population within the mid-Atlantic coastal plain. Journal of Raptor Research 49(4):359-366.
- Washington Department of Fish and Wildlife. 2008. Priority Habitat and Species List. Olympia, Washington.
- White, C.M., N.J. Clum, T.J. Cade, and W.G. Hunt. 2002. Peregrine Falcon (*Falco peregrinus*), In The Birds of North America Online, edited by A. Poole. Ithaca, New York: Cornell lab of Ornithology.

- Wolf, S. G., M. A. Snyder, W. J. Sydeman, D. F. Doaks, and D. A. Croll. 2010. Predicting population consequences of ocean climate change for an ecosystem sentinel, the seabird Cassin's auklet. *Global Change Biology* 16:1923-1935.
- Wootton, J. T. and D. A. Bell. 1992. A metapopulation model of the peregrine falcon in California: viability and management strategies. *Ecological Applications* 2:307-321.
- Zota, A.R., L. Linderholm, J.S. Park, M. Petreas, T. Guo, M.L. Privalsky, R.T. Zoeller, and T.J. Woodruff. 2013. A temporal comparison of PBDEs, OH-PBDEs, PCBs, and OH-PCBs in the serum of second trimester pregnant women recruited from San Francisco General Hospital, California. *Environ. Sci. Technol.* 47(20):11776-11784.

Appendix A. Occupancy and reproductive success at American peregrine falcon nesting territories included in the federal post-delisting monitoring, Washington, 2003-2012.

Year	Nesting Sites						Number of Young			
	Sites Checked (n)	Occupied (n) %		Occupied With Known Outcome (n) %		Successful (n) %		Observed (n)	/Occupied Known Outcome	/Successful Nesting Attempt (brood size)
2003	24	22	92	22	100	11	50	23	1.04	2.09
2006	24	18	75	18	100	11	61	29	1.61	2.64
2009	25	22	88	22	100	12	54	32	1.45	2.67
2012	25	21	84	21	100	16	76	38	1.81	2.38