

Draft
Washington State Status Report
for the
Sage Grouse

by

Michelle J. Tirhi
David W. Hays

Washington Department of Fish and Wildlife
600 Capitol Way North
Olympia, Washington 98501-1091

October 1997

The Washington Department of Fish and Wildlife maintains a list of endangered, threatened and sensitive species (Washington Administrative Codes 232-12-014 and 232-12-011, Appendix C). In 1990, the Washington Fish and Wildlife Commission adopted listing procedures developed by a group of citizens, interest groups, and state and federal agencies (Washington Administrative Code 232-12-297, Appendix C). The procedures include how species listing will be initiated, criteria for listing and de-listing, public review and recovery and management of listed species.

The first step in the process is to develop a preliminary species status report. The report includes a review of information relevant to the species' status in Washington and addresses factors affecting its status including, but not limited to: historic, current, and future species population trends, natural history including ecological relationships, historic and current habitat trends, population demographics and their relationship to long term sustainability, and historic and current species management activities.

The procedures then provide for a 90-day public review opportunity for interested parties to submit new scientific data relevant to the status report, classification recommendation, and any State Environmental Policy Act findings. During the 90-day review period, the Department holds statewide public meetings to answer questions and take comments. At the close of the comment period, the Department completes the Final Status Report and Listing Recommendation for presentation to the Washington Fish and Wildlife Commission. The Final Report and Recommendation are then released 30 days prior to the Commission presentation for public review.

This is a Draft Status Report for the Sage Grouse. **Submit written comments on this report by January 23, 1998 to: Endangered Species Program Manager, Washington Department of Fish and Wildlife, 600 Capitol Way N, Olympia, WA 98501-1091.** The Department will present the results of this status review to the Fish and Wildlife Commission for action at its April 3-4, 1998 meeting.

This report should be cited as:

Tirhi, M. J., and D. W. Hays. 1997. Washington State status report for the sage grouse. Wash. Dept. Fish and Wildl., Olympia. 62pp.

TABLE OF CONTENTS

LIST OF TABLES AND FIGURES	vi
ACKNOWLEDGMENTS	vii
EXECUTIVE SUMMARY	viii
TAXONOMY	1
DESCRIPTION	1
Plumage and Extremities	1
Measurements	1
GEOGRAPHICAL DISTRIBUTION	2
North America	2
Washington	3
NATURAL HISTORY	4
Behavior	5
Roost	5
Territoriality and sociality	5
Sexual behavior	5
Food, Nutrition, and Energetics	5
Water	6
Life Span and Mortality Rates	7
Density	7
Home Range	8
HABITAT REQUIREMENTS	9
General	9
Elevation	10
Breeding	10
Late-Spring, Summer, and Fall	11
Winter	12
POPULATION DYNAMICS	12
Reproduction	12
Age of first breeding	12
Seasons	13
Fidelity to leks	13
Mating	13
Nesting	13

Nesting success	14
Brood-rearing	14
Juveniles	14
Mortality	15
Predation	15
Hunting	15
Weather	16
Disease	16
Accidents and pesticides	16
POPULATION STATUS	17
Past	17
Harvest	17
Statewide population	19
Regional populations	22
Present	24
HABITAT STATUS	25
Past	25
Present	26
CONSERVATION STATUS	29
Legal Status	29
Management Activities in Washington	29
Management plan	29
Population monitoring	29
Habitat restoration/acquisition	30
Research	30
Coordination and partnership	31
Enforcement	31
Information and education	31
Habitat protection and enhancement	32
FACTORS AFFECTING CONTINUED EXISTENCE	32
Population Isolation	32
Potential for Continued Population Decline	33
Compatibility with Military Training	33
Habitat Security	34
Relationships with Private Landowners	35
Fire	35
Chemical Treatment	36
Grazing	37
Harassment and Disturbance	38

Adequacy Of Existing Regulatory Mechanisms 38

CONCLUSIONS AND RECOMMENDATION 38

REFERENCES CITED 39

PERSONAL COMMUNICATIONS 52

ACRONYMS 53

Appendix A. Museum specimens of sage grouse in Washington, 1853-1949 54

Appendix B. Sage grouse lek counts in Washington 1970-1997 55

Appendix C. Washington Administrative Code 232-12-297 58

LIST OF TABLES AND FIGURES

Table 1. Home range (km ²) estimates of sage grouse in Washington and Idaho.	9
Table 2. Hunting seasons and harvests of sage grouse in Washington, 1950-1987	18
Table 3. Historical and remaining shrub steppe (ha) habitat in Washington counties within the historic range of sage grouse	27
Table 4. Administration and ownership of sage grouse range in Washington	28
Table 5. Conservation Reserve Program (CRP) lands (ha) in eastern Washington counties in current sage grouse range through 1996	29
Figure 1. Historic and current range of the sage grouse in North America.	2
Figure 2. Historic and current range of the sage grouse in Washington.	4
Figure 3. The proportional change in sage grouse numbers estimated from annual lek counts, 1955-1997.	19
Figure 4. The ratio of males per lek and the number of leks included in counts, 1954-1997.	20
Figure 5. Male sage grouse counted statewide during lek counts in Washington, 1954-1997. Statewide counts are influenced by the number of leks included in the count.	21
Figure 6. Estimated current and historic population size of sage grouse based upon annual proportional changes in lek counts.	21
Figure 7. Lek counts in Lincoln County, 1970-1997.	22
Figure 8. Lek counts in Douglas County, 1961-1997.	23

ACKNOWLEDGMENTS

This status report uses information from the *Washington State Management Plan for the Sage Grouse*, written in 1995 by Michelle Tirhi.

Dave Ware, Michael Schroeder, Lisa Fitzner, Fred Dobler, Jerry Hickman, Ron Friesz, and Mark Quinn were members of the Washington Department of Fish and Wildlife Sage Grouse Steering Committee.

Valuable insights and critical reviews of earlier drafts were provided by Mike Schroeder, Upland Bird Research Biologist, and Dave Ware, Upland Bird Program Manager. Harriet Allen, Scott Richardson, Lee Stream and Tom Juelson also reviewed the draft.

Helpful information was offered by Steve Kruger and Margaret Pounds, U.S. Department of the Army, Environmental and Natural Resource Division, Yakima Training Center. Assistance was also provided by the state and regional offices of the U.S. Bureau of Land Management; the U.S. Department of Agriculture, Natural Resources Conservation Service; the U.S. Department of Agriculture, Agricultural Stabilization and Conservation Service; and the U.S. Department of Commerce, Bureau of the Census.

Kay Bujacich-Smith helped to prepare the maps.

Funding was provided in part by Federal Aid to Wildlife Restoration funds under the Pittman-Robertson Projects. These funds are derived from an excise tax on sporting guns and ammunition paid by hunters and recreational shooters. Funding was also provided by Wildlife Diversity funds derived from the sale of personalized license plates.

EXECUTIVE SUMMARY

Sage grouse inhabit the shrub steppe and meadow steppe region of eastern Washington. Typically, low rolling hills and adjacent valleys provide the best topography and habitat for sage grouse. Suitable sage grouse habitat contains medium to dense sagebrush stands with equal amounts of tall and short sagebrush plants and equal mixtures of forbs and grasses.

Sage grouse were historically distributed throughout much of the western United States and the southern border of three western Canadian provinces. Their range followed the distribution of sagebrush of the climax sagebrush and prairie ecosystems. From 1900 to the 1930's, sage grouse populations steadily declined throughout North America. From 1940 to 1950, sage grouse declines stabilized but remained permanently reduced because of habitat loss and degradation.

In Washington sage grouse historically ranged from the southern portion of the Columbia River, north to Oroville, west to the foothills of the Cascades, and east to the Spokane River. By 1860, sage grouse had declined and were rarely seen on some areas that had formerly contained numerous birds. By 1908, sage grouse had been extirpated from various counties historically containing small populations such as Spokane County. Breeding potential in Lincoln County was essentially eliminated by 1985 because of habitat alteration. The sage grouse population on the Fitzner and Eberhardt Arid Lands Ecology Reserve at Hanford (Hanford Site) in Benton County has evidently been extirpated due to catastrophic fires in 1981 and 1984. No sage grouse have been found there in recent surveys. A viable population of sage grouse no longer inhabits the Badger Pocket area of Kittitas County. Sage grouse in this area, historically known to support large numbers were extirpated by 1987.

Recent surveys indicate 2 remaining sage grouse populations in Washington: one within Douglas and Grant counties; and one in southern Kittitas and Yakima counties. These populations are relatively isolated, corresponding to partially intact areas of sagebrush habitat. Sage grouse are predominantly found on private lands in Douglas County and on the federally managed Yakima Training Center in Kittitas and Yakima counties. The largest contiguous block of shrub steppe remaining in Washington is located on the Yakima Training Center. Washington's sage grouse population is now considered isolated from surrounding populations in Idaho and Oregon.

The reduction in sage grouse numbers and distribution in Washington is primarily attributed to loss of habitat through conversion and degradation. Before the arrival of early settlers, the climax condition in the shrub steppe region of eastern Washington consisted of tracts of native sagebrush and bunchgrass species. Agricultural expansion, grazing, and sagebrush control through burning, mechanical removal, and chemical control severely degraded traditional sage grouse habitat. Approximately 40% of the estimated 4.16 million ha (10.4 million acres) of shrub steppe in eastern Washington before European settlement remains. Sage grouse habitat is a subset of this remaining acreage, and factors affecting occupancy include elevation, soil type, size of shrub steppe patch, and habitat quality.

Sage grouse sources of mortality include predation, weather, disease and parasitism, accidents, and environmental hazards such as pesticides. These natural and man-influenced factors become important management issues with small populations. Predation is considered a limiting factor throughout the annual sage grouse cycle. Raptors and coyotes are primary predators of sage grouse while corvids, badgers, and ground squirrels are primary nest predators. Weather influences nesting success and survival of young chicks. Several avian diseases and parasites have been reported for sage grouse; however, they do not appear to be a significant source of mortality.

Sage grouse populations have increased in Douglas County since 1992. This may represent the improving underlying habitat conditions or the cyclical patterns of population change indicated from past years. Sage grouse have been maintained over time on private lands in Douglas County due to portions of the land being poorly suited for agriculture. Sage grouse habitat has improved on lands under the federal Conservation Reserve Program (CRP). Listing sage grouse may be of concern for private landowners due to fears of regulation, and listing may result in strained relationships. It may also be a benefit to private landowners interested in the federal CRP program, as concerns for sage grouse habitat results in an increased likelihood of certain lands being enrolled in the CRP program.

Immediate threats to sage grouse populations include the uncertain future of the Conservation Reserve Program (especially in Douglas County), potential catastrophic fires, and the effects of military training on sage grouse and their habitat. Other threats to sage grouse habitat include continued conversion of shrub-steppe to cropland uses, and livestock grazing.

A significant downward population trend is indicated from lek counts and harvest information. Harvests averaged 1,842 sage grouse from 1951 to 1973, moved below 1,000 after 1974 to 18 in 1987. The season was closed in 1988. The number of males per active lek declined statewide from 35 males/lek in 1970 to 14 males/lek in 1994, before increasing somewhat to 16 males/lek in 1997.

The estimated sage grouse breeding population in Washington based upon 1997 survey results is approximately 900-1000 birds statewide. Approximately 600 of these birds are located in Douglas County and approximately 300 - 400 sage grouse are located in Kittitas and Yakima counties. Scattered sage grouse also exist on the periphery of the range but are not believed to play a significant role in the dynamics of the population in Washington.

Management activities in Washington include annual surveys by the Department of Fish and Wildlife to monitor populations, development of a statewide management plan for sage grouse, acquisition and restoration of habitat, and coordination of activities with other land management agencies. Research has and will continue to be conducted on both populations. A conservation agreement has been developed with the Yakima Training Center for management of lands critical to sage grouse on Department of Defense lands.

Washington farmers were allotted only 21 percent of the acreage they submitted for CRP funding during spring of 1997, one of the lowest acceptance rates in the nation. Contracts for approximately 785,000 acres are due to expire in 1997. Applications for approximately 819,000 acres were submitted, but only 172,000 acres were accepted. Applicants are re-applying during fall 1997, and increases in CRP enrollment are likely. The removal of important CRP habitat in Grant and Douglas counties would reverse habitat gains in recent years, and may cause further declines in sage grouse numbers.

The sage grouse population and corresponding sage grouse habitat in Washington has declined significantly. Sage grouse current range has declined to 10-15 percent of historic range. Population declines occurred in both the last century and this century. Local populations were extirpated as recently as the mid-1980's. Major threats that remain to the two relatively small populations include potential for catastrophic fire, impacts of military training, and the instability of the federal Conservation Reserve Program.

For these reasons, the Department recommends that the sage grouse be designated a State Threatened species.

TAXONOMY

Sage grouse belong to the family Phasianidae (pheasant-like birds) and subfamily Tetraoninae. Two subspecies of sage grouse are recognized: *Centrocercus urophasianus urophasianus* Bonaparte, or the eastern sage grouse, and *C. u. phaios* Aldrich, the western sage grouse. Genetic, morphological, and behavioral research is being conducted on sage grouse from different localities (M. Schroeder, pers. comm.). Preliminary evidence indicates sage grouse in a montane basin near Gunnison, Colorado, differ in several traits from conspecifics (Young 1991). The classification of a new species, *C. minimus* (Schroeder pers. comm.), is proposed. Common names of sage grouse include sage hen, sage fowl, spine-tail grouse, fool hen, cock of the plains, spiny-tailed pheasant, sage cock, and sage chicken (Coues 1893, Girard 1937, Patterson 1952, Jewett et al. 1953, Johnsgard 1973).

DESCRIPTION

Plumage and Extremities

Both sexes of adults have narrow, pointed tails, feathering to the base of the toes, and a variegated pattern of grayish brown, buffy, and black on the upper parts of the body (Johnsgard 1973). The flanks are pale and the abdomen exhibits a diffuse, black pattern. Males are larger and more colorful than females with a black throat and bib, scaly white foreneck feathers, and a large white ruff on the breast (Dunn et al. 1987). Males also exhibit two large, frontally directed air sacs (cervical apteria) of olive green skin and yellow superciliary combs which are enlarged during breeding display (Johnsgard 1973, Udvardy 1977). Females are characterized by grayish-white upper throats with an absence of black, no scaly feathers on the foreneck, and a partial extension of the white tips of the under tail coverts to the feather rachis (Pyrah 1963). The larger size, dark belly, and absence of white outer-tail feathers distinguish sage grouse from the sharp-tailed grouse (*Tympanuchus phasianellus*), a co-inhabitant of portions of sage grouse range in Washington.

Measurements

Sage grouse are the largest North American grouse species and Washington's largest native upland game bird. Sage grouse weights vary by sex, age, time of year, and population (Autenrieth 1981). For these reasons, comparisons of sage grouse weight can be misleading. Generally males weigh more than females and adults weigh more than yearlings. Adult males average 2-3.5 kg (4-7 lb), adult females average 1 to 2 kg (2-4 lb). Male weight is related to specialized body parts which are integral components of the male's display (Honest and Allred 1942, Clarke et al. 1942, Beck and Braun 1978). Adult males range in size from 66 to 76 cm (26-30 inch) and adult females range from 48 to 58 cm (19-23 inch) (Girard 1937, Jewett et al. 1953, Johnsgard 1973).

GEOGRAPHICAL DISTRIBUTION

North America

Historically, sage grouse were distributed throughout much of the western United States and along the southern border of three western Canadian provinces (Patterson 1952, Braun 1993) (Fig. 1). Their range followed the distribution of sagebrush (*Artemisia* spp.) north to British Columbia, south to New Mexico and Oklahoma, east into Nebraska, and west to California (Aldrich 1963, Guiquet 1970). Lewis and Clark first reported sage grouse at the head of the Missouri River, on the plains of the Columbia, and at the mouth of the Snake River (Coues 1893). Historical reports describe large numbers of sage grouse throughout their range (Coues 1893, Huntington 1897, Burnett 1905, Wilhelm 1970). Sage grouse populations declined throughout North America from 1900 to 1940 primarily due to habitat loss, uncontrolled grazing, and unrestricted hunting (Patterson 1952, Jewett et al. 1953).




Figure 1. Historic and current range of the sage grouse in North America.

Currently, sage grouse range from southeastern Alberta and southern Saskatchewan, south to central Washington, eastern Oregon and California, and east to Nevada, Utah, western Colorado, southeastern Wyoming, and the western borders of North and South Dakota. The core range includes Colorado, Idaho, Montana, Nevada, Oregon, and Wyoming, with remnant populations in other states (Fig. 1). Sage grouse have been extirpated in British Columbia, Nebraska, New Mexico, and Oklahoma (Braun 1991, 1993). Braun (1993) considered populations remaining in Alberta, North Dakota, Saskatchewan, South Dakota, California, Colorado, Utah, and Washington as greatly reduced and possibly marginal. The western subspecies of sage grouse (*C. u. phaios*) occurs in disjunct populations in central Washington and southeast Oregon (Johnsgard 1973).

Washington

Historically, sage grouse ranged from the southern portion of the Columbia River to Oroville in the north, west to the Yakima River and east to the Spokane River (Yocom 1956) (Fig. 2). Lewis and Clark described large numbers of sage grouse near the mouth of the Snake River in 1805 (Jewett et al. 1953). Early explorers also reported sage grouse in Priest Rapids, Pine Creek, Alder Creek, Horse Heaven Sand Hills, the Blue Mountains area, the sagebrush areas surrounding the Columbia River, and the Yakima and Simcoe Valley from 1840 to 1900 (Baird et al. 1874; Royal Historical Soc. 1914 and Ballou 1938, cited in Yocom 1956). Sage grouse also inhabited the Okanogan Valley in British Columbia and areas bordering the Palouse, Snake, Touchet, and Walla Walla rivers (Yocom 1956). Rare but resident sage grouse were found in Asotin County. Sage grouse occurred in 16 counties, with the largest concentrations likely in Adams, Douglas, Yakima, Franklin, Grant, and Lincoln counties in the Big Bend, Moses Coulee, Grand Coulee, and Crab Creek drainages southward to the Snake and Columbia rivers (Yocom 1956). By 1860, sage grouse had declined and were rarely seen on areas that had formerly contained numerous birds (Cooper 1869 and Cleman 1918, cited in Jewett et al. 1953). By 1908, sage grouse had been extirpated from counties historically containing small populations, such as Spokane County (Jewett et al. 1953).



Figure 2. Historic and current range of the sage grouse in Washington.

Historic sage grouse range (Yocom 1956, Aldrich 1963) compared to current areas reveals a 85-90% reduction. The current range includes both year-round and seasonal concentration areas and both contiguous and fragmented habitats (Fig. 2). Two remaining sage grouse populations are found in Douglas and northern Grant counties, and southern Kittitas and Yakima counties. These populations are relatively small and isolated, corresponding to partially-intact areas of sagebrush habitat. Although the population declined in Douglas County, there was little reduction in range. The largest numbers of sage grouse remain in Douglas County and a smaller population occurs in Kittitas and Yakima counties.

NATURAL HISTORY

Behavior

Roost. Sage grouse usually roost on the ground from evening until early morning, feed and rest during the afternoon, and return to their roosting site at night (Johnsgard 1973). In Nevada, sage grouse burrowed into the snow when snow depth was ≥ 25 cm (10 in) and minimum temperature was < -10 C° (14 F°) (Back et al. 1987).

Territoriality and sociality. Male and female sage grouse gather within the same area during spring for displaying and mating. The area is called a lek. At the beginning of the breeding season, male sage grouse establish small territories on the lek. Males occupying territories near the center of the lek may be more successful at mating (Davis 1978). Males seek leks that are visited by many females. The ability to attract females and copulate is determined through display and aggressive behavior by males (Hartzler and Jenni 1988). Adult males typically occupy and establish territories first and dominate the breeding activities.

During most of the year, sage grouse segregate into flocks according to sex (Hjorth 1970). In late summer and early fall, sage grouse congregate in heterogeneous flocks in preparation for movement to wintering grounds (Patterson 1952). Similar congregations occur in late winter or early spring before movement to breeding areas (Patterson 1952). Not all sage grouse populations undergo these movements, however (Connelly et al. 1988).

Sexual behavior. The most distinguishable sexual behavior is the courtship display performed by males on leks during spring. Display allows males to proclaim and defend a territory and attract females for copulation (Johnsgard 1973). Males stand 25 to 50 cm (10-20 in) apart on their territory with tail feathers spread, wings dropped, neck feathers ruffled, and the cervical apteria inflated (Johnsgard 1973). To begin the display a male stands erect, fans his tail, lowers his folded wings, and steps forward. The feathers of the tail contrast sharply with natural colors in the shrub-steppe environment; the tail attracts females and serves as a warning to males (Hjorth 1970). Ruffled neck feathers, inflated air sacs, and enlarged eye combs likely serve the same function (Hjorth 1970). The back is gradually raised, the white feathers above the eyes (filoplumes) are erected, and the cervical air sacs exposed. Next, the air sacs are pushed outward and the male jerks upwards. The sequence is then repeated with slight variations and more steps. Toward the end of the display, males pull their heads into the neck feathers and completely inflate their esophageal pouches (air sacs). The release of air from the pouches makes a *plopping* sound and marks the end of one display (Johnsgard 1973).

Food, Nutrition, and Energetics

Sagebrush, grasses, forbs, and insects comprise the annual diet of sage grouse. Sagebrush comprises 60 to 80% of the yearly diet of adult sage grouse (Martin et al. 1951, Patterson 1952,

Wallestad et al. 1975, Rasmussen and Griner 1938, Remington and Braun 1985) and as much as 95 to 100% of the winter diet (Roberson 1984). Forbs may constitute $\geq 50\%$ of the juvenile diet (up to 11 weeks of age) (Rasmussen and Griner 1938, Klebenow and Gray 1968, Peterson 1970). Forbs also appear to be important to nesting hens in the pre-laying period (Barnett and Crawford 1993).

Insects make up $\geq 50\%$ of the diet during the first and second week of life (Rasmussen and Griner 1938, Patterson 1952, Klebenow and Gray 1968, Peterson 1970). Johnson and Boyce (1990) found chicks < 3 weeks old required insects for survival and chicks > 3 weeks old had reduced growth rates when insects were removed from the diet. In highly fluctuating sage grouse populations (usually those found in the xeric 18 to 25 cm [7-10 in] precipitation zone), important post-hatch limiting factors are forb and insect availability (Autenrieth 1981). In Oregon, forbs and invertebrates composed 80% of the dietary mass of chicks in one area with higher grouse productivity, whereas sagebrush composed 65% of the dietary mass of chicks on a second lower productivity site.

In fall, sage grouse shift back to a sagebrush dominated diet (Girard 1937; Griner 1939 and Bean 1941, cited in Roberson 1984; Dargan et al. 1942; Patterson 1952; Trueblood 1954; Nelson 1955; Klebenow and Gray 1968; Savage 1969; Martin 1970; Peterson 1970; Oakleaf 1971; Wallestad et al. 1975; Autenrieth 1980). Remington and Braun (1985) observed sage grouse in winter selecting vegetative species with the highest protein levels, possibly because plants with high levels of protein are easily digestible and provide instant energy. Protein levels differ among subspecies of sagebrush; Wyoming big sagebrush (*Artemisia tridentata wyomingensis*) may have the highest levels (Remington 1983). In Washington, big sagebrush (*Artemisia tridentata*) is the predominant species consumed (Eberhardt and Hofmann 1991).

Sage grouse require a higher quality winter diet relative to other grouse (Roberson 1984, Welch et al. 1988) and all age and sex classes gain or maintain weight and fat over the winter (Beck and Braun 1978, Roberson 1984, Remington and Braun 1988). Sage grouse experience a negative energy balance during the breeding and nesting period and subsequently lose weight (Beck and Braun 1978, Roberson 1984).

Nutrition affects productivity of all Tetraoninae (Moss et al. 1975). Poor productivity may be caused by inadequate nutrition of hens during the breeding season (Moss et al. 1975, Barnett and Crawford 1993). Females in good nutritional condition contribute more nutrients to eggs (Jenkins et al. 1965), which leads to larger clutches and increased chick viability (Jenkins et al. 1963, Eastman and Jenkins 1970). Sage grouse may select both shrubs and forbs with high nutritional value.

Water. Some researchers consider water a key component of sage grouse habitat (Carr 1967, Savage 1969, Call and Maser 1985). Others have failed to find sage grouse preferring sites close to water (Wallestad 1975, Autenrieth 1981, Cadwell et al. 1994). Sage grouse need to consume water, but they typically obtain enough water by consuming vegetation that stores water, such as

succulent forbs. Sage grouse may concentrate in late summer and fall where water or succulent forbs are available. Water sources include streams, springs, water holes, and cattle troughs. Where water is available, sage grouse normally visit water sites in the morning and evening. Sage grouse that occupy areas with little precipitation may migrate to areas containing water during the summer and fall. Chicks require water soon after hatching (Girard 1937), so hens with broods often migrate to areas containing water. Petersen (1980) found that hens with broods remained in upland habitat until succulent forbs disappeared then moved to wet meadows in late summer.

Sex Ratios

Sex ratios are typically determined from information supplied by hunters or wing samples taken from harvested birds. A male to female sex ratio of 1:1 for all sage grouse (Girard 1937), and 1:1.2 and 1:2.3 for juvenile sage grouse (Patterson 1952) were reported in Wyoming. In Colorado, sex ratios of 1:1.5 for all sage grouse (Rogers 1964), 1:1.1 for juveniles, 1:1.6 for yearlings, and 1:2.6 for adults (Braun 1984) were reported. Because of their conspicuousness during breeding, more adult males may be killed by predators than females. This may result in adult sex ratios that are swayed towards females as compared with more equitable sex ratios for chicks and juveniles.

Life Span and Mortality Rates

Sage grouse can survive up to 10 years in the wild (C. Braun, pers. comm.; J. Connelly, pers. comm.). The average life span of sage grouse in both hunted and protected populations in one study was 1 to 1.5 yr (Elman 1974); in another study sage grouse 3 to 4 yr old were considered old (Wallestad 1975).

Braun (1975) reported that annual mortality rates for sage grouse were 50 to 55% and for greater prairie chickens 54%; these were the lowest annual mortality rates of all prairie grouse species (Braun 1975). In Colorado, Zablan (1990) found mortality rates of females to be higher than males (55 and 38%, respectively). In Washington, annual mortality rate for sage grouse was approximately 40% (M. Schroeder, pers. comm.). Chick mortality ranges from 40% (Wallestad 1975) to >60% (Braun 1975). In Montana, Wallestad (1975) estimated a 56% mortality rate for sage grouse chicks between time of hatch and the following fall, a 65% mortality rate for yearling hens, and a 60% mortality rate for adult hens. J. Connelly (pers. comm.) reported an annual mortality rate of 50 to 60% for all sage grouse in Idaho.

Density

Sage grouse density fluctuates in response to habitat availability and population dynamics. Patterson (1952) reported an average of one lek per 15 km² (6 mi²) with a density of 5 males/km² (13 males/mi²). Edminster (1954) analyzed Patterson's data to estimate 12 to 19 grouse/km² (30-50 grouse/mi²) or 5 to 8 ha (13-21 ac) per grouse. In Colorado, Rogers (1964) estimated a

fall density of 12 to 19 grouse/km² (30-50 grouse/mi²) in areas containing the best sage grouse habitat. Fall density in areas having marginal habitat ranged from 0.4 to 12 grouse/km² (1-30 grouse/mi²) (Rogers 1964). In Oregon, Gregg (1991) determined 3 grouse/km² (8 grouse/mi²) in good habitat and 1 grouse/km² (3 grouse/mi²) in an area representing typical, remaining habitat. In Nevada, Zunino (1987) determined a spring density of 0.1 to 3 males/km² (0.3-8 males/mi²) and a fall density of 0.4 to 5 males/km² (1-13 males/mi²). A compilation from the 13 states and provinces that contain sage grouse had an average count of 16 to 32 males/lek from 1988 to 1990 (Braun 1991).

Home Range

Sage grouse have large home ranges in comparison to other grouse species (Bergerud 1988). In Washington, home ranges of sage grouse in Douglas County were studied by Pederson (1982), and home ranges of sage grouse on the Yakima Training Center (YTC) in Kittitas and Yakima counties were studied by Eberhardt and Hoffman (1991) (Table 1). Estimated home ranges of sage grouse in Douglas County were significantly smaller than home ranges of sage grouse on the YTC. Eberhardt and Hoffman (1991) attributed the large home ranges of sage grouse on the YTC to repeated disturbance. However, significant differences exist between studies in number of relocations, number of females followed, and time of year followed. Maximum distances moved from leks ranged from 4 to 36 km (2-22 mi) in Washington (Pedersen 1982, Eberhardt and Hofmann 1991, Cadwell et al. 1994, Schroeder 1994).

Seasonal movements depend on topography, vegetative cover, winter weather, and availability of winter food (Beck 1975, Autenrieth 1986). Wintering females have overlapping ranges because they share refuge areas (Bradbury et al. 1989b). In Washington, most sage grouse migrated between breeding areas and winter areas, and winter areas were somewhat distinct from summer areas for both females and males (Schroeder 1994). Schroeder (1994) hypothesized that this was due to the overgrazed condition of the winter range. Adult sage grouse often return to specific wintering areas regardless of weather or food conditions.

Sage grouse occupying sagebrush communities at low elevation often do not migrate (Wallestad 1975), and those inhabiting mountain valleys or areas with distinct elevation gradients are often migratory (Dalke et al. 1980, Connelly et al. 1988). Migratory sage grouse generally move >16 km (10 mi) (Baker 1978, cited in Roberson 1984; Berry and Eng 1985). Migrations of 80 to 160 km (50-100 mi) from wintering areas to leks (Pyrah 1954, Dalke et al. 1963) and 81 km (50 mi) from leks to winter range (Connelly and Markham 1983) have been reported, but shorter distances are more common (Bradbury et al. 1989a). Within their range, sage grouse move daily between feeding, breeding, and roosting sites (Wallestad 1971).

Mating, nesting, brood rearing, loafing, roosting, and foraging occur within 3 km (2 mi) of a lek (Wallestad and Pyrah 1974, Wallestad 1975, Autenrieth 1981), in some areas and at distances >3 km in other areas (Connelly et al. 1988, Wakkinen et al. 1992, Schroeder 1994).

Table 1. Home range (km²) estimates of sage grouse in Washington and Idaho.

Location	Range	Sex/Age ^a	n ^b	Season	Source
Idaho	44	M	6	na	Sime (1991)
	6	F	na	na	
	12	Y	na	na	
Washington	4	M/A	13	Spring	Pedersen (1982)
	6	M/Y	1	Spring	
	8	M/Ad	14	Summer	
	2	M/Y	1	Summer	
	5	F/Ad	1	Spring-Summer	
	0.28	M/Ad	3	Winter	
Washington	2	F/Ad	1	Winter	Eberhardt and Hofmann (1991)
	29	C	25	Spring	
	26	C	25	Summer	
	44	C	25	Fall	

^a M=male, F=female, Ad=adult, Y=yearling, C=sex and age classes combined

^b na=not available

HABITAT REQUIREMENTS

General

Sage grouse inhabit the shrub steppe and meadow steppe region of eastern Washington. Shrub steppe is a descriptive term for plant communities consisting of one or more layers of perennial grass with a conspicuous, but discontinuous, layer of shrubs above (Daubenmire 1988). Shrub-steppe communities in Washington typically contain shrubs such as big sagebrush (*Artemisia tridentata*), three-tipped sagebrush (*A. tripartita*), and bitterbrush (*Purshia tridentata*), and a variety of grasses and forbs. Meadow steppe is a descriptive term for plant communities that are dense at ground level, support many grasses and forbs with broad leaves, have few shrubs, are barely dry enough to exclude trees (Daubenmire 1988), and generally have meadow characteristics (Franklin and Dyrness 1973). Typical meadow-steppe communities in Washington have several grasses, including bluebunch wheatgrass (*Agropyron spicatum*) and Idaho fescue (*Festuca idahoensis*) (Daubenmire 1988).

Canopy coverage is defined as the percentage of the ground surface covered with vegetation. In this report, four levels of sagebrush canopy coverage are differentiated: low (5 to 14%), medium (15 to 25%), high (26 to 40%), and very high (>40%). Big sagebrush, commonly the dominant shrub in Washington's shrub steppe, seldom grows with canopy coverage >40%, except in areas having deep soil and local moisture (J. Connelly, pers. comm.; C. Perry, pers. comm.).

Low rolling hills and adjacent valleys provide the best topography for sage grouse (Call and Maser 1985). Sage grouse prefer slopes <30% (Call and Maser 1985) facing south year round (Rogers 1964). Suitable habitat consists of sagebrush/bunchgrass stands having medium to high canopy cover of sagebrush in a variety of height classes and a diverse grass and forb understory (Peterson 1970, Wallestad 1971, Eng and Schladweiler 1972). Sagebrush is used as food and cover all year. Winter habitat is typically the most limited seasonal habitat within the range of sage grouse (Patterson 1952, Eng and Schladweiler 1972, Beck 1977).

Elevation

Sage grouse are found at elevations of 1,200 to 2,400 m (4,000-8,000 ft) in Oregon (Call and Maser 1985), 2,400 to 2,600 m (7,800-8,500 ft) in Colorado (Rogers 1964), 1,200 to 2,900 m (3,900-9,500 ft) in Idaho (Girard 1937, Klebenow and Gray 1968), and 2,500 to 2,700 m (8,000-9,000 ft) in Wyoming. In Washington, sage grouse on the Yakima Training Center (YTC) were found at elevations of 500 to 900 m (1,650 to 2,970 ft) (Cadwell et al. 1996b).

Breeding

Leks are the focal point of the breeding season and range in size from 0.04 to 40 ha (0.1-99 ac) (Scott 1942, Call 1979, Call and Maser 1985). Leks are often near nesting habitat and between areas used in winter and summer (Wallestad 1975, Klebenow 1985, Connelly et al. 1988). They are found in gravel pits, burned areas, cultivated fields, air strips, abandoned homesteads, roads, bare ridges, grassy swales, natural and irrigated meadows devoid of grass, knolls, small buttes, openings in sagebrush stands, dry-lake beds, and areas denuded of vegetation by livestock (Roberson 1984, Call and Maser 1985, Autenrieth 1986). Given the diverse habitats where leks are placed, lek habitat availability is likely not a limiting factor for sage grouse (M. Schroeder, pers. comm.).

Most leks contain a central area that is barren and a surrounding area containing shrubs (Klebenow 1985, Ellis et al. 1989, Klott and Lindzey 1989). However, in Washington, numerous active leks are devoid of surrounding shrubs (L. Fitzner, pers. comm.; Schroeder 1994). Most leks on the Yakima Training Center contain big sagebrush-bluebunch wheatgrass, occur on loamy soils, are moderately sloped (0-10%), and average 36 ha (89 ac) in size (Eberhardt and Hofmann 1991). Visibility is important on a lek (Gill 1965, Wiley 1973) and is necessary for females to observe displaying males and for all sage grouse to observe predators. In Washington, sage grouse preferred 0 to 10% slopes during spring (Cadwell et al. 1994). Water is not necessary on a lek (M. Schroeder, pers. comm.), although leks are often located near water (Call 1979). In areas containing both sage grouse and livestock, sage grouse may form leks near water because the vegetation has been flattened or removed by repeated livestock use (J. Connelly, pers. comm.).

Shrubs surrounding leks are used extensively by sage grouse. Males use shrubs ≤ 1 km (0.6 mi) from a lek for foraging, loafing, and shelter (Rothenmaier 1979, Emmons and Braun 1984,

Autenrieth 1981). Shrub stands with medium to very high shrub cover are primarily used by sage grouse for foraging and loafing (Autenrieth 1981, Emmons and Braun 1984, Roberson 1984). Males select shrub stands 18 to 38 cm (7-15 in) high (Call and Maser 1985, Rothenmaier 1979) with a canopy coverage of 20 to 50% (Wallestad and Schladweiler 1974, Autenrieth 1981, Ellis et al. 1989).

Late-Spring, Summer, and Fall

After mating, sage grouse hens leave the lek to nest. Most hens build nests under shrubs (Jarvis 1974, Wallestad and Pyrah 1974, Roberson 1984), specifically in areas with medium-high shrub cover and residual grass (Schoenberg 1982, Gregg 1991, Sime 1991). However, females occasionally nest in grassland (Sveum 1995), cultivated fields that contain abundant insects for chicks after hatching (Autenrieth 1981), or in idle cropland, such as land enrolled in the Conservation Reserve Program (CRP) (Schroeder 1994). Sage grouse may require a balance of shrubs and grasses for greatest nest success (Sveum 1995). Shrubs located in nesting habitat act as an umbrella, which protects the nest and increases nest success (Autenrieth 1981, Connelly et al. 1991, Gregg et al. 1994). In Washington, big sagebrush/bunchgrass is the predominant habitat selected for nesting (Schroeder 1994, Sveum 1995). In other states, additional species used for nesting include rabbitbrush (*Chrysothamnus* sp.), bitterbrush, three-tipped sagebrush, silver sagebrush (*A. cana*), and mountain big sagebrush (*A. tridentata vaseyana*) (Gregg 1991, Sime 1991, Gregg et al. 1994).

Tall, dense vegetation provides visual, scent, and physical barriers between predators and the nests of ground-nesting birds (Redmond et al. 1982; Sugden and Beyersbergen 1986, 1987; Crabtree et al. 1989; Sveum 1995). Grass, especially tall grass, and forbs interspersed with sagebrush increase nest success (Autenrieth 1981, Wakkinen 1990, Gregg 1991, Sveum 1995). Grass may increase nest success by hiding the nest from ground predators and forming a microclimate that is warmer than the air above (Autenrieth 1981). Nests on the Yakima Training Center were characterized by a more even mixture of grass and shrubs in contrast to areas where unsuccessful nests occurred (Cadwell et al. 1996b). Hens also have been found nesting under the tallest sagebrush available (Wallestad and Pyrah 1974, Braun et al. 1977, Sime 1991) in other states, but this has not been found in Washington (M. Schroeder, pers. comm.). In Oregon, Crawford and DeLong (1993) found that nests placed in tall grass (>15 cm [6 in]) and medium high shrubs (40-80 cm [16-32 in]) had the least predation. This study included both real sage grouse nests and artificial nests placed by researchers. A separate study in Oregon also found that sage grouse nests placed in medium high shrubs had the least predation (Gregg et al. 1994). Nonpredated nests had a grass canopy coverage of 18% and shrub coverage of 41% within 1 m (3 ft) of the nest; grass coverage averaged 5% and shrub coverage averaged 29% at predated nests (Gregg et al. 1994).

In Washington, most females nested in sagebrush and grass that were relatively tall (124 cm [49 in] and 108 cm [43 in] respectively) and had medium to very high canopy coverage (20% and 51%, respectively) (Schroeder 1994). Both sagebrush and tall grasses are important components

of nesting habitat on the Yakima Training Center (Sveum 1995). The sagebrush/bunchgrass habitat type on the YTC accounted for 71 percent of nests during 1992 and 1993 (Sveum 1995).

During summer in Washington, Pedersen (1982) observed sage grouse moving from sagebrush communities to wet areas that contained annual forbs in fallow fields. Sage grouse on the Yakima Training Center did not frequent springs, nor did they prefer major streams and associated riparian areas for water and food (Cadwell et al. 1994). Sage grouse broods used both big sagebrush/bunchgrass and grasslands on the YTC (Sveum 1995). In Oregon, sage grouse were generally observed feeding on forbs near playas, water holes, and meadows in summer (Willis et al. 1993).

Winter

Sagebrush, which constitutes nearly 100% of the winter diet, is tall, evergreen, and nutritious. These properties become important when snow depth exceeds 30 cm (12 in) (Autenrieth 1981, Hupp and Braun 1989, Willis 1991). Sage grouse prefer sagebrush ≥ 25 cm (10 in) high with $\geq 15\%$ canopy coverage (Wallestad and Schladweiler 1974, Autenrieth 1981, Schoenberg 1982) and forage in the tallest sagebrush with the highest canopy cover (Beck 1977).

Deep snow limits food availability and may prevent a flock from using a site in winter. The best wintering sites are often located at the lowest elevations (Rogers 1964) in areas having flat or gentle slopes with $<15\%$ gradient (Jarvis 1974, Beck 1977, Autenrieth 1981). Winter sites typically face south or west (Beck 1977, Autenrieth 1981, Hupp and Braun 1989), possibly because less snow accumulates than on north or east aspects (Beck 1977, Hupp and Braun 1989). Drainage basins with abundant sagebrush (Pedersen 1982, Schoenberg 1982, Hupp and Braun 1989), or dry areas that may be unsuitable other times of the year, are often used during winter. In Washington, sage grouse were observed feeding on steep ($>15\%$) south-facing slopes and roosting on gradual slopes (15%) and ridgetops during winter (M. Schroeder, pers. comm.). On the Yakima Training Center, males used areas with more grass and less shrub cover in winter when compared with nest sites (Cadwell et al. 1996b).

POPULATION DYNAMICS

Reproduction

Age of first breeding. Female sage grouse are sexually mature their first fall and nest the following spring (Patterson 1952). Males are sexually mature their first spring after hatching. [Juvenile males engage in display and breeding but devote less time and energy to courtship activities than adults \(Wiley 1974\).](#)

Seasons. The mating season generally begins at the same time each year but ultimately depends on weather and vegetative conditions. In Washington, most females returned to breeding areas in late February to early March (Eberhardt and Hofmann 1991, Schroeder 1994), while most males returned from early to mid-April (Eberhardt and Hofmann 1991). Pedersen (1982) recorded the highest number of male and female sage grouse on leks from mid-March to mid-April.

Fidelity to leks. Most male sage grouse return to the same lek (Wiley 1978) or lek complex (Schlatterer 1960, Jarvis 1974, Braun and Beck 1976) each spring. In Montana, almost all males returned to the same lek regardless of their success at mating (Hartzler and Jenni 1988). Male sage grouse in Washington demonstrate strong fidelity to leks. All males ($n=43$) in Pederson's (1982) study remained at the same lek throughout the breeding season, and all males ($n=4$) in a study by Schroeder (1994) that were monitored during consecutive breeding seasons attended the same lek. An arrowhead used to hunt sage grouse and made approximately 105- to 110-years ago was found at an active lek in Idaho (Dalke et al. 1963), which demonstrates the longevity of some leks.

A male may choose a lek based on the number of females likely to visit (Bradbury et al. 1989b). Two scenarios could account for males returning to the same lek each year and concurrently searching for leks with many females. Individual females and males may attend the same lek in consecutive years by preference or coincidence, or individual females and males may attend different leks in consecutive years but are limited by the number of leks in a lek complex. Slight fluctuations in annual lek counts (Wash. Dept. Fish and Wildl., unpubl. data) and observations of sage grouse moving among leks in the same area (Dalke et al. 1963; Eng, unpubl. data cited in Wallestad and Schladweiler 1974; Braun and Beck 1976) support the second scenario.

Juveniles establish new leks and other leks become obsolete because of habitat changes or disintegration of local populations. Males may also move to a new location near a primary lek (J. Connelly, pers. comm.); the new location is called a "satellite," and males may or may not return to the new location in subsequent years. Females do not return to the same lek every year; instead, they search for active males and a good nesting site (Bradbury et al. 1989b).

Mating. Mating begins after males and females congregate on a lek. Hens form clusters near a centrally-located, dominant male (Hartzler and Jenni 1988), and these few dominant males participate in most of the mating (Eng and Schladweiler 1972). Males spend early morning and late evening at leks and remain nearby the rest of the day (Batterson and Morse 1948, Wallestad and Schladweiler 1974). After mating, males spend the summer alone or in small flocks. In Washington, males began to leave leks in late April and early May (Pedersen 1982) and moved directly to summer habitat that averaged 12 km (7 mi) from leks (Cadwell et al. 1994).

Nesting. After mating, females devote most of their time to building nests, laying eggs, and raising chicks; males do not assist in these activities (Rasmussen and Griner 1938, Patterson 1952, Harrison 1978). Females build nests approximately 10 days after mating (Autenrieth 1980).

Nests are typically constructed 2 to 6 km (1-4 mi) from leks (Gill 1965, Martin 1970, Jarvis 1974, Wallestad and Pyrah 1974, Petersen 1980, Pedersen 1982, Berry and Eng 1985, Eberhardt and Hofmann 1991, Wakkinen et al. 1992, Fischer et al. 1993), although movements <1 to 19 km (0.62-12 mi) were reported (Cadwell et al. 1994). In Washington, the average distance between the nest and the lek where the female was captured was 6.9 km (4.3 mi) (Schroeder 1994). Nest placement likely depends on habitat quality and not distance to the lek (Wakkinen et al. 1992). In California, hens chose nest sites before they chose a lek, they traveled farther from winter grounds to select a nest site than to select a lek, and after mating, they nested on the chosen site rather than nesting around the lek (Bradbury et al. 1989b). Hens returned to the same nest site each year in Idaho (Gates 1983, Fischer et al. 1993), Montana (Berry and Eng 1985), and Wyoming (Berry and Eng 1985).

Nesting success. Sage grouse nesting success (percentage of nests having eggs that hatch) ranges from 10 to 64% (Batterson and Morse 1948, Patterson 1952, Nelson 1955, Gill 1966, Wallestad and Pyrah 1974, Wallestad 1975, Petersen 1980, Connelly et al. 1988, Zunino 1987, Wakkinen 1990, Gregg 1991, Sime 1991). Bergerud (1988) analyzed nesting-success rates from 12 studies and reported an average of 35%. In Washington, Schroeder (1997) reported overall nesting success of 37%. Eighty seven percent of females re-nested following nest failure in Washington, consequently overall breeding success was 61% (Schroeder 1997). Eberhardt and Hofmann (1991) reported a nesting-success rate of 38% on the Yakima Training Center. Success rates were 39% (Nelson 1955) and 15% (Crawford et al. 1992) in Oregon, 61% (Wakkinen 1990) and 52% (Connelly et al. 1993) in Idaho, and 64% (Wallestad and Pyrah 1974) in Montana. Wallestad and Pyrah (1974) and Autenrieth (1981) reported greater productivity in female adults than juveniles; however, Connelly et al. (1993) found no difference in nest success or re-nesting success among age class.

Brood-rearing. After hatching, chicks wait until they are dry, then they leave the nest. Sage grouse hens attempt to raise one brood in a season (Girard 1937). Hens spend considerable time keeping chicks warm and guarding them for the first 4 to 5 weeks, but chicks feed themselves (Patterson 1952). Chicks remain with hens until late summer or early fall, then they congregate with other sage grouse in winter flocks. Brood size depends on nest success and chick survival. Average brood size was 4.6 chicks/hen in Washington (Pedersen 1982), 2.3 to 4 chicks/hen in Nevada (Zunino 1987), and 6.1 chicks/hen in Wyoming (Girard 1937). Forty-five to 75 days after eggs hatched in Washington, Schroeder (1997) observed broods with 3.8 chicks/hen. However, this may not accurately reflect brood size since some chicks leave broods within 60 days after hatching (Dunn and Braun 1986).

Juveniles. Sage grouse are considered juveniles when they lose their down. Juveniles become independent 10 to 12 weeks after hatching. Juveniles weigh the same as adults after 6 or 7 months (Patterson 1952). Broods separate by late August and early September, and juveniles join adult sage grouse in flocks. Juvenile males become sexually mature in April and May (Patterson 1950) and although they display on leks, they seldom mate (Dalke et al. 1963, Patterson 1950).

Mortality

Predation. Autenrieth (1986) reported nest predation was the most important constraint of population growth, followed by chick and adult predation. Habitat quality, specifically the amount and type of vegetation available to hens when building nests, ultimately determines the number of nests destroyed by predators (Gregg et al. 1994). Approximately 50% of sage grouse mortality is caused by predators (Bergerud 1988). Predation during winter depends on the availability of alternative prey for predators, such as cottontail rabbits (*Sylvilagus* sp.) and jackrabbits (*Lepus* sp.) (Autenrieth 1986).

Raptors and crows are the primary avian predators of sage grouse (Patterson 1952, Lumsden 1968, Wiley 1973). In Montana, ground predators were seen less frequently than aerial predators but killed more grouse (Hartzler 1974). Ground predators include coyotes (*Canis latrans*), bobcats (*Lynx rufus*), minks (*Mustela vison*), badgers (*Taxidea taxus*), and ground squirrels (*Spermophilus* sp.). Schroeder (1994) found ravens, coyotes, badgers, and other small mammals preyed on sage grouse eggs and were primarily responsible for nest failure in north central Washington. On the Yakima Training Center in Washington, four instances of sage grouse predation were documented: three by coyotes and one by an avian predator (Eberhardt and Hofmann 1991).

Hunting. Historically, the sage grouse was the leading game bird in 9 of the 15 western states it occupied (Girard 1937). By 1937, sage grouse was the leading game bird in three western states and of minor importance in a fourth state (Girard 1937, Rasmussen and Griner 1938, Patterson 1952).

Studies of the effect of hunting on sage grouse populations conflict. Leopold (1933) believed unregulated hunting could reduce any wildlife population to a level that was unstable. However, regulated harvest likely has little effect on stability in healthy populations (Autenrieth et al. 1982), unless there is no harvest limit or the season extends from July to October (Patterson 1952). In Oregon, there was no correlation between the number of sage grouse killed during hunting seasons and the size of the spring population (Crawford 1982). In Colorado, Braun and Hoffman (1979) believed hunting affected sage grouse stability only when $\geq 30\%$ of the population was killed. In Nevada, regulated hunting had no effect on the stability of a sage grouse population at low density (Zunino 1987). In 10 studies involving 8 species of grouse, Bergerud (1988) reported that hunting increased annual mortality by adding to, rather than compensating for, mortality during winter. Most of the studies cited by Bergerud (1988) reported a harvest of $>30\%$ of the population, which may have caused the additive effect.

In Washington, Yocom (1956) believed healthy populations of sage grouse annually produced more birds than were needed to sustain the population. Surplus sage grouse died from predation or exposure to inclement weather if they were not harvested, and protecting habitat, rather than eliminating hunting, insured the survival of the population (Yocom 1956).

Weather. Weather influences nesting success and survival of young chicks (Rasmussen and Griner 1938, Crawford 1960, Schlatterer 1960, Gill 1966, Rothenmaier 1979). However, in Montana, Wallestad and Watts (1972) found no correlations between productivity of sage grouse and rainfall or temperature. Weather affected nest success in Idaho (Dalke et al. 1963, Autenreith 1981), but its impact depended on the availability of forbs and insects for broods immediately following hatch (Autenreith 1981). In Colorado, Gill (1966) reported good sage grouse production when mean average temperature in spring exceeded 45° F and total precipitation was ≥5 cm (2 in). In Wyoming, Patterson (1952) found no nest failure resulting from low temperatures or snow but chicks apparently died from several consecutive days of cold rain, sleet, and snow accompanied by low temperatures. It appears adult sage grouse endure the winter reasonably well, provided wintering habitat contains adequate amounts of suitable sagebrush (Patterson 1952).

The effect of weather on habitat quality is a possible explanation for the population fluctuations exhibited by sage grouse (M. Schroeder, pers.comm.). Bradbury et al. (1989a) observed a decline in male lek attendance during the breeding season in response to bad weather conditions. Precipitation and wind during the morning display correlated with reduced male attendance, and low average daily temperatures on the day prior to lek attendance correlated with reduced male attendance on the display day. Declines in male lek attendance could result in decreased production rates for the season (Bradbury et al. 1989a).

Disease. Coccidiosis is the most prevalent disease affecting sage grouse (Simon 1940, Thorne 1969), although it is not continuously epidemic (Honest 1947). Coccidiosis is transmitted by the single-celled *Eimeria angusta* and *E. centroceri* in contaminated water and is more noticeable near drying water holes (Simon 1940). Maggots and beetles that feed on sage grouse droppings, and then are consumed by sage grouse, are responsible for disease transmission (Grover 1944).

Accidents and pesticides. In Idaho, Bean (1941) reported 11 male sage grouse killed by automobiles. Bean noted sage grouse use roads to dust which may make them vulnerable to vehicular accidents. In Washington, M. Schroeder (pers. comm.) reported one radio-marked female and two other females killed by automobiles in 1993. In Idaho, sage grouse have also been observed flying into windows and powerlines (J. Connelly, pers. comm.).

Pesticides may also directly kill sage grouse. Blus et al. (1989) found organophosphorus insecticides (dimethoate or methamidophos) directly responsible for the death of sage grouse occupying or being near sprayed fields in southeastern Idaho.

POPULATION STATUS

Past

Sage grouse numbers declined from the late 1800's to the early 1900's because of habitat conversion, grazing, and unrestricted hunting (Yocom 1956). In the 1920's and 1930's, grazing restrictions and the change from horse-plow to tractor farming reduced grazing by horses and allowed some recovery of rangeland (Yocom 1956). This resulted in more grouse during the 1940's and 1950's (Yocom 1956). However, the population likely remained depressed in comparison to historic descriptions.

Harvest. When the Washington Department of Game (WDFW) was created in 1933, a moratorium was placed on sage grouse hunting. In 1950, a 2-day sage grouse hunting season was re-opened in the Badger Pocket area of Kittitas County, with a daily and possession limit of one bird. The 1950 hunt resulted in the harvest of an estimated 2,700 sage grouse (Table 2). From 1951 to 1973, the season varied from 2 to 11 days throughout eastern Washington, with daily and possession limits of one or two. Due to declining harvest and lek counts after 1973, the season was shortened to 2 days with a daily limit of one and possession limit of two. Possession limits for hunting were further reduced to one in 1977 and the area open for hunting was reduced in 1978. Continuing declines in the sage grouse population and the lack of sufficient biological information resulted in a statewide harvest closure in 1988. Harvest figures show a decline in the number of sage grouse harvested from 1951 to 1987. In just 7 years, between 1974 and 1980, hunter success was cut nearly in half, from 0.43 grouse/hunter to 0.23 grouse/hunter. During this time, the hunting season was primarily 2 days/year (Table 2).

Harvest estimates were derived from the Washington Department of Wildlife (WDFW) Annual Game Harvest Questionnaire and wing envelopes (Pedersen 1982). Autenrieth et al. (1981) discussed the inadequacy of harvest questionnaires that sampled <25 to 30% of all hunters, which tend to overestimate harvest. Overestimates of $\geq 100\%$ may result when sampling 1 to 10% of a state's hunters (Pedersen 1982), so Washington sampling may not have been adequate. Furthermore, season restrictions could be partly responsible for harvest declines. Despite sampling inadequacies and changes in seasons, a declining trend in the sage grouse harvest in Washington from 1950 to 1987 is apparent.

Table 2. Hunting seasons and harvests of sage grouse in Washington, 1950-1987^a.

Year	Season length (days)	Counties open for harvest ^b	Harvested grouse ^c	Number of hunters ^d
1950	2	K	2,700	na ^c
1951	2	D, G, K	1,625	na
1952	2	D, G, K	2,500	na
1953	2	K, Y	2,550	na
	8	D, G		
1954	2	K, Y	2,375	na
	8	D, G		
1955	2	B, K, Y	2,275	na
	10	D, G		
1956	2	B, K, Y	1,550	na
	8	D, G		
1957	0	Closed Season	na	na
1958	2	D, G	1,400	na
1959	2	B, K, Y	1,450	na
	8	D, G, L		
1960	2	B, K, Y	2,700	na
	8	D, G, L		
1961	2	B, K, Y	2,175	na
	10	D, G, L		
1962	8	B, D, G, K, L, Y	1,600	na
1963	8	B, D, G, K, L, Y	1,900	na
1964	8	B, D, G, K, L, Y	1,750	na
1965	8	Eastern Washington	1,650	na
1966	8	Eastern Washington	1,750	na
1967	8	Eastern Washington	1,750	na
1968	8	Eastern Washington	1,400	na
1969	8	Eastern Washington	1,450	3,120
1970	8	Eastern Washington	2,065	3,300
1971	8	Eastern Washington	1,250	2,500
1972	8	Eastern Washington	1,250	2,700
1973	8	Eastern Washington	1,275	2,200
1974	2	Eastern Washington	900	2,100
1975	4	Eastern Washington	875	2,000
1976	2	Eastern Washington	865	2,200
1977	2	Eastern Washington	500	1,700
1978	2	E Wash., except A, F, G, O	410	2,000
1979	2	E Wash., except A, F, G, O	740	1,500
1980	2	D, K, L, Y	460	2,000
1981	2	D, K, L, Y	680	na
1982	2	D, K, L, Y	580	na
1983	2	D, K, L, Y	871	na
1984	2	D, K, L, Y	147	na
1985	2	K, Y	132	na
1986	2	K, Y	81	na
1987	2	K, Y (on Yakima Training Center)	18	na

^a Information obtained from past WDFW hunting pamphlets and departmental records.

^b A=Adams, B=Benton, D=Douglas, F=Franklin, G=Grant, K=Kittitas, L=Lincoln, O=Okanogan, Y=Yakima.

^c Harvest and hunter estimates acquired from the WDFW Annual Game Harvest Questionnaire and WDFW license sales (Pedersen 1982, WDFW unpubl. data). Approximately 10% of all licensed Washington hunters were sent questionnaires from 1967 to 1980, 5% from 1958 to 1966, and 3% from 1957 to 1958. Responses, based on the total number of licensed Washington hunters, averaged 6.0% from 1967 to 1980. Numbers of grouse harvested and hunters were adjusted using a 26% error factor that had previously been reported for sage grouse questionnaire responses in Washington (Zeigler 1978).

^d Number of sage grouse hunters were not tallied separately from other grouse hunters prior to 1969 or after 1980.

^e na=data not available

Statewide population. The average number of males per lek indicates trends and is useful in monitoring local populations. The total number of leks used by sage grouse can remain the same even though the population is declining. For these reasons, it is important for biologists to count the number of males at each lek rather than simply recording whether the lek was active or not. A decline in the number of males per lek likely indicates a declining population.

Possible biases in lek count data include: 1) new leks found over time may be smaller than previously located leks, 2) smaller leks may be less likely to be monitored, 3) the maximum number of males located may increase with increasing survey effort, and 4) the same sample of leks are not necessarily monitored in consecutive years.

Lek counts performed since 1955 show annual changes in the population (Fig. 3). Population fluctuations may result from changes in weather or other factors. Sage grouse populations fluctuate on a 10-year cycle, based on lek counts and harvest data from several western states (Rich 1985). In Washington (Wash. Dept. Fish and Wildl. 1995) and Oregon (Willis et al. 1993), state and county lek counts peaked approximately every 10 years. Surveys indicate that sage grouse subpopulations can decline to 20 to 25% of previous counts several years following peak counts.




Figure 3. The proportional change in sage grouse numbers estimated from annual lek counts, 1955-1997.

Efforts to perform standardized lek counts increased in 1970. The total number of males counted statewide increased after 1970. This was in large part because biologists included more leks in lek counts. However, increased frequency of lek counts, standardized counting procedures after

1970, and the discovery of satellite leks likely influenced trends. Although biologists visited more leks after 1969, the number of males per active lek continued to decline in Washington (Fig. 4). Males per active lek declined statewide from 35 males/lek in 1970 to 14 males/lek in 1994, then increased to 16 males/lek in 1997 (Fig. 4).




Figure 4. The ratio of males per lek and the number of leks included in counts, 1954-1997.

To roughly estimate historic population size and rate of decline, we looked at counts of active leks surveyed only in consecutive years. We used only consecutive year counts because frequently-monitored leks are the best indicators of trends in populations. Rate of decline was calculated from the annual change in mean number of birds per lek (Fig. 5). The estimated population declined an annual average of 3.4% between 1970 and 1997.

Past population estimates are also based upon back-calculation, following the estimated percent annual population change in lek counts. Overall, the population appears to have declined from a high of approximately 3,800 in 1970 to about 1,000 in 1997 (Fig. 6). Following this analysis back to 1954, an estimated population would be roughly 9,000 birds. Although lek counts have been conducted since 1954, the data since 1970 are likely best due to an increased number of leks counted and standardized counting procedures implemented that year. This analysis has inherent sources of bias, is limited by the lack of complete historical survey information, and therefore numbers should be considered rough estimates. The principal assumption is that changes in historic lek counts reflect changes in population size.




Figure 5. Male sage grouse counted statewide during lek counts in Washington, 1954-1997. Statewide counts are influenced by the number of leks included in the count.




Figure 6. Estimated current and historic population size of sage grouse based upon annual proportional changes in lek counts.

Regional populations. Sage grouse in Lincoln County illustrate how rapidly populations can disappear. Lek counts declined from 79 males in 1981 to 0 in 1987 (Fig. 7). The cause of the decline is unknown, but may be related to continued habitat removal (J. Hickman, pers. comm.) and degradation over time. Although no current leks are known, at least one sighting of a sage grouse brood was reported in 1993, 1994, and 1995.




Figure 7. Lek counts in Lincoln County, 1970-1997.

The last peak in the Grant and Douglas county population was in 1988; since that time the population declined to a low of 110 males in 1993 before rebounding to 235 in 1997 (Fig. 8). Some of the recent increase may be from newly discovered leks. Thirteen leks were active in 1995, up from 11 between 1988 and 1992. The average number of males per active lek increased from a low of 10.5 in 1994 to 18 in 1997, but is still below levels that occurred between 1988 and 1992 (28 males/lek in 1990, for example).

Over the past 10 years, habitat conditions in Douglas County have likely improved for sage grouse (M. Schroeder, pers comm.), largely as a result of the Conservation Reserve Program on private lands. Although sage grouse numbers have increased in Douglas County since 1992, it is too early to tell whether this reflects the improving underlying habitat conditions or represents the fluctuating patterns of population change indicated from past years (Fig. 8) Populations have increased significantly in the past, and then plummeted.




Figure 8. Lek counts in Douglas County, 1961-1997.

Yakima, Benton, and Kittitas county hunters took approximately 900 sage grouse annually between 1967 and 1972. Limited lek counts began in 1971. These intensified to encompass all known leks on the U.S. Army Yakima Training Center (YTC) in 1989. With more intensive survey effort conducted on the YTC since 1989, use of maximum counts from leks likely results in some double counting of males. Because wildlife staff at the YTC count all leks during a single day, high counts of all leks on any single day may alleviate some sampling bias, and provides a more conservative population estimate.

A trend toward a greater number of active leks with fewer males per lek is evident from survey data on the YTC. Numbers of males counted since 1989 declined from a high of 168 in 1991 to 76 in 1996, and increased to 158 in 1997 (S. Kruger, pers. comm.). Similarly, the average number of males per lek declined from 28 in 1991 to 7.6 in 1996, then increased to 14 in 1997 (S. Kruger, pers. comm.). The total number of active leks has increased from 5 in 1989 to 11 in 1997. The population on the YTC is the only remaining known population in Yakima and Kittitas counties.

The sage grouse population on the Fitzner and Eberhardt Arid Lands Ecology Reserve at Hanford (Hanford Site) in Benton County has been extirpated because of habitat degradation, development, powerline construction, and wildfires (C. Braun, letter dated 4 March 1992 to D. Ware; L. Fitzner pers. comm.). It is unknown when grouse were abundant at the Hanford site, but there were few grouse present after 1978 (L. Cadwell, pers. comm.). Three historic leks are

recorded from Benton County. One, near the southern edge of the Hanford reservation, was last known to be active in 1991.

A viable population of sage grouse no longer inhabits the Badger Pocket area of Kittitas County. This area, adjacent to the YTC, was known to support large numbers of sage grouse historically (L. Stream, pers. comm.). Sage grouse persisted at the site until 1987. During this time much of the native shrub steppe habitat was converted to agriculture.

In summary, the sage grouse population has declined substantially. This assessment is based on historical measures of abundance, pressures resulting from an annual average harvest of 1,842 grouse between 1951 and 1973, a decline in harvest from 1974 to 1987 (averaging 500 grouse annually), a corresponding steep decline in hunter success, a decline in the number of males per lek, the absence of grouse at traditional leks in Lincoln County, and a range reduction of approximately 85-90%.

Present

Currently two subpopulations of sage grouse remain in Washington; one on the Yakima Training Center in Yakima and Kittitas counties and one in Douglas and Grant counties (Fig. 2). The two populations appear to be separated by approximately 13 miles, which includes the Columbia River and lands dominated by orchards and croplands. While unlikely, it is possible that there may be interchange between these populations.

The 1997 breeding population of sage grouse in Washington has been estimated through lek counts and a population model. During spring surveys, 393 grouse were counted on 24 leks in 4 counties.

A model based on observations of the WDFW field researcher (M. Schroeder, pers. comm.), input and survey data from WDFW biologists, review of research outside Washington, and peer review by other sage grouse researchers was used to estimate the size of the 1997 breeding population (Table 3). The model assumed all leks were known and surveyed, all males were on leks during counts, and the male to female sex ratio was 1:6.

Table 3. Estimated size of the Washington sharp-tailed grouse breeding population in 1997 (based on preliminary data, September 1997).

Sex	Population estimate	Estimate source	Reference
Male	393	Statewide lek counts	WDFW unpubl. data
Female	629	1:1.6 sex ratio	Schroeder, pers. comm.
Total	1022	Males + Females	

This model would underestimate actual population size if some leks were not located, if all males were not on leks during counts, if the sex ratio was not 1:1, and if surveys were flawed (e.g., bad weather, incomplete counts, etc.). The model would overestimate actual population size if lek counts included females or if individual males were counted at more than one lek.

Double-counting of males on the Yakima Training Center is a concern. Estimates of approximately 312 - 411 sage grouse occur on the Yakima Training Center (YTC) in Kittitas and Yakima counties, based on 1997 lek counts (Kruger, pers. comm.). The lower estimate represents counts from the highest single day (as all leks on the YTC are counted in one day). The higher estimate represents the maximum number of grouse counted on any day. The higher estimate is reflected in Table 3.

Washington's sage grouse breeding population is estimated to be 923 - 1022 birds in 1997. Populations and resulting estimates will fluctuate from year to year. Approximately 611 of these birds are located in Douglas County and 312 - 411 birds are estimated on the Yakima Training Center.

HABITAT STATUS

Past

Much of Washington's shrub steppe has been converted to agriculture, converted to rangeland for livestock, or overgrazed, all of which reduce or degrade sage grouse habitat. Most shrub steppe has been sprayed, plowed, mechanically treated, burned, cut, or flattened to grow crops or forage for livestock (Table 4). Before settlement, much of eastern Washington was covered with large tracts of sagebrush/bunchgrass vegetation (Daubenmire 1988). Sagebrush coverage ranged from 5 to 26% and perennial grass coverage ranged from 69 to 100% on undisturbed sites (Daubenmire 1988).

Few ungulates grazed these areas since the last glaciation (Mack and Thompson 1982, Daubenmire 1988), and Native Americans seldom burned shrub steppe (Daubenmire 1988). Therefore, large tracts of sagebrush/bunchgrass vegetation evolved without intense livestock grazing and without frequent, intentional burning. Grazing of cattle, sheep, and horses by settlers and Native Americans began the era of rangeland degradation. Horses, which were introduced to Native Americans around 1730, were the first to intensively graze eastern Washington in recent history (Harris and Chaney 1984). Cattlemen were the first settlers in the Palouse region; they introduced cattle in 1834, sheep in the 1880's, and expanded wild-horse herds from 1830 to 1880 (Daubenmire 1988). Where shrub-steppe vegetation was grazed excessively by domestic animals, the density and canopy cover of native grasses was reduced, which allowed non-native species to invade (Daubenmire 1988).

The Homestead Act of 1862 led to the proliferation of small farms in eastern Washington between 1863 and 1910 (Harris and Chaney 1984). Plowing and burning of shrub steppe for agricultural expansion were widespread (Yocom 1956). Most of the land conversion for dryland farming occurred from 1900 through the 1940's, and for irrigated farming after 1950. The development of dryland farming required large herds of horses, which grazed freely on rangelands when they were not being used for farming (Harris and Chaney 1984). The combination of agricultural expansion and horses used in farming operations caused the most serious damage and deterioration to eastern Washington's shrub steppe in the late 1800's and early 1900's (Harris and Chaney 1984). Initially, agriculture increased food and water supplies, which allowed sage grouse to expand their range to unused areas (Yocom 1956). The introduction of tractor farming in the 1920's and 1930's reduced the need for horses and allowed some recovery of rangeland, but it also increased the area under cultivation. The conversion of native habitat to cropland intensified, and sage grouse began to decline (Yocom 1956).

Large-scale reclamation projects in the 1950's and 1960's further reduced sage grouse habitat (Hofmann 1991). From 1947 to 1982, 301,500 ha (744,705 ac) of brush control were documented under the federal Agricultural Conservation Program (ACP) and the Columbia Basin Project in Washington (Pedersen 1982). This brush control included 88,393 ha (218,331 ac) of sagebrush chemically or mechanically controlled and 213,120 ha (526,406 ac) converted to facilities and irrigated cropland. Twenty percent (60,800 ha [150,176 ac]) of all brush control occurred in Douglas, Lincoln, Kittitas, and Yakima counties; these counties also contained the most sage grouse. Sagebrush may re-invade areas following treatment, but sagebrush control in eastern Washington is an on-going activity. Although significant, the amount of sagebrush removed under federal programs was small compared to sagebrush removed by private landowners for agriculture (Pederson 1982).

Present

Based upon habitat classification of Daubenmire (1988) and LANDSAT analysis by Dobler et al. (1996), approximately 40% of the estimated 4.16 million ha (10.4 million acres) of shrub steppe in eastern Washington before European settlement remains. Additional shrub steppe exists in Okanogan County, but is outside of the range of sage grouse (Table 3). The remaining sage grouse range covers approximately 578,000 hectares (1,430,000 acres) (Table 4). Sage grouse habitat is a subset of the remaining shrub steppe; occupancy is dependent upon elevation, soil type, slope, size of shrub steppe area, and habitat quality. Three of the largest blocks of shrub-steppe occur on the Yakama Indian Nation, the Hanford Site in Benton County, and the U.S. Army Yakima Training Center in Yakima and Kittitas counties (Dobler et al. 1996). Of these, sage grouse remain only on the Yakima Training Center.

The largest contiguous block of shrub steppe habitat remaining in Washington is on the Yakima Training Center (YTC) in Yakima and Kittitas counties (Hofmann 1991). The YTC is used for military training exercises. Cadwell et al. (1994) found approximately 85% of all sage grouse use was within a L-shaped area in the southeast portion of the YTC, generally south and east of

Umtanum Ridge. Grouse generally used elevations between 500 and 900 m (1650-300 ft) on slopes less than 16° (Cadwell et al. 1996b). Sage grouse used a variety of shrub-steppe habitats, but tended to use areas with greater cover of sagebrush more frequently, especially for nesting (Sveum 1995, Cadwell et al. 1996b).

Another large parcel of shrub-steppe is located on the Department of Energy Hanford site, an area encompassing 560 sq. miles north of Richland in Benton County. Historic fires have decreased the abundance of sagebrush on the Hanford site, and likely contributed to the loss of sage grouse from Benton County. Currently, shrub-steppe at Hanford is in 3 general conditions; mature sagebrush, recovering shrub steppe (grass-dominated), and recovering agricultural lands (Dept. Of Energy 1996).

Table 3. Historical and remaining shrub steppe (ha) habitat in Washington counties within the historic range of sage grouse (Dobler et al. 1996)^a.

County	Historical	Remaining	% Loss
Adams	474,960	111,903	76
Benton	412,875	201,009	51
Chelan	80,770	30,761	62
Douglas	438,006	201,084	54
Franklin	301,486	92,311	69
Grant	645,822	228,732	65
Kittitas	232,466	129,578	44
Lincoln	504,013	189,470	62
Okanogan ^b	172,998	106,520	38
Walla Walla	308,007	71,215	77
Yakima	595,469	343,092	42
Total	4,166,870	1,705,674	59

^a Values based on LANDSAT data analyzed by Jim Eby at the WDFW Remote Sensing Laboratory using predictions of plant community distributions from Daubenmire (1988). Data is not available for the five other counties in the historical sage grouse range.

^b Analysis for only 20% of Okanogan County has been completed.

Land administration and ownership within the sage grouse range are based on the Washington Department of Natural Resources 1992 public lands survey and WDFW 1995 ownership/control data obtained from the WDFW Land Resources Program (Table 4). both owned and managed by federal or state agencies. Recent acquisitions or changes in management control since 1992 are not represented (except for WDFW), so current areas and percentages may differ.

The amount of land used primarily as farmland has remained relatively constant in the major counties that constitute sage grouse range. Cropland includes land harvested, land used for pasture or grazing, land used for cover crops, legumes, and soil improvement grasses, land where all crops failed, land used in cultivated summer fallow, and idle cropland.

The federal Conservation Reserve Program can benefit sage grouse by removing land from crop production, replacing those crops with perennial vegetation, and allowing sagebrush to invade. The CRP pays private landowners and public agencies that have highly-erodible cropland (usually dryland wheat) to establish grass for a minimum of 10 years. Payments to individual landowners or land managers are described in a signed contract, which specifies a date of termination. More than 160,000 hectares (395,000 acres) of cropland within the sage grouse range were enrolled in the CRP in the late 1980's. Douglas County enrolled the most at 60,000 hectares (150,000 acres) (Table 5).

Table 4. Administration and ownership of sage grouse range in Washington, 1992^a.

Agency/Ownership	Area (ha)	% of Total
Private	344,563	60
U.S. Army	104,624	18
Washington Department of Natural Resources	89,110	15
Washington Department of Fish and Wildlife	23,051	4
U.S. Bureau of Land Management	12,948	2
Washington State Parks and Recreation Commission	2,520	<1
U.S. Department of Energy	1,106	<1
Total	577,922	100

^a Ownership and administration tallies generated by the WDFW Geographical Information System Program based on WDNR public land surveys, January 1992.

Because CRP establishes permanent cover, it provides more year-round security to wildlife than land under cultivation. Sage grouse were observed displaying, feeding, roosting, and nesting in CRP grassland in eastern Washington (Schroeder 1994). M. Schroeder (pers. comm.) located 40% of nests in CRP fields in Douglas County from 1992 to 1996 (approximately 60 total nests were found), and these nests were as successful as ones built in other cover. The CRP fields that appear to be most important are those near small islands of shrub steppe (Schroeder 1994). These islands of shrub steppe are privately owned ranches with poor suitability for agricultural conversion and too small in area for efficient livestock use, but are selectively used by sage grouse (M. Schroeder, pers. comm.).

Sage grouse likely use CRP fields because the cover is continuous and can provide good nesting habitat. However, sage grouse select only quality CRP and typically avoid CRP fields in poor condition (M. Schroeder, pers. comm.). The quality of a CRP field depends on the type of vegetation planted and the length of time the field has been in CRP. In general, the longer a field is in CRP, the better its quality. In Douglas County, sagebrush has invaded many CRP fields (R. Friesz, pers. comm.), which may have increased habitat for sage grouse.

Table 5. Conservation Reserve Program (CRP) lands (ha) in eastern Washington counties in current sage grouse range through 1996^a.

County	Approximate county land area	Land enrolled in CRP	Percent of total CRP
Douglas	465,165	61,476	58
Grant	680,904	29,199	28
Yakima	1,097,398	14,333	14
Kittitas	590,917	836	<1
Total	2,825,384	105,844	100

^a Figures obtained from the U.S. Department of Agriculture, Agricultural Stabilization and Conservation Service, Spokane, Washington.

CONSERVATION STATUS

Legal Status

Sage grouse are listed as a game species in Washington by the WDFW, although the season has been closed since 1988. Sage grouse are designated a priority species and their habitat designated a priority habitat by the WDFW Priority Habitat and Species (PHS) Program.

Sage grouse are not protected under the federal Migratory Bird Treaty Act and jurisdiction is the state's responsibility. The U.S. Fish and Wildlife Service has placed the western sage grouse subspecies on its species of concern list.

Management Activities in Washington

Management plan. A statewide management plan for sage grouse was developed by the WDFW in 1995 (Wash. Dept. Fish and Wildl. 1995). The management plan establishes goals, objectives, and strategies for securing sage grouse populations in Washington. The management plan primarily directs WDFW activities and holds no authority over other public agencies. Specific tasks and objectives in the management plan such as habitat acquisition, enhancement, and monitoring are being implemented.

Population monitoring. The WDFW has conducted counts of sage grouse to assess population status and trends since the 1950's. Survey data were also used for establishing hunting seasons and bag limits. More intensive surveys were initiated in 1971 because of a recognized decline in the sage grouse population and the need to closely monitor the remaining population. The WDFW conducts lek counts and surveys for new leks each spring. Lek counts and searches for new leks are also conducted by the U.S. Army on the Yakima Training Center.

Winter surveys are conducted by the U.S. Bureau of Land Management (BLM) in cooperation with the WDFW in Lincoln County on sites acquired by the BLM. The reliability and feasibility of winter surveys in other areas are being determined as part of a current WDFW research project.

Habitat restoration/acquisition. The WDFW is acquiring and restoring habitat for sage grouse in eastern Washington. Funding here comes from the Bonneville Power Administration (BPA) mitigation funds and the Washington Wildlife and Recreation Program (WWRP). The Bonneville Power Administration (BPA) is obligated to mitigate for habitat and wildlife, including sage grouse, that were impacted during the construction and operation of federal dams on the Columbia River. The following criteria are used to prioritize acquisition areas for sage grouse:

- ▶ Areas of high-quality shrub steppe currently occupied by sage grouse
- ▶ Overlapping leks and winter-use areas on remaining shrub steppe
- ▶ Key wintering areas
- ▶ Shrub steppe ≤ 8 km (5 mi) from active leks
- ▶ Areas supporting many shrub-steppe obligates including sage grouse
- ▶ Historic use areas and travel corridors

The WDFW is restoring sagebrush on the Wells Wildlife Area in northern Douglas County and planted 30,000 sagebrush seedlings on the Hanford Site in Benton County during 1995. Lands purchased by the WDFW in Douglas and Lincoln counties that are designated Wildlife Areas are also being enhanced for sage grouse, specifically through grass and forb seeding and planting of shrubs and trees. WDFW Wildlife Areas that contain high quality shrub steppe can potentially be used for reintroduction/augmentation projects in the future. The WDFW works with landowners and the Natural Resources Conservation Service to promote and extend CRP contracts. Landowners are also encouraged to plant native forbs, grasses, and sagebrush on CRP land.

Research. The WDFW began a project entitled, "Productivity and Habitat Use of Sage Grouse in North-central Washington" in 1992 and it is still in progress. This project targets the sage grouse population centered in Douglas County. Current sage grouse research is focused on habitat-use assessment, population status, and identifying mortality rates and population recruitment. An index of habitat suitability is also being developed to refine habitat objectives and evaluate acquisition, augmentation, and reintroduction. Another WDFW study by Pederson (1982) evaluated distribution, home range and seasonal habitat use of sage grouse primarily in Douglas County.

The Battelle Memorial Institute Pacific Northwest Laboratory has been studying sage grouse on the Yakima Training Center (YTC) since 1989. In addition, since 1994, the sage grouse lek searches, lek counts, and population studies have been undertaken by the wildlife staff of the Directorate of Environment and Natural Resources (DENR) of the YTC. Research includes investigating seasonal habitat use, lek locations and annual use, population levels, survey techniques, and management techniques. Hoffman (1991) examined habitat selection at the

landscape scale on the YTC. Sveum (1995) examined sage grouse nesting and brooding habitat selection on the YTC (Sveum 1995).

Coordination and partnership. A conservation agreement was developed with the U.S. Army for the protection of sage grouse and their habitat on the Yakima Training Center. The agreement focuses on fire, military training, maintenance and construction of roads and facilities, stabilization of soil using exotic vegetation, grazing, invasion of introduced knapweed (*Centaurea* sp.), training equipment, hunting of other upland game birds and possible illegal harvest of sage grouse, disease and predation, restoration of disturbed sites, and potential changes in training levels or type of equipment used. Research guides management decisions on the training center and is used to solve conflicts between sage grouse and military training.

The WDFW coordinates with federal and other state agencies on habitat management and enhancement. Some public land that is administered by the U.S. Bureau of Reclamation and the U.S. Fish and Wildlife Service is managed by the WDFW. The WDFW currently works with the U.S. Department of Agriculture (through the NRCS and the Consolidated Farm Service Agency) on reducing the effect of brush control on sage grouse and other wildlife.

House Bill-1309 (Washington State Legislature 1993) requires the WDFW to develop goals to preserve, protect, and perpetuate fish and wildlife on state land used for agriculture, rangeland, or woodland used for grazing. The goals are submitted to the Washington State Conservation Commission, the legislative body responsible for coordinating the county conservation districts. A technical advisory committee appointed by the Conservation Commission has developed ecosystem standards for managing WDFW and WDNR land. These state agencies are required to implement practices on managed land to help achieve the standards for the site.

Enforcement. Illegal harvest of sage grouse is insignificant in Washington at this time, based on reports from WDFW enforcement agents. Furthermore, no documented mortality due to hunting or illegal harvest has been discovered during the current research project conducted by the WDFW in north-central Washington. Enforcement activities currently focus on discouraging harassment of sage grouse.

Information and education. The WDFW Priority Habitats and Species management recommendations for sage grouse are distributed to county planning departments, state and federal agencies, Native American tribes, and the public. The WDFW *Sage Grouse Fact Sheet* describes sage grouse and their habitat, life history, status, and management. Landowners are also encouraged to protect and enhance habitat for upland wildlife through the WDFW Upland Wildlife Restoration Program (UWRP). Copies of these materials, the management plan, and technical assistance are available through regional offices of the WDFW.

Habitat protection and enhancement. Sage grouse may also benefit from UWRP projects focused on other species. The UWRP was designed to work directly with private landowners on the protection and enhancement of private lands for upland wildlife. Although no projects specific

to sage grouse have been conducted, habitat enhancement for pygmy rabbits in Douglas County and for sharp-tailed grouse in Lincoln County is expected to benefit sage grouse.

FACTORS AFFECTING CONTINUED EXISTENCE

A variety of factors affect the continued existence of sage grouse in eastern Washington. General threats include broad scale habitat change and conversion, and management practices that adversely affect sagebrush habitat. Additional threats to habitat may also include livestock grazing and habitat loss caused by drifting herbicides applied during aerial application on wheat crops (Pederson 1982).

Primary threats to remaining sage grouse populations include potential for elimination or significant reduction of lands in the Conservation Reserve Program and the potential for large-scale fires eliminating large stands of sagebrush, especially at the Yakima Training Center. Additional threats of habitat reduction and disturbance occur from military training exercises at the Yakima Training Center. In Douglas County, continuation of the Conservation Reserve Program and protection of remnant patches of native habitat are the most critical needs for sage grouse. On the Yakima Training Center, fire prevention and training activity management are critical to maintaining sage grouse.

Population Isolation

Population isolation is potentially a significant factor influencing the continued existence of sage grouse in Washington. As grouse populations naturally fluctuate due to environmental conditions, the lower the population level, the greater the risk of extirpation. For example, the potential for compounded effects of habitat change are great when populations have dropped to low levels exhibited in 1988 and 1994.

Both sage grouse subpopulations have fluctuated to estimated lows of 100-150 females during the 1990's. Many authors indicate that long-term survival (greater than 100 years) of isolated populations may require many more individuals (Lande and Barrowclough 1987, Dawson et al. 1987, Grumbine 1990) although a needed effective population size for sage grouse has not been determined. The isolation of populations may have important ramifications for their genetic quality and recruitment (Lacy 1987) that may require human transport of individuals to counteract loss of fitness in small populations.

Potential for Continued Population Decline

Sage grouse populations in Washington have declined in virtually every decade this century, and their range has declined by approximately 85-90%. Neither of the two remaining populations is considered secure; significant threats remain in each case. Both populations exhibit relatively low

numbers of males at leks, which makes them vulnerable to predation, inclement weather, fire, and increased grazing pressure. Small reductions in habitat quality may have significant effects on lek maintenance. Without continued conservation effort to address the remaining threats, the sage grouse population in Washington will likely not increase substantially over current levels and may decline.

Over half of the remaining sage grouse occur in Douglas County. Sage grouse in Douglas County remain due to two primary factors: non-suitability of land for agriculture (which has resulted in the maintenance of remnant patches of shrub-steppe habitat) and the enrollment of private agricultural lands in the CRP. The CRP program has helped improve habitat conditions adjacent to remnant patches of shrub-steppe in Douglas County. Many of the island patches of shrub steppe have been maintained by private landowners for the past several decades. The principal difference between lands in Franklin County (where sage grouse were recently extirpated) and Douglas County is the presence of remnant habitat patches in Douglas County. Both areas have significant acreages enrolled in the CRP and have similar CRP habitats. The continued protection of leks, nesting grounds, and wintering sites in Douglas County, as well as continued monitoring are high priorities.

Compatibility with Military Training

The majority of sage grouse in Kittitas and Yakima counties are situated on the Yakima Training Center. The U.S. Army, U.S. Fish and Wildlife Service, and the WDFW have developed a cooperative agreement for the protection of sage grouse habitat on the training center. Key areas for sage grouse breeding, nesting, and wintering on the Yakima Training Center were identified (Eberhardt and Hofmann 1991, U.S. Army, unpubl. data). Year-round habitat for sage grouse occurs in the Lmumma Creek (formerly Squaw Creek) and Selah Creek watersheds. These areas contained the majority of active leks and nest sites (S. Kruger, pers. comm.) and are a priority for habitat and population protection. They do not, however, include all the areas used by sage grouse on the YTC. The U.S. Army has also initiated a comprehensive sage grouse research, monitoring, and habitat restoration program to mitigate impacts of training exercises.

The removal of sagebrush from the training center would adversely impact the sage grouse population in Kittitas and Yakima counties (Eberhardt and Hofmann 1991). Troop and vehicle activity on the training center has been shown to cause atypical and extensive movements of sage grouse and may be responsible for large home ranges (Eberhardt and Hofmann 1991). The prospect of sage grouse survival in Kittitas and Yakima counties in the future is questionable, depending on the maintenance of quality habitat on the Yakima Training Center and restriction of military activities in critical areas. However, because of increased training activity and potential fires, there is no guarantee that current efforts to conserve habitat by the U.S. Army will result in stable or increased populations of sage grouse.

Sagebrush levels on the Yakima Training Center are low when compared to studies from other states (Cadwell et al. 1996a). Sage grouse appear to use high quality habitat patches in a

fragmented environment. Sveum (1995) found that shrub cover at sage grouse nests on the YTC averaged 51% while random sites averaged 6%. A recent training exercise called Cascade Sage impacted approximately 14% of the big sagebrush occurring in the primary sage grouse habitat, causing death of 1.7% of sage grouse plants, and major structural damage to 7.8% (Cadwell et al. 1996a). Military vehicle use near active sage grouse leks was similar to that elsewhere. The removal of sagebrush from the exercise likely had some negative impact on the sage grouse population (Cadwell et al. 1996a). Future training exercises similar to Cascade Sage would likely result in additional negative impact (Cadwell et al. 1996a).

The actions taken by the U.S. Army are significant in the protection of sage grouse on the YTC. However, potential adverse impact from training exercises like Cascade Sage still threaten sage grouse inhabiting the YTC. Additionally, fires resulting from training exercises could have a devastating effect on the population.

Habitat Security

The declining distribution of sage grouse in Washington and likely the decline in sage grouse numbers appear to be related to the removal of shrub steppe. There has been a 85-90% reduction in the distribution of sage grouse in Washington. They survive in 4 of 16 counties they originally occurred in. The loss of shrub steppe in these counties ranges from 38 to 77% (Table 3).

Sage grouse may survive in Washington primarily because the remaining sagebrush habitat occurs on lands unsuitable for farming. Examples include the scablands of Grant and Douglas counties and the steep, rocky slopes of southern Kittitas and northern Yakima counties (Pedersen 1982). Future conversion of shrub steppe to agriculture is limited in Washington because the current sage grouse range is mostly unsuitable for farming (Pederson 1982). An important exception may be the Department of Energy's Hanford site. Significant portions of the Hanford Site may be converted to agricultural uses through transferral of management from federal to local authorities. Along with Lincoln County, the Hanford Site may hold Washington's greatest potential for expanding the current sage grouse population.

In Douglas County, however, where agriculture is the major land use, brush control and shrub steppe conversion continue. Remaining shrub steppe in Douglas County is believed to be vulnerable because of the extent of private lands and the intensity of land use for agriculture. Sagebrush has begun to invade many CRP lands, which may also result in better habitat conditions for sage grouse. Habitat improvement on CRP lands is thought to be a significant factor in the increasing population in north-central Washington. The CRP appears to benefit sage grouse by providing essential cover for nesting, cover that would otherwise be less available (M. Schroeder, pers.comm.). Beneficial CRP lands are those adjacent to remnant shrub-steppe patches.

The removal of CRP habitat in Grant and Douglas counties could cause further declines in sage grouse numbers. Thousands of acres currently enrolled in CRP may be placed back into production in the near future because contracts will expire; contracts representing approximately

318,000 ha (785,000 ac) are due to expire in 1997. CRP applications for approximately 332,000 ha (819,000 ac) were submitted by Washington farmers in spring 1997, but only 70,000 ha (172,000 ac) were accepted. The 21% acceptance rate was one of the lowest in the nation. Although landowners are reapplying during fall 1997, and we expect improvement in CRP enrollment, the future status of these areas is uncertain.

Relationships with Private Landowners

Sage grouse in Douglas County are entirely dependent upon private lands. A good working relationship between agencies trying to conserve sage grouse and private landowners needs to be maintained and developed. WDFW needs access to private lands, in some cases, to complete lek surveys and other research. Listing sage grouse may result in strained relationships with private landowners due to fears of regulation. It may also be a benefit to private landowners interested in the federal CRP program, as concerns for sage grouse habitat results in an increased likelihood of lands being enrolled in the CRP program.

Fire

Controversy exists regarding the frequency of fire in shrub steppe prior to settlement and increased grazing by livestock. Some authors believe fire maintained some bunchgrass-sagebrush ecosystems (Blaisdell et al. 1982). Braun (1987), however, contends this was possible only in mesic areas where silver and sand (*A. filifolia*) sagebrush dominated. C. Perry (pers. comm.) considers a fire-maintained sagebrush community in Washington unlikely, because the predominant species (big sagebrush) is fire intolerant. The fact that most sagebrush species are sensitive to fire and early explorers found sagebrush abundant in the sagebrush-grass region (Tisdale et al. 1969, Vale 1975) suggests that fire was infrequent (Tisdale and Hironaka 1981). Tisdale and Hironaka (1981) found it likely that fire was uncommon in the drier sagebrush types but more frequent in mesic communities where the fuel supply would be greater.

Following white settlement, the combination of fire and overgrazing greatly reduced the sagebrush and the understory species of grasses and palatable forbs (Tisdale and Hironaka 1981). The affected areas were then invaded by various aggressive, less-palatable species, especially introduced cheatgrass (Pickford 1932, Stewart and Hull 1949). The result was the domination of several million acres of the sagebrush-grass region by cheatgrass (Hironaka and Tisdale 1981). Cheatgrass is highly flammable, and its presence greatly increased the incidence of wildfire in the sagebrush-grass region (Klemmedson and Smith 1964, Kauffman 1990).

Harniss and Murray (1973) reported sagebrush only re-invades burned areas by seed, so 30 years or more may be required to regain pre-burn densities. Natural reseeding would also depend on a seed source in the general vicinity. Burning may also result in the invasion of noxious weeds (Autenrieth 1986) and introduced cheatgrass, which usually out-competes other grasses and forbs (C. Perry, pers. comm.; E. Bracken, pers. comm.). In Idaho and Nevada, wildfires converted large tracts of sagebrush to cheatgrass monocultures that were unsuitable as sage grouse habitat

(Drut 1994). In Idaho, Sime (1991) observed sage grouse using non-burned areas for cover and feeding during the winter and nesting period; vegetation in recovering burned areas provided overhead canopies for young grouse seeking palatable forbs during brood rearing.

Fire is a constant threat in the Yakima Training Center, where training activities occur. The mission of the Yakima Training Center is to train soldiers. Fire is an inevitable outcome of training activities. A fire on the YTC in August 1996 started in a military impact zone and burned over 15,000 ha (37,000 ac) on the YTC and approximately 5,000 ha (13,000 ac) on adjacent lands (M. Pounds, pers. comm.). Some areas known to be used by sage grouse were burned, but the fire missed critical lek and nesting habitats managed for sage grouse. The YTC is implementing a program to accelerate re-vegetation of burned and disturbed areas. Even though the training center has the highest intentions through a comprehensive fire management plan to control fires, fire remains a significant threat to the sage grouse population.

Chemical Treatment

Insecticides applied to agricultural and shrub steppe communities are often detrimental to sage grouse. Approximately 91,000 km² (35,000 mi²) of western rangelands were sprayed for grasshopper control from 1985 to 1990 (Johnson and Boyce 1990). Areas sprayed were commonly used by nesting sage grouse. Insects such as grasshoppers were found to be key food items in the diet of chicks (Rasmussen and Griner 1938, Patterson 1952, Klebenow and Gray 1968, Peterson 1970, Johnson and Boyce 1990), and chicks older than 3 weeks old show reduced growth rates when insects are removed from their diet (Johnson and Boyce 1990). Blus et al. (1989) also reported detrimental effects of organophosphorus insecticides (dimethoate and methamidophos) on sage grouse in southeastern Idaho.

Chemical treatment of vegetation has been found to reduce wintering, breeding, nesting, and brood-rearing activities of sage grouse (Enyeart 1956, Rogers 1964, June and Higby 1965, Kufeld 1968, Klebenow 1970, Martin 1970, Peterson 1970, Pyrah 1972, Wallestad 1975, Blus et al. 1989). However, Autenrieth (1969) found sprayed leks will continue to be used if the surrounding nesting and brood-rearing habitat has not been made uninhabitable by the removal of sagebrush and forbs. The herbicide 2,4-D is commonly used to reduce sagebrush and has been found to detrimentally impact sage grouse by altering the vegetational composition (Blaisdell and Mueggler 1956, Martin 1970, Autenrieth et al. 1982). Future efforts to control grasshoppers and sagebrush in Washington's shrub-steppe habitat may contribute to depressed sage grouse populations.

Grazing

Grazing has been identified as a factor in the decline in sage grouse numbers throughout their range. This decline coincided with the maximum livestock use of range resources between 1900 and 1915 (Patterson 1952). Ungulate pressure played no significant part in the evolution of ecotypes of the shrub steppe and steppe plants of Washington prior to the arrival of early settlers

(Mack and Thompson 1982, Daubenmire 1988). Human restriction of grazing animals confined them to sites subject to the vagaries of localized storms and periodic drought; the inability to determine grazing capacity led to either periodic or continuous overgrazing (Autenrieth et al. 1977). The result was an eventual decline in range production (Autenrieth et al. 1977). Yocom (1956) believed overgrazing during the cattle, sheep, and horse era may have had a depressive effect on sage grouse population levels.

The livestock era impacted habitat primarily through the reduction and alteration of native vegetation (Daubenmire 1988). Since most of the herbaceous species are more palatable than sagebrush, the former were reduced while the shrubs flourished (Tisdale and Hironaka 1981). Each period of overuse by domestic animals reduced the density of the large perennial grasses and forbs to a lower level than the preceding, and highly-adapted alien species (including noxious weeds) invaded. Lowered water tables through headcutting in meadow areas and erosion also resulted (Cottam and Stewart 1940, West 1983). Native shrub steppe was later converted to pastureland by livestock managers through burning, cutting, and spraying (Yocom 1956). By the 1930's, federal range personnel estimated that 84% of the sagebrush-grass region in the United States was severely depleted (USDA For. Serv. 1936).

Excessive grazing of sagebrush-bunchgrass communities that results in a low stubble and/or destruction of deciduous shrubs used for nesting and brood rearing can negatively impact sage grouse. Vegetative cover, specifically the provision of an understory of native grasses and forbs, contributes to sage grouse nesting success (Hein et al. 1980, Autenrieth 1981, Crawford and Delong 1993, Gregg et al. 1994). In Oregon, the availability of tall, dense grass cover and medium shrub cover decreased the likelihood of predation (Crawford and Delong 1993, Gregg et al. 1994) and land management practices that reduced herbaceous cover were discouraged (Crawford and Delong 1993). Grazing of tall grasses to <18 cm (7 in) would decrease their value for nest concealment (Gregg et al. 1994). The trampling of vegetation near riparian areas is detrimental to most upland wildlife. In general, valley bottoms and areas around water sources are used first by cattle; only after depletion of the vegetation at these mesic sites are cattle forced to use less-accessible, rougher terrain (Stoddart and Smith 1955). Upland meadows containing mesic sites (streams, springs, and associated meadow-type vegetation) provide critical habitat for sage grouse in arid regions (Savage 1969, Oakleaf 1971, Autenrieth et al. 1982).

Grazing was a serious concern on the Yakima Training Center until it was halted in 1995. The return of grazing in the future to the YTC could have significant adverse impacts to sage grouse. In addition, continued heavy grazing of shrub steppe habitats in Douglas and Grant counties will likely result in more degraded sage grouse habitats and depress populations further.

Harassment and Disturbance

The hunting season for sage grouse in Washington was closed in 1988; all current recreational use of sage grouse is viewing. Uncontrolled viewing could disrupt breeding populations and should be monitored and controlled if necessary. During the breeding season, daily disturbance at a lek

has the potential to reduce mating opportunities and cause decreased production. When humans approach the display site, grouse often flush and may or may not return again that day (Call 1979). Viewing at a distance from automobiles does not appear to disrupt courtship activity; but grouse flush when people leave cars to get a closer look. In Washington, viewing tours occur on the Yakima Training Center using strict guidelines; no reduction in lek attendance or disruption of breeding activities due to controlled public viewing has been reported (M. Pounds, pers. comm.).

Adequacy Of Existing Regulatory Mechanisms

Sage grouse individuals are protected from killing by Washington Department of Fish and Wildlife regulations; the Fish and Wildlife Commission closed the hunting of sage grouse in 1988. Populations have stayed at low levels or declined since then. There are no existing state or federal regulatory mechanisms to protect sage grouse habitat. The statewide management plan for sage grouse holds no authority over activities on public or private lands outside of those managed by the WDFW.

CONCLUSIONS AND RECOMMENDATION

The sage grouse population and corresponding sage grouse habitat have declined significantly in Washington. Sage grouse current range has declined to 10 - 15 percent of historic range. Population declines have occurred in the 1980's and 1970's as well as early in this century. Local populations were extirpated as recently as the mid-1980's. Only 2 populations, numbering between 900 and 1000 individuals remain in the state. Major threats to the two populations are potential for catastrophic fire, impacts of military training, agricultural conversion, potential for increased grazing, and the instability of the federal Conservation Reserve Program. Because of these factors, it is recommended that the sage grouse be designated a threatened species in Washington.

There are three principal reasons why the Department does not recommend "endangered" status for the sage grouse. These are: the relative size of the remaining populations; efforts by the Yakima Training Center to reduce impacts to sage grouse; and the existence of the Conservation Reserve Program. Some private lands critical to maintenance of sage grouse were enrolled during spring 1997. Other critical lands are expected to be enrolled during fall 1997.

REFERENCES CITED

- Aldrich, J.W. 1963. Geographic orientation of American Tetraonidae. *J. Wildl. Manage.* 27:529-545.
- Autenrieth, R. E., W. R. Brigham, W. Molini, P. Shields, J. Slosson, and M. Wickersham. 1977. Livestock and upland wildlife. Pages 76-86 in J. W. Menke, ed. Proc. of a workshop on livestock and wildlife, fisheries relationships in the Great Basin. Spec. Publ. 3301.
-))))). 1980. The sagebrush connection. *Id. Wildl.* 2:9-11.
-))))). 1981. Sage grouse management in Idaho. *Id. Dept. Fish and Game Wildl. Bull.* 9.
-))))), W. Molini, and C. Braun, editors. 1982. Sage grouse management practices. *West. States Sage Grouse Comm. Tech. Bull.* 1.
-))))). 1986. Sage grouse. Pages 762-779 in A. S. Eno, R. L. Di Silvestro, and W. J. Chandler, compilers. Audubon wildlife report 1986. *Natl. Audubon Soc., New York, N.Y.*
- Baird, S. F., T. M. Brewer, and R. Ridgway. 1874. *The land birds of North America.* Vol. 3. Little, Brown and Co., Boston, Mass.
- Barnett, J. K. and J. A. Crawford. 1993. Diet and nutrition of sage grouse hens during the pre-laying period in Oregon. *Oreg. Agric. Res. Stn. Rep.* 10-123.
- Batterson, W. M. and W. B. Morse. 1948. Oregon sage grouse. *Oreg. Game Comm. Fauna Ser.* 1.
- Bean, R. W. 1941. Life history studies of the sage grouse (*Centrocercus urophasianus*) in Clark County, Idaho. B.S. Thesis, Utah St. Agric. Coll., Logan.
- Beck, T. D. I. 1975. Attributes of a wintering population of sage grouse, North Park, Colorado. M.S. Thesis, Colorado State Univ., Fort Collins.
-))))). 1977. Sage grouse flock characteristics and habitat selection in winter. *J. Wildl. Manage.* 41:18-26.
-))))). and C. E. Braun. 1978. Weights of Colorado sage grouse. *Condor* 80:241-243.
- Bergerud, A. T. 1988. Population ecology of North American grouse. Pages 578-648 in A. T. Bergerud and M. W. Gratson, eds. *Adaptive strategies and population ecology of northern grouse.* Univ. Minnesota Press., Minneapolis.

- Berry, J. D. and R. L. Eng. 1985. Interseasonal movements and fidelity to seasonal use areas by female sage grouse. *J. Wildl. Manage.* 49:237-240.
- Blaisdell, J. P. and W. F. Mueggler. 1956. Effect of 2,4-D on forbs and shrubs associated with big sagebrush. *J. Range Manage.* 9:30-40.
- Blaisdell, J.P., R. B. Murry, and E. D. McArthur. 1982. Managing intermountain rangelands--sagebrush-grass ranges. U.S. Dept. Agric., Inter. For. and Range Exp. Sta. Gen. Tech. Rep. INT-134.
- Blus, L. J., C. S. Staley, C. J. Henny, G. W. Pendleton, T. H. Craig, E. H. Craig, and D. K. Halford. 1989. Effects of organophosphorus insecticides on sage grouse in southeastern Idaho. *J. Wildl. Manage.* 53:1139-1146.
- Bradbury, J. W., S. L. Vehrencamp, and R. M. Gibson. 1989a. Dispersion of displaying male sage grouse. Part I. Patterns of temporal variation. *Behav. Ecol. Sociobiol.* 24:1-14.
-))))), R.M. Gibson, C. E. McCarthy, and S.L. Vehrencamp . 1989b. Dispersion of displaying male sage grouse. Part II. The role of the female dispersion. *Behav. Ecol. Sociobiol.* 24:15-24.
- Braun, C. E. 1975. Mortality, survival and effects of hunting on grouse, partridge, pheasants and quail, an annotated bibliography. *Colo. Div. Wildl. Rep.* 3.
-)))) T. D. I. Beck. 1976. Effects of sagebrush control on distribution and abundance of sage grouse. *Fed. Aid Wildl. Restor. Proj. W-37-R, Work Plan 3, Job 8a.* Colo. Div. Wildl., Fort Collins.
-))))), T. Britt, and R. O. Wallestad. 1977. Guidelines for maintenance of sage grouse habitats. *Wildl. Soc. Bull.* 5:99-106.
-)))) and D. M. Hoffman. 1979. Vulnerability and population characteristics of sage grouse in Moffat County. *Fed. Aid Wildl. Restor. Proj. W-37-R-32.* Colo. Div. Wildl., Fort Collins.
-)))) . 1984. Attributes of a hunted sage grouse population in Colorado, USA. Pages 148-162 *in* P. J. Hudson and T. W. I. Lorel, eds. Third international grouse symposium. World Pheasant Assoc., York Univ., U.K.
-)))) . 1987. Current issues in sage grouse management. *Proc. western association fish and wildlife agencies.* Colo. Div. Wildl., Fort Collins.

-))))), compiler. 1991. Western states sage grouse and Columbian sharp-tailed grouse questionnaire survey, 1988-90. West. States Sage Grouse and Columbian Sharp-tailed Grouse Tech. Comm.
-))))). 1993. The status of sage grouse: Are they endangered, threatened, or ?. Abstract first joint meeting prairie grouse technical council and western states sage and Columbian sharp-tailed grouse workshop. Colo. Div. Wildl., Fort Collins.
- Burnett, L. E. 1905. The sage grouse, *Centrocercus urophasianus*. The Condor 7:102-105.
- Cadwell, L. L., M. A. Simmons, J. L. Downs, and C. M. Sveum. 1994. Sage grouse on the Yakima Training Center: A summary of studies conducted during 1991 and 1992. Pac. Northwest Lab., Richmond, Washington.
-))))),))))), W.R. Reid, and J.J. Nugent. 1996a. An analysis of sage grouse habitat change at the Yakima Training Center after the Cascade Sage Training Exercise. PNNL-10999 Pac. Northwest Lab., Richmond, Washington
-))))),))))), J.J. Nugent, and V.I. Cullinan. 1996b. Sage grouse habitat on the Yakima Training Center: Part II, Habitat Modeling. Pac. Northwest Lab., Richmond, Washington
- Call, M. W. 1979. Habitat requirements and management recommendations for sage grouse. U.S. Dept. Inter., Bur. Land Manage. Tech. Note 330.
-)))) and Maser, C. 1985. Wildlife habitats in managed rangelands, the Great Basin of southeastern Oregon: Sage grouse. U.S. Dept. Agric., Pacific Northwest For. and Range Exp. Stn., Gen. Tech. Rep. PNW-187.
- Carr, H. D. 1967. Effects of sagebrush spraying on abundance, distribution, and movements of sage grouse. M.S. Thesis, Colorado State Univ., Fort Collins.
- Clark, L. F., H. Rahn, and M. D. Martin. 1942. Sage grouse studies. Part II. Seasonal and sexual dimorphic variation in the so-called "air sacs" region of the sage grouse. Wyo. Game and Fish Comm. Bull. 2:13-27.
- Connelly, J. W. Jr. and O. D. Markham. 1983. Movements and radionuclide concentrations of sage grouse in southeastern Idaho. J. Wildl. Manage. 47:169-177.
-))))), H. W. Browsers, and R. J. Gates. 1988. Seasonal movements of sage grouse in southeastern Idaho. J. Wildl. Manage. 52:116-122.
-))))), W. L. Wakkinen, A. D. Apa, and K. P. Reese. 1991. Sage grouse use of nest sites in southeastern Idaho. J. Wildl. Manage. 55:521-524.

-))))), R. A. Fischer, A. D. Apa, K. P. Reese, and W. L. Wakkinen. 1993. Renesting by sage grouse in southeastern Idaho. *The Condor* 95:1041-1043.
- Cooper, J. G. 1860. Report upon the birds collected on the survey: Land Birds. Ch. 1., U.S. Pacific Railroad Explor. and Surv. 12. Book 2, Zoology. 3:140-226.
- Cottam, W. P. and G. Stewart. 1940. Plant succession as a result of grazing and of meadow desiccation by erosion since settlement in 1892. *J. For.* 38:613-626.
- Coues, E. 1893. Birds of the Northwest. A handbook of the ornithology of the region drained by the Missouri River and its tributaries. U.S. Dept. Inter., U.S. Geol. Surv. of Territories Misc. Publ. 3:400-407.
- Crabtree, R. L., L. S. Broome, and M. L. Wolfe. 1989. Effects of habitat characteristics on gadwall nest predation and nest-site selection. *J. Wildl. Manage.* 53:129-137.
- Crawford, J. A. 1982. Factors affecting sage grouse harvest in Oregon. *Wildl. Soc. Bull.* 10:374-377.
-))))), M. A. Gregg, M. S. Drut, and A. K. DeLong. 1992. Habitat use by female sage grouse during the breeding season in Oregon. Final rep. submitted to Bur. Land Manage., Oregon State Univ., Corvallis.
-)))) and A. K. DeLong. 1993. Relationships between vegetative structure and predation rates of artificial sage grouse nests. Final rep. submitted to Bur. Land Manage., Oregon State Univ., Corvallis.
- Dalke, P. D., D. B. Pyrah, D. C. Stanton, J. E. Crawford, and E. F. Schlatterer. 1960. Seasonal movements and breeding behavior of sage grouse in Idaho. *Trans. twenty-fifth North Am. Wildl. Conf.:*396-407.
-))))),))))),))))),))))), and))))). 1963. Ecology, productivity, and management of sage grouse in Idaho. *J. Wildl. Manage.* 27:811-841.
- Daubenmire, D. 1988. Steppe vegetation of Washington. Washington State Univ. Coop. Ext. Bull. EB 1446.
- Davis, N. B. 1978. Ecological questions about territorial behavior. Pages 317-350 in J. R. Krebs and N. B. Davies, eds. *Behavioral ecology*. Blackwell Sci. Publ., Oxford.
- Dobler, F. C. J. Eby, C. Perry, S. Richardson, and M. Vander Haegen. 1996. Status of Washington's shrub-steppe Ecosystem: Extent, ownership, and wildlife/vegetation relationships. *Wildl. Manage. Div., Wash. Dept. Fish and Wildl., Olympia*.

- Donoho, H. S. and J. Roberson, editors. 1985. Summary of sage grouse questionnaires, Volume I through XXXIII. West. States Sage Grouse and Columbian Sharp-tailed Grouse Tech. Comm.
- Drut, M. S. 1994. Status of sage grouse in North America with emphasis on populations in Oregon and Washington. Audubon Soc., Portland, Oregon.
- Dunn, P. O. and C. E. Braun. 1986. Late summer-spring movements of juvenile sage grouse. *Wilson Bull.* 98:83-92.
- Dunn, J. L., E. A. T. Blom, G. E. Watson, and J. P. O'Neill, consultants. 1987. Field guide to the birds of North America. Natl. Geogr. Soc., Wash. D.C.
- Eastman, D. S. and D. Jenkins. 1970. Comparative food habits of red grouse in northeast Scotland, using fecal analysis. *J. Wildl. Manage.* 34:612-620.
- Eberhardt, L. E. and L. A. Hofmann. 1991. Sage grouse on the Yakima Training Center: A summary of studies conducted during 1989 and 1990. Pac. Northwest Lab., Richland, Washington.
- Edminster, F. C. 1954. American game birds of field and forest. Charles Scribner's Sons, New York, N.Y.
- Ellis, K. L., J. R. Parrish, J. R. Murphy, and G. H. Richins. 1989. Habitat use by breeding male sage grouse: a management approach. *Great Basin Nat.* 49:404-407.
- Elman, R. 1974. The hunters field guide to the game birds and animals of North America. Alfred A. Knopf, New York, N.Y.
- Emmons, S. R. and C. E. Braun. 1984. Lek attendance of male sage grouse. *J. Wildl. Manage.* 48:1023-1028.
- Enyeart, G. W. 1956. Responses of sage grouse to grass reseeding in the Pines Area, Garfield County, Utah. M.S. Thesis, Utah State Univ., Logan.
- Eng, and P. Schladweiler. 1972. Sage grouse winter movements and habitat use in central Montana. *J. Wildl. Manage.* 36:141-146.
- Fischer, R. A., A. K. Apa, W. L. Wakkinen, K. P. Reese, and J. W. Connelly. 1993. Nesting-area fidelity of sage grouse in southeastern Idaho. *The Condor* 95:1038-1041.
- Franklin, J. F. and C. T. Dyrness. 1973. Natural vegetation of Oregon and Washington. U.S.D.A. For. Serv. Gen. Tech. Rep., PNW-8. 417 pp.

- Gates, R. J. 1983. Sage grouse, lagomorph, and pronghorn use of sagebrush grassland burn site on the Idaho National Engineering Laboratory. M.S. Thesis, Montana State Univ., Bozeman.
- Gill, R. B. 1965. Distribution and abundance of a population of sage grouse in North Park, Colorado. M.S. Thesis, Colorado State Univ., Fort Collins.
-))))). 1966. Weather and sage grouse productivity. Colo. Game, Fish and Parks Dept. Outdoor Facts 37.
- Girard, G. L. 1937. Life history, habits, and food of the sage grouse, *Centrocercus urophasianus* Bonaparte. Univ. Wyoming Publ. 3:1-56.
- Gregg, M. A. 1991. Habitat use and selection of nesting habitat by sage grouse in Oregon. M.S. Thesis, Oregon State Univ., Corvallis.
-)))))), J. A. Crawford, M. S. Drut, and A. K. DeLong. 1994. Vegetational cover and predation of sage grouse nests in Oregon. J. Wildl. Manage. 58:162-166.
- Griner, L. A. 1939. A study of the sage grouse, *Centrocercus urophasianus*, with special reference to life history, habitat requirements and numbers and distribution. M.S. Thesis, Utah State Agric. Coll., Logan.
- Grover, G. 1944. Disease among sage grouse. Colo. Conserv. Comments 7:14-15.
- Grumbine, R.E. 1990. Viable populations, reserve size, and federal lands management: a critique. Conservation Biology (4): 127-134.
- Guiquet, C. J. 1970. Birds of British Columbia. Queen's Printers, Victoria, British Columbia.
- Harris, G. A. and M. Chaney. 1984. Washington State grazing land assessment. Wash. Range Land Comm. and Washington State Univ. Coop. Ext. Ser., Pullman.
- Harrison, C. 1978. A field guide to the nests, eggs and nestlings of North American birds. The Stephen Green Press, Brattleboro, Vt.
- Hartzler, J. E. 1974. Predation and the daily timing of sage grouse leks. Auk 91:532-536.
-)))) and D. A. Jenni. 1988. Mate choice by female sage grouse. Pages 240-269 in A. T. Bergerud and M. W. Gratson, eds. Adaptive strategies and population ecology of northern grouse. Volume I: Population studies. Univ. Minnesota Press, Minneapolis.
- Hjorth, I. 1970. Reproductive behavior in tetraonidae with special reference to males. Viltrevy Swedish Wildl. 7:432-461.

- Hofmann, L. 1991. The western sage grouse (*Centrocercus urophasianus phaios*) on the Yakima Training Center in central Washington. M.S. Thesis, Central Washington Univ., Ellensburg.
- Honess, R. F. 1947. Coccidiosis as a decimating factor among sage grouse. Fed. Aid Wildl. Restor. Proj. 33-R. Wyo. Game and Fish Comm., Cheyenne.
-)))) and W. J. Allred. 1942. Sage grouse studies. Part I. Structure and function of the neck muscles in inflation and deflation of the esophagus in sage grouse. Wyo. Game and Fish Comm. Bull. 2:1-12.
- Huntington, D. W. 1897. The sage grouse. Osprey 2:17-18.
- Hupp, J. W. and C. E. Braun. 1989. Topographic distribution of sage grouse foraging in winter. J. Wildl. Manage. 53:823-829.
- Jarvis, J. M. 1974. Sage grouse population studies on the Parker Mountain in south central Utah. Fed. Aid Wildl. Restor. Proj. W-65-R, Job c-1. Ut. Dept. Nat. Resour., Div. Wildl. Resour., Salt Lake City.
- Jenkins, D., A. Watson, and G. R. Miller. 1963. Population studies on red grouse, *Lagopus lagopus scoticus* (Lath.), in north-east Scotland. J. Anim. Ecol. 32:317-376.
-)))) , A. Watson, and N. Picozzi. 1965. Red grouse chick survival in captivity and in the wild. Trans. sixth Congr. Int. Union Game Biol.: 63-70.
- Jewett, S. G., W. P. Taylor, W. T. Shaw, and J. W. Aldrich. 1953. Birds of Washington State. Univ. Washington Press, Seattle.
- Johnsgard, P. A. 1973. Grouse and quail of North America. Univ. Nebraska Press, Lincoln.
- Johnson, G. D. and M. S. Boyce. 1990. Feeding trials with insects in the diet of sage grouse chicks. J. Wildl. Manage. 54:89-91.
- June, J.W. and L. W. Higby. 1965. Proc. western states sage grouse 1965 workshop. West. States Sage Grouse Comm.
- Kauffman, J. B. 1990. The ecology of fire in rangelands: Historical and current contexts. Pages 2-6 in T. E. Bedell, ed. Proc. 1990 Pacific Northwest Range Management Short Course: Fire in Pacific Northwest Ecosystems. Dept. Rangeland Resour., Oregon State Univ., Corvallis.

- Klebenow, D. A. and G. M. Gray. 1968. Food habits of juvenile sage grouse. *J. Range Manage.* 21:80-83.
-))))). 1969. Sage grouse nesting and brood habitat in Idaho. *J. Wildl. Manage.* 33:649-662.
-))))). 1970. Sage grouse versus sagebrush control in Idaho. *J. Range Manage.* 23:396-400.
-))))). 1981. Livestock grazing interactions with sage grouse. Pages 113-123 *in* J. M. Peek and P. D. Dalke, eds. *Wildlife-livestock relationships symposium: Proceedings 10.* Univ. Idaho, For., Wildl. and Range Exp. Stn., Moscow.
-))))). 1985. Habitat management for sage grouse in Nevada. *World Pheasant Assoc.* 10:34-46.
- Klemmedson, J. O. and J. G. Smith. 1964. Cheatgrass (*Bromus tectorum* L.). *Bot. Rev.* 30:226-262.
- Klott, J. H. and F. G. Lindzey. 1989. Comparison of sage and sharp-tailed grouse leks in south central Wyoming. *Great Basin Nat.* 49:275-278.
-)))) and)))). 1990. Brood habitat of sympatric sage grouse and Columbian sharp-tailed grouse in Wyoming. *J. Wildl. Manage.* 54:84-88.
- Kufeld, R. C. 1968. Range type conversion programs in Colorado and their impact on deer, elk and sage grouse. Pages 173-186 *in* Proc. forty-eighth annual conference western association state game and fish commissions.
- Lacy, R.C. 1987. Loss of genetic diversity from managed populations: interacting effects of drift, mutation, immigration, selection, and population subdivision. *Conservation Biology* (2):143-158.
- Lande, R., and G.F. Barrowclough. 1987. Effective population size, genetic variation, and their use in population management. Pages 87-123 *in* M.E. Soule, ed. *Viable populations.* Cambridge University Press, New York.
- Leopold, A. 1933. *Game management.* Oxford Univ. Press, New York, N.Y.
- Lumsden, H. G. 1968. The displays of the sage grouse. *Ont. Dept. Lands and For. Res. Rep.* (Wildl.) 83.
- Mack, R. N. and J. N. Thompson. 1982. Evolution in steppe with few large, hooved mammals. *Am. Nat.* 119:757-773.

- Martin, N. S. 1970. Sagebrush control related to habitat and sage grouse occurrence. *J. Wildl. Manage.* 34:313-320.
- Moss, R., A. Watson, and R. Pam. 1975. Maternal nutrition and breeding success in red grouse (*Lagopus lagopus scoticus*). *J. Anim. Ecol.* 44:233-244.
- Nelson, O. C. 1955. A field study of the sage grouse in southeastern Oregon with special reference to reproduction and survival. M.S. Thesis, Oregon State Univ., Corvallis.
- Oakleaf, R. J. 1971. The relationship of sage grouse to upland meadows in Nevada. M.S. Thesis, Univ. Nevada, Reno.
- Patterson, R. L. 1950. The sage grouse in the Upper Green River Basin. Thirtieth ann. conf. west. assoc. state game and fish comm.
-))))). 1952. The sage grouse in Wyoming. Sage Books, Inc., Denver, Colo.
- Pedersen, W. T. 1982. Sage grouse status, distribution, movement, seasonal use of habitat, and habitat status in eastern Washington. Fed. Aid Wildl. Restor. Proj. W-70-R-21, Study VI, Job 1 and 2. Wash. Dept. Game, Olympia.
- Petersen, B. E. 1980. Evaluation of the effects of hunting regulations on sage grouse populations: evaluation of census of females. Fed. Aid Wildl. Restor. Proj. W-37-R-33. Colo. Div. Wildl., Denver.
- Peterson, J. G. 1970. The Food Habits and Summer Distribution of Juvenile Sage Grouse in Central Montana. *J. Wildl. Manage.* 34:147-155.
- Pickford, G. D. 1932. The influence of continued heavy grazing and of promiscuous burning on spring-fall ranges in Utah. *Ecology* 13:159-171.
- Pyrah, D. B. 1954. A preliminary study toward sage grouse management in Clark and Fremont counties based on seasonal movements. M.S. Thesis, Univ. Idaho, Moscow.
-))))). 1963. Sage grouse investigations. Fed. Aid Wildl. Restor. Proj. W-125-R-2, Id. Fish and Game Dept., Boise.
-))))). 1972. Effects of chemical and mechanical sagebrush control on sage grouse. Subject: Dropping counts. Pages 16-25 in *Ecological effects of chemical and mechanical sagebrush control*. Mont. Fish and Game Dept., Helena.

- Rasmussen, D. I. and L. A. Griner. 1938. Life history and management studies of the sage grouse in Utah, with special reference to nesting and feeding habits. Trans. North Am. Wildl. Conf. 3:852-864.
- Redmond, G. W., D. M. Keppie, and P. W. Herzog. 1982. Vegetative structure, concealment, and success at nests of two races of spruce grouse. Can. J. Zool. 60:670-675.
- Remington, T. E. 1983. Food selection, nutrition, and energy reserves of sage grouse during winter, North Park, Colorado. M.S. Thesis, Colorado State Univ., Fort Collins.
-)))) and C. E. Braun. 1985. Sage grouse food selection in winter, North Park, Colorado. J. Wildl. Manage. 49:1055-1061.
-)))) and)))). 1988. Carcass composition and energy reserves of sage grouse during winter. The Condor 90:15-19.
- Rich, T. 1985. Sage grouse population fluctuations: evidence for a 10-year cycle. U.S. Dept. Int., Bur. Land Manage., Id. State Off. Tech. Bull. 85-1.
- Roberson, J. A. 1984. Sage grouse-sagebrush relationships: A review. Biol. of *Artemisia* and *Chrysothamnus*, Provo, Utah.
- Rogers, G. E. 1964. Sage grouse investigations in Colorado. Fed. Aid Wildl. Restor. Proj. W-37-R, number 16. Colo. Game, Fish and Parks Dept., Denver.
- Rothenmaier, D. 1979. Sage grouse reproductive ecology: breeding season movements, strutting ground attendance and site characteristics, and nesting. M.S. Thesis, Univ. Wyoming, Laramie.
- Savage, D. E. 1969. The relationship of sage grouse to upland meadows in Nevada. Nev. Coop. Wildl. Res., Nev. Fish and Game Comm., and Nev. Agr. Exp. Stn., Reno.
- Schlatterer, E. F. 1960. Productivity and movements of a population of sage grouse in southeastern Idaho. M.S. Thesis, Univ. Idaho, Moscow. 87 pp.
- Schoenberg, T. J. 1982. Sage grouse movements and habitat selection in North Park, Colorado. M.S. Thesis, Colorado State Univ., Fort Collins.
- Schroeder, M.A. 1994. Progress report: Productivity and habitat use of sage grouse in north-central Washington. Washington Dept. of Wildlife (Unpubl.), Olympia, WA.
- Schroeder, M.A. 1997. Unusually high reproductive effort by sage grouse in a fragmented habitat in north-central Washington. Condor 99:933-941.

- Scott, J. W. 1942. Mating behavior of the sage grouse. *Auk* 59:477-498.
- Sime, C. A. 1991. Sage grouse use of burned, non-burned, and seeded vegetation on the Idaho National Engineering Laboratory, Idaho. M.S. Thesis, Montana State Univ., Bozeman.
- Simon, F. 1940. The parasites of the sage grouse. *Univ. Wyoming Publ.* 7:77-100.
- Sugden, L. G. and G. W. Beyersbergen. 1986. Effect of density and concealment on American crow predation of simulated duck nests. *J. Wildl. Manage.* 50:9-14.
-)))) and)))). 1987. Effect on nesting cover density on American crow predation of simulated duck nests. *J. Wildl. Manage.* 51:481-485.
- Thorne, E. T. 1969. Diseases in Wyoming sage grouse. *Proc. West. States Sage Grouse Workshop* 6:192-198.
- Tisdale, E. W. and M. Hironaka. 1981. The sagebrush-grass region: A review of the ecological literature. U.S. For. Serv., For. and Range Exp. Stn. Bull. 33.
- Trueblood, R. W. 1954. The effect of grass reseeding in sagebrush lands on sage grouse populations. M.S. Thesis, Utah State Agric. Coll., Logan.
- Udvardy, M. D. F. 1977. The Audubon Society field guide to North American birds: western region. Alfred A. Knopf, New York, N.Y.
- United States Department of Agriculture. 1986. Conservation Reserve Program: outline and administration handbook. Conserv. Reserve Program, Deputy Adm., State and County Operations, Washington, D.C.
- United States Department of Energy. 1992. Environmental assessment: sharp-tailed grouse and pygmy rabbit wildlife mitigation project. Bonneville Power Adm., Portland, Oregon.
-))))). 1996. Hanford site biological resources Management Plan. U.S. Dept. of Energy, Richland, WA DOE/RLL 96-32.
- Vale, T. R. 1975. Resettlement vegetation in the sagebrush grass areas of the intermountain west. *J. Range Manage.* 28:32-36.
- Wakkinen, W. L. 1990. Nest site characteristics and spring-summer movements of migratory sage grouse in southeastern Idaho. M.S. Thesis, Univ. Idaho, Moscow.

-))))), K.P. Reese, and J.W. Connelly. 1992. Sage grouse nest locations in relation to leks. *J. Wildl. Manage.* 56:381-383.
- Wallestad, R. O. 1971. Summer movements and habitat use by sage grouse broods in central Montana. *J. Wildl. Manage.* 35:129-136.
-)))) and R. C. Watts. 1972. Factors effecting annual sage grouse productivity in central Montana. Fed. Aid Wildl. Restor. Proj. W-120-R-3, Job G.B.-2.01, Plan III. Mont. Dept. Fish and Game, Helena.
-)))) and D. Pyrah. 1974. Movement and nesting of sage grouse hens in central Montana. *J. Wildl. Manage.* 38:630-633.
-)))) and P. Schladweiler. 1974. Breeding season movements and habitat selection of male sage grouse. *J. Wildl. Manage.* 38:634-637.
-)). 1975. Life history and habitat requirements of sage grouse in central Montana. Mont. Dept. Fish and Game, Game Manage. Div., Helena.
-)))) , J. G. Peterson, and R. L. Eng. 1975. Foods of adult sage grouse in central Montana. *J. Wildl. Manage.* 39:628-630.
- Washington Department of Fish and Wildlife. 1995. Washington State management plan for sage grouse. Game Div., Wash. Dept. Fish and Wildl., Olympia.
- Washington State Legislature. 1993. House Bill 1309. First Special Session, Wash. State Leg., Olympia.
- Welch, B. L., J. C. Pederson, and R. L. Rodriguez. 1988. Selection of big sagebrush by sage grouse. *Great Basin Natl.* 48:274-279.
- West, N. E. 1983. Western intermountain sagebrush steppe. Pages 351 to 397 in N. E. West, ed. *Ecosystems of the World 5: Temperate deserts and semi-deserts.* Elsevier Scientific Publ. LTD, Sussex, U.K.
- Wiley, R. H. 1973. Territoriality and non-random mating in sage grouse, *Centrocercus urophasianus*. *Anim. Behav. Monogr.* 6:85-169.
-)). 1974. Evolution of social organization and life-history patterns among grouse. *Q. Rev. Biol.* 49:201-227.
-)). 1978. The lek mating system of sage grouse. *Sci. Am.* 238:114-125.

Willis, M. J. 1991. Sage grouse winter habitat use and survival. Fed. Aid Wildl. Restor. Proj. W-87-R-4. Oreg. Dept. Fish and Wildl, Portland.

))))), G. P. Keister, Jr., D. A. Immell, D. M. Jones, R. M. Powell, and K. R. Durbin. 1993. Sage grouse in Oregon. Oreg. Dept. Fish and Wildl., Wildl. Res. Sect., Wildl. Res. Rep. 15.

Yocom, C. F. 1956. The sage hen in Washington State. *Auk* 73:540-550.

Young, J. R. 1991. Can sexual selection cause speciation among sage grouse populations? Page 22 in J. W. Connelly, ed. Seventeen western states sage and columbian sharp-tailed grouse workshop transactions. Western States Sage Grouse and Columbian Sharp-tailed Grouse Tech. Comm.

Zablan, M.A. 1990. Progress report: sage grouse survival estimation, North Park, Colorado. Colorado Div. of Wildlife, Fort Collins, 15p.

Zeigler, D. L. 1978. Distribution and status of sage grouse in eastern Washington. Fed. Aid Wildl. Restor. Proj. W-70-R-17, Job 1. Wash. Dept. Fish and Wildl., Olympia.

Zunino, G. W. 1987. Harvest effect on sage grouse densities in northwest Nevada. M.S. Thesis, Univ. Nevada, Reno.

PERSONAL COMMUNICATIONS

Ed Bracken
Habitat Biologist
Washington Dept. Fish and Wildlife
Yakima, Washington

Clait Braun
Wildlife Research Leader
Colo. Div. Wildl.
Wildlife Research Center
Fort Collins, Colorado

Larry Cadwell
Research Scientist
Battelle Pacific Northwest Laboratories
Richland, Washington

Jack Connelly
Research Biologist
Idaho Dept. Fish and Game
Pocatello, Idaho

Lisa Fitzner
Wildlife Biologist
Washington Dept. Fish and Wildlife
Yakima, Washington

Ron Friesz
Wildlife Biologist
Washington Dept. Fish and Wildlife
Ephrata, Washington

Jerry Hickman
Wildlife Biologist
Washington Dept. Fish and Wildlife
Spokane, Washington

Steve Kruger
Director
Environmental and Natural Resource
Division
U.S. Dept. Army
Yakima Training Center
Yakima, Washington

Margaret Pounds
Environmental and Natural Resource
Division
U.S. Dept. Army
Yakima Training Center
Yakima, Washington

Chuck Perry
Habitat Biologist
Washington Dept. Fish and Wildlife
Ephrata, Washington

Mike Schroeder
Upland Bird Research Specialist
Washington Dept. Fish and Wildlife
Ephrata, Washington

Lee Stream
Wildlife Biologist
Washington Dept. Fish and Wildlife
Yakima, Washington

ACRONYMS

Agricultural Conservation Program (ACP)
Bonneville Power Administration (BPA)
Conservation Reserve Program (CRP)
Priority Habitat and Species Program (PHS)
United State Bureau of Land Management (USBLM)
United States Bureau of Reclamation (USBOR)
United States Fish and Wildlife Service (USFWS)
United States Department of the Army (U.S. Army)
United States Department of Energy (USDOE)
United States Dept. of Agriculture, Agricultural Stabilization and Conservation Service (ASCS)
United States Department of Agriculture, Natural Resources Conservation Service (NRCS)
Upland Wildlife Restoration Program (UWRP)
Washington Department of Fish and Wildlife (WDFW)
Washington Department of Natural Resources (WDNR)
Washington Wildlife Recreation Coalition (WWRC)

Appendix A. Museum specimens of sage grouse in Washington, 1853-1949.

Location	County	Date ^a	Sex or set	<i>n</i>	Source ^b
Crab Creek	Lincoln	na	na		USNM # 429139
na	na	9/16/1853	na		USNM # A10019
Spokane Plain	Spokane	11/1853	na		USNM # A10021
Yakima River	Yakima	9/14/1853	na		USNM # A10022
Coulee City	Grant	6/30/1902	female		WSUCM # 682
White Bluffs	na	6/28/1903	female		WSUCM # 421
Yakima	Yakima	6/22/1910	female		MCZ # 253835
Yakima	Yakima	7/1910	male		MCZ # 253836
Yakima	Yakima	10/25/1910	male		AMNH # 751239
Kiona	Benton	2/28/1919	male		FMNH # 157308
Priest Rapids	Yakima	4/29/1919	egg set	9	PSM # 13559
Tampico	Yakima	4/11/1921	egg set	11	PSM # 13561
Kiona	Benton	11/15/1921	male		FMNH # 157307
na	Yakima	4/7/1926	egg set	11	PSM # 13560
Rattlesnake Hills	Benton	10/16/1927	male		PSM # 05890
Selah	Yakima	4/1/1928	egg set	7	WFVZ # 309a
Beverly	Kittitas	4/18/1928	egg set	1	WFVZ # 309c
Squaw Creek	Kittitas	4/16/1933	male		UWBM # 11462
Squaw Creek	Kittitas	4/28/1935	male		PSM # 07065
Rattlesnake Hills	Benton	3/30/1941	egg set	8	WFVZ # 309b
Badger Pocket	Kittitas	10/5/1941	male		PSM # 08429
10 mi W of Ephrata	Grant	7/15/1949	female		WSUCM # 49-355
Badger Pocket	Kittitas	6/24/1950	male		PSM # 07061
Badger Pocket	Kittitas	6/24/1950	female		PSM # 07064
Tieton	Yakima	9/18/1955	female		PSM # 07067
Selah	Yakima	3/19/1964	female		PSM # 07066
Bridgeport	Douglas	10/20/1973	female		UWBM # 35310
7 mi E of Winthrop	Douglas	4/23/1979	male		UWBM # 32148

^a na=information not available.

^b USNM=Smithsonian Institute, National Museum of Natural History, Washington D.C.; MCZ=Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts; WSUCM=Washington State University, Conner Museum, Pullman, Washington; AMNH=American Museum of Natural History, New York, New York; FMNH=Field Museum of Natural History, Chicago, Illinois; WFVZ=Western Foundation of Vertebrate Zoology, Camarillo, California; UWBM=University of Washington, Burke Museum, Seattle, Washington; PSM=University of Puget Sound, Slater Museum, Tacoma, Washington.

Appendix B. Sage grouse lek counts in Washington 1970-1997^a.

Lek Sites	Year																											
	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97
Douglas County																												
1	22	38	33	23	19	12		11	6	5	3	2	0	0		0	0	0	0	0	0	0	0	0	9	6	8	
2			6		2			3	10	26	16	12	14	10	5	1	1	7	0	3	1	0	1	1	0	0	0	0
3		0																			0							0
4			5	40	2			24	28	27	25	29	19	23	14	0	0	0	0	0	0	2	0	0	0	0	0	0
5	24	40	20	21	19	13	7	7	6	4	7	0			2	0	1	0	3	0		0	13	7	10	14	23	25
6		67	34	51	27			16	15	20	24	35	32	33		16	14	24	57	63	64	69	13	29	33	34	60	58
7																											9	11
8					>1											4										21	14	11
9	50	52	59	54	21		17	30	24	46	47	31	27	30	31	9	11	10	13	11	14	5	6	4	5	6	6	2
10			55																									0
11		25	13	11	19		17	19	16	13	12	21	23	13	15	5	5	5	14	14	6	4	2	0	0	0	0	0
12										>1																		
13																					6	1	0	0	0	0	0	0
14									16	20	67	71	47	39	23	12	61	71	90	91	88	71	36	11	18	25	22	22
15		26	15	20	0		0																0	0	0	0	0	0
16																							0	0	4	1	0	0
17																						2		1	0	0	0	1
18										28	33	38	30	31	17	2	3	15	45	46	55	49	26	14	8	26	22	36
19									33	45	48	35	61	57	43	22	19	37	52	63	57	53	24	26	19	20	21	26
20										4													0	0	0	0	0	0
21										>1															2	0	0	0
22								9	4	3	7	7		0		0	0		0	0	0	0	0	0	0	0	0	0
23								>1														0	0	0	0	0	0	0
24																						3	0	0	0	0	0	0
25				13	0			3	0	0	0						0				0	0	0	0	0	0	0	0
26			21	17	10			11	14	16	37	33	19	23	15	8	5	19	22	25	16	19	16	13	11	11	14	23
27	63	42	36	23	11	3	1	0	0	0			3	0	0	0	0	0	0	2	1	0	0	2	0	0	0	0
28		15							9	0		0							23	5			9	4	>1	21	20	8

Appendix B. Cont'd.

Lek Sites	Year																												
	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	
Kittitas County																													
1 *																													0
2														14		15	2	10	0	2	1	0							0
3														3															
4 *																													
5																					4								
6																					7								
7						18	20	36	29	39	83	94	96	88	80	73	42	54	54	53	50	74	67	49	43	33	24	34	
8 *																													
9														6															
Yakima County																													
1																					6	7	14	19	22	13	8	7	5
2																													25
3																												5	21
4																												17	18
5																												11	14
6											15																		0
7																	10			22	17	33	18	18	15	12	8	10	
8																				15	15	16	9	15	5	3	3	4	
9																													13
10														28	15	26	17	19	11										0
11																					12	11	31	22	25	19	8	4	4
12			20	18	12	8	5	4	9	8	11	10	19	26	30	30	25	18	15	5	2								
13																		0	2										
14																													10

^a 0 = the lek was surveyed, but no birds were present; >1= more than one bird was seen, but no counts are available.

* Historic site; lek counts prior to 1970..

Appendix C. Washington Administrative Code 232-12-297.

WAC 232-12-011 Wildlife classified as protected shall not be hunted or fished.

Protected wildlife are designated into three subcategories: Threatened, sensitive, and other.

(1) Threatened species are any wildlife species native to the state of Washington that are likely to become endangered within the foreseeable future throughout a significant portion of their range within the state without cooperative management or removal of threats. Protected wildlife designated as threatened include:

Common Name	Scientific Name
Western gray squirrel	<i>Sciurus griseus</i>
Steller (northern) sea lion	<i>Eumetopias jubatus</i>
North American lynx	<i>Lynx canadensis</i>
Aleutian Canada goose	<i>Branta canadensis leucopareia</i>
Bald eagle	<i>Haliaeetus leucocephalus</i>
Ferruginous hawk	<i>Buteo regalis</i>
Marbled murrelet	<i>Brachyramphus marmoratus</i>
Green sea turtle	<i>Chelonia mydas</i>
Loggerhead sea turtle	<i>Caretta caretta</i>

(2) Sensitive species are any wildlife species native to the state of Washington that are vulnerable or declining and are likely to become endangered or threatened in a significant portion of their range within the state without cooperative management or removal of threats. Protected wildlife designated as sensitive include:

Common Name	Scientific Name
Gray whale	<i>Eschrichtius robustus</i>
Larch Mountain salamander	<i>Plethodon larselli</i>

(3) Other protected wildlife include:

Common Name	Scientific Name
Cony or pika	<i>Ochotona princeps</i>
Least chipmunk	<i>Tamias minimus</i>
Yellow-pine chipmunk	<i>Tamias amoenus</i>
Townsend's chipmunk	<i>Tamias townsendii</i>
Red-tailed chipmunk	<i>Tamias ruficaudus</i>
Hoary marmot	<i>Marmota caligata</i>
Olympic marmot	<i>Marmota olympus</i>
Cascade golden-mantled ground squirrel	<i>Spermophilus saturatus</i>
Golden-mantled ground squirrel	<i>Spermophilus lateralis</i>
Washington ground squirrel	<i>Spermophilus washingtoni</i>
Red squirrel	<i>Tamiasciurus hudsonicus</i>
Douglas squirrel	<i>Tamiasciurus douglasii</i>
Northern flying squirrel	<i>Glaucomys sabrinus</i>
Fisher	<i>Martes pennanti</i>
Wolverine	<i>Gulo gulo</i>
Painted turtle	<i>Chrysemys picta</i>
California mountain kingsnake	<i>Lampropeltis zonata</i> ;

All birds not classified as game birds, predatory birds or endangered species, or designated as threatened species or sensitive species; all bats, except when found in or immediately adjacent to a dwelling or other occupied building; mammals of the order *Cetacea*, including whales, porpoises, and mammals of the order *Pinnipedia* not otherwise classified as endangered species, or designated as threatened species or sensitive species. This section shall not apply to hair seals and sea lions which are threatening to damage or are damaging commercial fishing gear being utilized in a lawful manner or when said mammals are damaging or threatening to damage commercial fish being lawfully taken with commercial gear.

[Statutory Authority: RCW 77.12.020. 97-18-019(Order 97-167), § 232-12-011, filed 8/25/97, effective 9/25/97. Statutory Authority: RCW 77.12.040, 77.12.020, 77.12.030 and 77.32.220. 97-12-048, § 232-12-011, filed 6/2/97, effective 7/3/97. Statutory Authority: RCW 77.12.020. 93-21-027 (Order 615), § 232-12-011, filed 10/14/93, effective 11/14/93; 90-11-065 (Order 441), § 232-12-011, filed 5/15/90, effective 6/15/90. Statutory Authority: RCW 77.12.040. 89-11-061 (Order 392), § 232-12-011, filed 5/18/89; 82-19-026 (Order 192), § 232-12-011, filed 9/9/82; 81-22-002 (Order 174), § 232-12-011, filed 10/22/81; 81-12-029 (Order 165), § 232-12-011, filed 6/1/81.]

WAC 232-12-014 Wildlife classified as endangered species.

Endangered species include:

Pygmy rabbit	<i>Brachylagus idahoensis</i>
Gray wolf	<i>Canis lupus</i>
Grizzly bear	<i>Ursus arctos horribilis</i>
Sea otter	<i>Enhydra lutris</i>
Sei whale	<i>Balaenoptera borealis</i>
Fin whale	<i>Balaenoptera physalus</i>
Blue whale	<i>Balaenoptera musculus</i>
Humpback whale	<i>Megaptera novaeangliae</i>
Black right whale	<i>Balaena glacialis</i>
Sperm whale	<i>Physeter catodon</i>
Columbian white-tailed deer	<i>Odocoileus virginianus leucurus</i>
Woodland caribou	<i>Rangifer tarandus</i>
American white pelican	<i>Pelecanus erythrorhynchos</i>
Brown pelican	<i>Pelecanus occidentalis</i>
Peregrine falcon	<i>Falco peregrinus</i>
Sandhill crane	<i>Grus canadensis</i>
Snowy plover	<i>Charadrius alexandrinus</i>
Upland sandpiper	<i>Bartramia longicauda</i>
Spotted owl	<i>Strix occidentalis</i>
Western pond turtle	<i>Clemmys marmorata</i>
Leatherback sea turtle	<i>Dermochelys coriacea</i>
Oregon silverspot butterfly	<i>Speyeria zerene hippolyta</i>
Oregon spotted frog	<i>Rana pretiosa</i>

[Statutory Authority: RCW 77.12.020. 97-18-019 (Order 97-167), § 232-12-014, filed 8/25/97, effective 9/25/97; 93-21-026 (Order 616), § 232-12-014, filed 10/14/93, effective 11/14/93. Statutory Authority: RCW 77.12.020(6). 88-05-032 (Order 305), § 232-12-014, filed 2/12/88. Statutory Authority: RCW 77.12.040. 82-19-026 (Order 192), § 232-12-014, filed 9/9/82; 81-22-002 (Order 174), § 232-12-014, filed 10/22/81; 81-12-029 (Order 165), § 232-12-014, filed 6/1/81.]

WAC 232-12-297

Endangered, threatened, and sensitive wildlife species classification.

PURPOSE

- 1.1 The purpose of this rule is to identify and classify native wildlife species that have need of protection and/or management to ensure their survival as free-ranging populations in Washington and to define the process by which listing, management, recovery, and delisting of a species can be achieved. These rules are established to ensure that consistent procedures and criteria are followed when classifying wildlife as endangered, or the protected wildlife subcategories threatened or sensitive.

DEFINITIONS

For purposes of this rule, the following definitions apply:

- 2.1 "Classify" and all derivatives means to list or delist wildlife species to or from endangered, or to or from the protected wildlife subcategories threatened or sensitive.
- 2.2 "List" and all derivatives means to change the classification status of a wildlife species to endangered, threatened, or sensitive.
- 2.3 "Delist" and its derivatives means to change the classification of endangered, threatened, or sensitive species to a classification other than endangered, threatened, or sensitive.
- 2.4 "Endangered" means any wildlife species native to the state of Washington that is seriously threatened with extinction throughout all or a significant portion of its range within the state.
- 2.5 "Threatened" means any wildlife species native to the state of Washington that is likely to become an endangered species within the foreseeable future throughout a significant portion of its range within the state without cooperative management or removal of threats.
- 2.6 "Sensitive" means any wildlife species native to the state of Washington that is vulnerable or declining and is likely to become endangered or threatened in a significant portion of its range within the state without cooperative management or removal of threats.
- 2.7 "Species" means any group of animals classified as a species or subspecies as commonly accepted by the scientific community.
- 2.8 "Native" means any wildlife species naturally occurring in Washington for purposes of breeding, resting, or foraging, excluding introduced species not found historically in this state.
- 2.9 "Significant portion of its range" means that portion of a species' range likely to be essential to the long term survival of the population in Washington.

LISTING CRITERIA

- 3.1 The commission shall list a wildlife species as endangered, threatened, or sensitive solely on the basis of the biological status of the species being considered, based on the preponderance of scientific data available, except as noted in section 3.4.
- 3.2 If a species is listed as endangered or threatened under the federal Endangered Species Act, the agency will recommend to the commission that it be listed as endangered or threatened as specified in section 9.1. If listed, the agency will proceed with development of a recovery plan pursuant to section 11.1.
- 3.3 Species may be listed as endangered, threatened, or sensitive only when populations are in danger of failing, declining, or are

vulnerable, due to factors including but not restricted to limited numbers, disease, predation, exploitation, or habitat loss or change, pursuant to section 7.1.

- 3.4 Where a species of the class Insecta, based on substantial evidence, is determined to present an unreasonable risk to public health, the commission may make the determination that the species need not be listed as endangered, threatened, or sensitive.

DELISTING CRITERIA

- 4.1 The commission shall delist a wildlife species from endangered, threatened, or sensitive solely on the basis of the biological status of the species being considered, based on the preponderance of scientific data available.
- 4.2 A species may be delisted from endangered, threatened, or sensitive only when populations are no longer in danger of failing, declining, are no longer vulnerable, pursuant to section 3.3, or meet recovery plan goals, and when it no longer meets the definitions in sections 2.4, 2.5, or 2.6.

INITIATION OF LISTING PROCESS

- 5.1 Any one of the following events may initiate the listing process.
 - 5.1.1 The agency determines that a species population may be in danger of failing, declining, or vulnerable, pursuant to section 3.3.
 - 5.1.2 A petition is received at the agency from an interested person. The petition should be addressed to the director. It should set forth specific evidence and scientific data which shows that the species may be failing, declining, or vulnerable, pursuant to section 3.3. Within 60 days, the agency shall either deny the petition, stating the reasons, or initiate the classification process.
 - 5.1.3 An emergency, as defined by the Administrative Procedure Act, chapter 34.05 RCW. The listing of any species previously classified under emergency rule shall be governed by the provisions of this section.
 - 5.1.4 The commission requests the agency review a species of concern.
- 5.2 Upon initiation of the listing process the agency shall publish a public notice in the Washington Register, and notify those parties who have expressed their interest to the department, announcing the initiation of the classification process and calling for scientific information relevant to the species status report under consideration pursuant to section 7.1.

INITIATION OF DELISTING PROCESS

- 6.1 Any one of the following events may initiate the delisting process:
- 6.1.1 The agency determines that a species population may no longer be in danger of failing, declining, or vulnerable, pursuant to section 3.3.
 - 6.1.2 The agency receives a petition from an interested person. The petition should be addressed to the director. It should set forth specific evidence and scientific data which shows that the species may no longer be failing, declining, or vulnerable, pursuant to section 3.3. Within 60 days, the agency shall either deny the petition, stating the reasons, or initiate the delisting process.
 - 6.1.3 The commission requests the agency review a species of concern.
- 6.2 Upon initiation of the delisting process the agency shall publish a public notice in the Washington Register, and notify those parties who have expressed their interest to the department, announcing the initiation of the delisting process and calling for scientific information relevant to the species status report under consideration pursuant to section 7.1.

SPECIES STATUS REVIEW AND AGENCY RECOMMENDATIONS

- 7.1 Except in an emergency under 5.1.3 above, prior to making a classification recommendation to the commission, the agency shall prepare a preliminary species status report. The report will include a review of information relevant to the species' status in Washington and address factors affecting its status, including those given under section 3.3. The status report shall be reviewed by the public and scientific community. The status report will include, but not be limited to an analysis of:
- 7.1.1 Historic, current, and future species population trends.
 - 7.1.2 Natural history, including ecological relationships (e.g., food habits, home range, habitat selection patterns).
 - 7.1.3 Historic and current habitat trends.
 - 7.1.4 Population demographics (e.g., survival and mortality rates, reproductive success) and their relationship to long term sustainability.
 - 7.1.5 Historic and current species management activities.
- 7.2 Except in an emergency under 5.1.3 above, the agency shall prepare recommendations for species classification, based upon scientific data contained in the status report. Documents shall be prepared to determine the environmental consequences of adopting the recommendations pursuant to requirements of the State Environmental Policy Act (SEPA).
- 7.3 For the purpose of delisting, the status report will include a review of recovery plan goals.

PUBLIC REVIEW

- 8.1 Except in an emergency under 5.1.3 above, prior to making a recommendation to the commission, the agency shall provide an opportunity for interested parties to submit new scientific data relevant to the status report, classification recommendation, and any SEPA findings.
- 8.1.1 The agency shall allow at least 90 days for public comment.

- 8.1.2 The agency will hold at least one public meeting in each of its administrative regions during the public review period.

FINAL RECOMMENDATIONS AND COMMISSION ACTION

- 9.1 After the close of the public comment period, the agency shall complete a final status report and classification recommendation. SEPA documents will be prepared, as necessary, for the final agency recommendation for classification. The classification recommendation will be presented to the commission for action. The final species status report, agency classification recommendation, and SEPA documents will be made available to the public at least 30 days prior to the commission meeting.
- 9.2 Notice of the proposed commission action will be published at least 30 days prior to the commission meeting.

PERIODIC SPECIES STATUS REVIEW

- 10.1 The agency shall conduct a review of each endangered, threatened, or sensitive wildlife species at least every five years after the date of its listing. This review shall include an update of the species status report to determine whether the status of the species warrants its current listing status or deserves reclassification.
- 10.1.1 The agency shall notify any parties who have expressed their interest to the department of the periodic status review. This notice shall occur at least one year prior to end of the five year period required by section 10.1.
- 10.2 The status of all delisted species shall be reviewed at least once, five years following the date of delisting.
- 10.3 The department shall evaluate the necessity of changing the classification of the species being reviewed. The agency shall report its findings to the commission at a commission meeting. The agency shall notify the public of its findings at least 30 days prior to presenting the findings to the commission.
- 10.3.1 If the agency determines that new information suggests that classification of a species should be changed from its present state, the agency shall initiate classification procedures provided for in these rules starting with section 5.1.
 - 10.3.2 If the agency determines that conditions have not changed significantly and that the classification of the species should remain unchanged, the agency shall recommend to the commission that the species being reviewed shall retain its present classification status.
- 10.4 Nothing in these rules shall be construed to automatically delist a species without formal commission action.

RECOVERY AND MANAGEMENT OF LISTED SPECIES

- 11.1 The agency shall write a recovery plan for species listed as endangered or threatened. The agency will write a management plan for species listed as sensitive. Recovery and management plans shall address the listing criteria described in sections 3.1 and 3.3, and shall include, but are not limited to:
- 11.1.1 Target population objectives.
 - 11.1.2 Criteria for reclassification.

- 11.1.3 An implementation plan for reaching population objectives which will promote cooperative management and be sensitive to landowner needs and property rights. The plan will specify resources needed from and impacts to the department, other agencies (including federal, state, and local), tribes, landowners, and other interest groups. The plan shall consider various approaches to meeting recovery objectives including, but not limited to regulation, mitigation, acquisition, incentive, and compensation mechanisms.
 - 11.1.4 Public education needs.
 - 11.1.5 A species monitoring plan, which requires periodic review to allow the incorporation of new information into the status report.
- 11.2 Preparation of recovery and management plans will be initiated by the agency within one year after the date of listing.
- 11.2.1 Recovery and management plans for species listed prior to 1990 or during the five years following the adoption of these rules shall be completed within five years after the date of listing or adoption of these rules, whichever comes later. Development of recovery plans for endangered species will receive higher priority than threatened or sensitive species.
 - 11.2.2 Recovery and management plans for species listed after five years following the adoption of these rules shall be completed within three years after the date of listing.
 - 11.2.3 The agency will publish a notice in the Washington Register and notify any parties who have expressed interest to the department interested parties of the initiation of recovery plan development.
 - 11.2.4 If the deadlines defined in sections 11.2.1 and 11.2.2 are not met the department shall notify the public and report the reasons for missing the deadline and the strategy for completing the plan at a commission meeting. The intent of this section is to recognize current department personnel resources are limiting and that development of recovery plans for some of the species may require significant involvement by interests outside of the department, and therefore take longer to complete.
- 11.3 The agency shall provide an opportunity for interested public to comment on the recovery plan and any SEPA documents.

- 13.1 The commission has the authority to classify wildlife as endangered under RCW 77.12.020. Species classified as endangered are listed under WAC 232-12-014, as amended.
- 13.2 Threatened and sensitive species shall be classified as subcategories of protected wildlife. The commission has the authority to classify wildlife as protected under RCW 77.12.020. Species classified as protected are listed under WAC 232-12-011, as amended. [Statutory Authority: RCW 77.12.020. 90-11-066 (Order 442), § 232-12-297, filed 5/15/90, effective 6/15/90.]

CLASSIFICATION PROCEDURES REVIEW

- 12.1 The agency and an ad hoc public group with members representing a broad spectrum of interests, shall meet as needed to accomplish the following:
 - 12.1.1 Monitor the progress of the development of recovery and management plans and status reviews, highlight problems, and make recommendations to the department and other interested parties to improve the effectiveness of these processes.
 - 12.1.2 Review these classification procedures six years after the adoption of these rules and report its findings to the commission.

AUTHORITY