


# Periodic Status Review for the Greater Sage-grouse



**Derek W. Stinson**  
*Washington Department of  
FISH AND WILDLIFE  
Wildlife Program*

The Washington Department of Fish and Wildlife maintains a list of endangered, threatened, and sensitive species (Washington Administrative Codes 220-200-100 and 220-610-010). In 1990, the Washington Wildlife Commission adopted listing procedures developed by a group of citizens, interest groups, and state and federal agencies (Washington Administrative Code 220-610-110). The procedures include how species listings will be initiated, criteria for listing and delisting, a requirement for public review, the development of recovery or management plans, and the periodic review of listed species.

The Washington Department of Fish and Wildlife is directed to conduct reviews of each endangered, threatened, or sensitive wildlife species at least every five years after the date of its listing by the Washington Fish and Wildlife Commission. The periodic status reviews are designed to include an update of the species status report to determine whether the status of the species warrants its current listing status or deserves reclassification. The agency notifies the general public and specific parties who have expressed their interest to the Department of the periodic status review so that they may submit new scientific data to be included in the review. The agency notifies the public of its recommendation at least 30 days prior to presenting the findings to the Fish and Wildlife Commission. In addition, if the agency determines that new information suggests that the classification of a species should be changed from its present state, the agency prepares documents to determine the environmental consequences of adopting the recommendations pursuant to requirements of the State Environmental Policy Act.

This is the final Periodic Status Review for the Greater Sage-grouse. It contains an update of information pertaining to the status of the Greater Sage-grouse in Washington since the publication of the last periodic status review (Stinson et al. 2016). The draft was reviewed by species experts and was available for a 90-day public comment period from 28 September to 30 December 2020; comments received were considered during preparation of this final document. The Department presented the results of this periodic status review to the Fish and Wildlife Commission at a meeting on 26 March 2021. The Commission voted on 23 April 2021 to up-list the Greater Sage-grouse to endangered in Washington.

**This report should be cited as:**

Stinson, D. W. 2020. Periodic status review for the Greater Sage-grouse in Washington. Washington Department of Fish and Wildlife, Olympia, Washington. 20+ iv pp.


*On the cover: photos of male sage-grouse by Mike Schroeder; background by Mark Teske; black and white illustrations by Darrell Pruett*



*This work was supported in part by personalized and endangered species license plates*



# Periodic Status Review for the Greater Sage-grouse in Washington



Prepared by  
Derek W. Stinson

Washington Department of Fish and Wildlife  
Wildlife Management Program  
600 Capitol Way North  
Olympia, WA 98501-1091

**April 2021**

## TABLE OF CONTENTS

EXECUTIVE SUMMARY .....	iii
INTRODUCTION .....	1
<i>Distribution</i> .....	1
LIFE HISTORY .....	1
POPULATION STATUS .....	3
HABITAT STATUS .....	5
FACTORS AFFECTING GREATER SAGE-GROUSE IN WASHINGTON .....	8
Adequacy of Regulatory Mechanisms .....	8
Continued Habitat Loss, Fragmentation or Degradation .....	9
Other Factors Affecting Sage-grouse .....	10
MANAGEMENT ACTIVITIES .....	11
CONCLUSION AND RECOMMENDATION .....	13
REFERENCES CITED .....	15
PERSONAL COMMUNICATIONS .....	20
APPENDIX A. PUBLIC COMMENTS .....	20
Washington State Status Reports, Periodic Status Reviews, Recovery Plans, and Conservation Plans .....	21

## LIST OF FIGURES AND TABLES

Figure 1. Historical range and breeding range (Habitat Concentration Areas, circa 2012) of sage-grouse in Washington. ....	1
Figure 2. State-wide population estimate and significant events for sage-grouse in Washington, 1969-2020 (spring, before the fires). ....	3
Figure 3. Estimates for the Douglas County and JBLM-YTC populations of sage-grouse in Washington, 1980-2020. ....	4
Figure 4. Land cover and Sage-grouse Management Units, Habitat Concentration Areas (Robb and Schroeder 2012), and the 2020 fires. ....	6

## ACKNOWLEDGMENTS

Copies of reports or survey data were provided by Jason Lowe and Colin Leingang. Preparation was made easier by recent reports by Colleen Stinson, Jason Lowe, and Kourtney Stonehouse, Kyle Ebenhoch, and Kevin White. The draft was improved by reviews by Kevin White, Ellie Mangelinckx, Lisa Shipley, Grant Casady, Jason Lowe, Mike Atamian, and Mike Schroeder. Many thanks to our partners and cooperators in Sage-grouse conservation, including Bureau of Land Management, Oregon Department of Fish and Wildlife, Washington State University, U.S. Fish and Wildlife Service, Joint Base Lewis-McChord-Yakima Training Center, Spokane Audubon, Yakima Audubon, Yakama Nation, USDA Farm Service Agency, Natural Resource Conservation Service, The Sage-grouse Initiative, Wenatchee Sportsmen, Inland Northwest Wildlife Council, Foster Creek Conservation District, and Lincoln County Conservation District.

## EXECUTIVE SUMMARY

The Greater Sage-grouse (*Centrocercus urophasianus*) historically was found throughout the shrub-steppe areas of eastern Washington. The species is now limited in distribution in the state to Douglas County, the Joint Base Lewis-McChord Yakima Training Center (JBLM-YTC), and Lincoln County where a reintroduction project re-established a small population. The state-wide population estimate, based on lek counts, was 775 in spring 2020, an increase from 688 birds in 2019, as a result of an increase in Douglas County from 597 to 697, while Lincoln County decline from 13 to 10, and the JBLM-YTC declined from 78 to 68. Subsequent to those counts, the habitat of all three populations were affected by wildfires. Preliminary assessments suggest that the Douglas County population will be reduced by ~50% due to loss of sagebrush on half the occupied habitat, and mortalities primarily from high predation due to lack of cover. The struggling Lincoln County population will probably be extirpated.

The sage-grouse was state-listed as threatened in 1998, and a state recovery plan was completed in 2004. From 2001–2015, the Columbia Basin sage-grouse population was a candidate for listing as a threatened Distinct Population Segment (DPS) under the U.S. Endangered Species Act. In September 2015, the U.S. Fish and Wildlife Service issued a decision that the population in Washington did not meet the criteria for a DPS and listing of the Greater Sage-grouse across its entire range was not warranted. However, since that decision, genomic analyses highlighted the unique nature of the Washington population.

The potential for wildfires to eliminate sagebrush (*Artemisia* spp.) on extensive areas has been the greatest ongoing threat to sage-grouse in Washington, as we have seen in 2020. However, with the continued decline, all of Washington's populations are now likely suffering from problems with genetic health and fitness related to small population size. Uncertainty about the long-term maintenance of habitat that depends on Farm Bill programs (CRP/SAFE) is also a concern. Other major management issues include habitat that is fragmented by roads, agriculture, and development and degraded by past wildfires, historical excessive livestock grazing, fencing, electrical transmission lines, and exotic vegetation. Sage-grouse may suffer mortality rates above historical levels as a result of collisions with fences, powerlines, and vehicles, and higher populations of some generalist predators, especially ravens and coyotes.

The Washington Department of Fish and Wildlife (WDFW) and several partner organizations are working on habitat and other aspects of sage-grouse recovery. Without these efforts, the sage-grouse would likely decline to extinction in Washington. In Spring 2020, sage-grouse had not yet declined to populations levels indicated in the 2004 state recovery plan for up-listing (<650 birds); however, that was before the devastating fires of September, and the threshold assumed that the Douglas County and JBLM-YTC populations were connected, which now appears unjustified. Due in part to their polygynous mating system, the effective size of the three populations are ~108 birds for Douglas County and 10 birds for JBLM-YTC. Extinction of the Lincoln County population is all but certain, and of the JBLM-YTC within a decade or so is likely unless they can be increased substantially. The hope of any reintroductions in the future is tempered by the recent failure of the reintroduction project by the Yakama Nation, the probable failure of the Lincoln County population, and the continued loss of habitat in suitable condition by wildland fire.

Concurrent with this troubling decline, genomic analysis has indicated that Washington's population is more distinct than the Bi-state population that was proposed for listing as a threatened 'Distinct Population Segment' under the Endangered Species Act (USFWS 2019). For these reasons, it is recommended the sage-grouse be up-listed to endangered in Washington.

## INTRODUCTION

The Greater Sage-grouse (*Centrocercus urophasianus*; hereafter sage-grouse), the largest grouse species in North America, was once abundant in the shrub-steppe of eastern Washington. Sage-grouse are closely associated with sagebrush (*Artemisia* spp.) and populations require extensive areas of sagebrush habitat to persist. Sage-grouse hunting seasons were closed in Washington in 1987 due to population concerns. The species was state-listed as threatened in 1998, and a recovery plan was completed in 2004 (Stinson et al. 2004). Sage-grouse in Washington are the most genetically unique of any subpopulation in North America and might serve as important genetic reservoirs of adaptive diversity (Oh et al. 2019), if they can be recovered. The Washington birds are more likely to nest, more likely to re-nest, and they lay more eggs than is typical elsewhere (Schroeder 1997), and they are also 15% larger than sage-grouse to the south (M. Schroeder, pers. comm.). Based on their genomic analyses, Oh et al. (2019) stated that, “highly differentiated populations like the Washington greater sage-grouse may warrant recognition and protection as a genetically distinct conservation unit.”

This document is an update of the 2016 periodic status review. It includes the most recent information and estimates for the populations in Washington, as well as a brief synopsis of recent research and management activity. The Population and Habitat Status section have been updated substantially and there is a recommendation for a change in classification, particularly in light of the September 2020 fires.

**Distribution.** Sage-grouse persist in two main areas in Washington: one primarily on the U.S. Army’s Joint Base Lewis McChord Yakima Training Center (JBLM-YTC) in Kittitas and Yakima counties and the other, often referred to as the Moses Coulee population, in Douglas County and potentially, adjacent parts of Grant County (Fig. 1). Populations were being re-established in Lincoln County (i.e. Crab Creek), and on the Yakama Indian Reservation, but with the 2020 fire, both have now probably failed.




Figure 1. Historical range and breeding range (Habitat Concentration Areas, circa 2012) of sage-grouse in Washington.

## LIFE HISTORY

The spring courtship display of males at specific locations, called ‘leks’, is the most conspicuous behavior of sage-grouse. Sage-grouse are polygynous, with the dominant males mating with multiple females. In Douglas County, most birds return to breeding areas in late February or March. Females generally return to the same nesting area (Schroeder et al. 1999), and probably visit the same lek or leks each year. Males begin to leave leks in late April and early May and move to summer habitat (Stinson et al. 2004). After mating, females devote most of their time to nesting and brood-rearing; males do not assist in these activities. Sage-grouse females attempt to raise one brood per year (Schroeder et al. 1999). The precocial chicks feed themselves, but females spend considerable time keeping chicks warm and guarding them for the first four to five weeks.

**Habitat requirements.** Sage-grouse rely upon shrub-steppe habitat, with sagebrush comprising 60–80% of the yearly diet of adults and up to 95–100% of the winter diet (Schroeder et al. 1999). That is despite sagebrush’s high concentrations of monoterpenes, sesquiterpene lactones, and phenolics, substances thought to have evolved as chemical defenses to herbivory (Kelsey et al. 1982). Wyoming Big Sagebrush (*Artemisia tridentata wyomingensis*) and Three-tip Sagebrush (*Artemisia tripartita*) are the most important sagebrush species to sage-grouse in Washington. There is considerable geographic variation in composition and concentration of secondary compounds among and within sagebrush varieties (Welch 2005), suggesting that sage-grouse populations across the species’ ranges may be adapted for selecting and digesting the distinct chemistry of local sagebrush varieties (Oh et al. 2019).

Sage-grouse need large areas of shrub-steppe with sagebrush canopy (~15–35%), and a healthy herbaceous understory for nest concealment and brood-rearing cover and food (>10% forb cover, >15% grass cover, >18 cm grass height; Connelly et al. 2000a, Stinson et al. 2004, Hagen et al. 2007). A diverse herbaceous layer supplies food to females during the pre-laying period, and forbs and insects, particularly grasshoppers, beetles, and ants, are important during early chick development and thus are necessary for recruitment (Gregg and Crawford 2009, Connelly et al. 2011b). Later in summer, the diet of juveniles shifts from insects to more forbs, and broods often move to higher elevations or more mesic sites, such as seeps, riparian areas, and alfalfa fields that stay green when the vegetation of surrounding areas has dried (Connelly et al. 2011b). Sage-grouse broods in Washington do not seem to move to higher elevations and riparian habitats in late summer (Sveum et al. 1998, Stonehouse et al. 2015, G. Casady, pers. comm.). In winter, sage-grouse rely on the sagebrush that remains accessible above the snow for food and shelter (Connelly et al. 2011b). Sage-grouse have large home ranges; single ‘season’ home ranges (i.e. spring-summer) in Douglas County averaged 2–44 km<sup>2</sup> (0.8–12 mi<sup>2</sup>; Stinson et al. 2004:14), and in Lincoln County, spring-summer home ranges averaged 33 km<sup>2</sup> (range 4–94 km<sup>2</sup>; n=22) for males and 28 km<sup>2</sup> (range 2–174 km<sup>2</sup>; n=28) for females (Stonehouse et al. 2015). Annual home ranges averaged 94 km<sup>2</sup> (range 11–292; n=18 m, 48 f) on the JBLM-YTC (K. White, pers. comm.). Sage-grouse also tend to avoid roads, electrical distribution lines, and vertical structures (distribution poles and trees) in their spring-summer home ranges and when selecting nest sites (Schroeder and Vander Haegen 2014, Shirk et al. 2015, Stonehouse et al. 2015).

**Predation and population dynamics.** In north-central Washington, the survival rate for chicks to 50 days old was 33.4% (n=515; Stinson et al. 2004). In grouse species (subfamily Tetraoninae), predation typically accounts for about 85% of reported non-hunting mortalities and 79–94% of nest failures (Bergerud 1988: p. 615, 684; Moynahan et al. 2007). Nest success for 2012–2019 on JBLM-YTC was 24.9% calculated using nest survival models in Rmark (E. Mangelinckx, pers. comm.). Moynahan et al. (2007) reported that average seasonal nest success rate was 0.24–0.32 for early nests and 0.32–0.42 for late nests in Montana. Habitat quality, specifically the amount and type of vegetation available to conceal nests from visually hunting predators, like Common Ravens (*Corvus corax*), ultimately affects the number of nests destroyed by predators (Gregg et al. 1994, Ritchie et al. 1994, Rebholz 2007). Recent studies suggest that predation on young sage-grouse chicks can be high, and in fragmented landscapes or in areas with subsidized predators, predation can limit population growth (Hagen 2011). In Washington, ravens, Coyotes (*Canis latrans*), and American Badgers (*Taxidea taxus*), preyed on sage-grouse eggs and were responsible for many nest failures (Stinson et al. 2004, Lannoye and White 2014a, Harris Environmental Group 2015).

The annual survival rate for adult males in Douglas County was 56.9% (n=29) and 72.5% for adult females (n=88; Schroeder 2000). On JBLM-YTC, annual survival for males was 31.7% (n=24) and for females was 75.2% (n=71) from 2012–2018 (all radio-marked individuals; E. Mangelinckx, pers. comm.). Overwinter survival is generally high, and most mortalities occur in spring, summer and early fall

(Connelly et al. 2011a). Sage-grouse are relatively long-lived for gamebirds with individuals up to 9 years old recorded in the wild, and they may be able to live 14 or 15 years.

## POPULATION STATUS

Based on changes in number of males counted on lek complexes, the state-wide sage-grouse population in Washington has declined ~80% since 1970 (Fig. 2). The population estimate for spring 2020 was 775 birds associated with 21 known active leks before the late summer fires. The overall population increased 32% (510 to 710) between 2017 and 2018 but decreased 5% to 688 between 2018 and 2019. The birds were distributed between 3 populations including 697 birds with 17 leks in Douglas County, 68 birds with 3 leks on JBLM-YTC, and 10 birds with 1 lek in Crab Creek (Fig. 3). The population in Douglas County increased 17% between 2019 and 2020. The increase is a notable reversal following consistent declines between 2010 and 2017. The 2020 population is still substantially lower than the population observed in 2010. A fourth population resulting from a reintroduction project of the Yakama Nation, was lost between 2018 and 2019.




Figure 2. State-wide population estimate and significant events for sage-grouse in Washington, 1969-2020 (spring, before the fires).

The relative stability of the population from ~1997 to 2011 could probably be attributed to the maturation of Conservation Reserve Program (CRP) fields in Douglas County (Schroeder and Vander Haegen 2011). CRP enrolled lands allowed the Douglas County population to remain relatively stable, whereas the JBLM-YTC population has continued a long decline since 1983, despite inhabiting one of the largest areas (1,300 km<sup>2</sup>, 502 mi<sup>2</sup>) of shrub-steppe remaining in the state (Fig. 3). The number of active leks declined to 3 in 2019 and 2020 on the JBLM-YTC, down from 6 in most recent years. Also, sage-grouse numbers in Douglas County were affected by the wildfires of 2012, and cropland coming out of CRP, or tilling of older CRP that did not comply with stricter planting requirements of the Sage-grouse and Sharp-tailed Grouse State Acres for Wildlife (SAFE) program. An anticipated increase with habitat improvements may have been just starting to be evident in 2018 but SAFE and CRP acreage is expected to decline in the next few years.






Figure 3. Estimates for the Douglas County and JBLM-YTC populations of sage-grouse in Washington, 1980-2020.

With the 2020 fires, sage-grouse will almost certainly have declined to populations levels below that indicated in the 2004 state recovery plan for up-listing by 2021 (<650 birds); that threshold also assumed that the Douglas County and JBLM-YTC populations were connected. In the ~17 years since that uplisting objective was developed (Stinson et al. 2004), we have seen very little movement between populations, and the populations appear to be functionally isolated. Over 50% of the birds were impacted by wildfire and the fires likely reduced the carrying capacity of their habitat for some time. Lack of sagebrush food and cover may result in additional mortalities if severe winter conditions, such as crusted snow and extreme cold, occurs. The JBLM-YTC population will also likely be down further as a result of the Taylor Ponds Fire, and the struggling Lincoln County population will likely be extirpated by the Whitney Fire. The full impact will not be known until surveys are done in 2021 and 2022, but Foster et al. (2019) reported reduced adult female survival and low nest survival after a large fire in Oregon in 2012. Connelly et al. (2000b) also reported an accelerated population decline and loss of active leks after a prescribed burn in southeastern Idaho.

Perhaps a more important metric than census population estimates for predicting population persistence is the ‘effective population’ size. The effective population [ $N_e$ ] is the proportion of a population that can be expected to pass on their genetic information from one generation to the next (Frankham 1995). Schroeder (2000) estimated the ratio of census and effective populations sizes ( $N_c/N_e$ ) for Washington’s sage-grouse was 0.156. Therefore, the Douglas County population (~697 individuals in spring 2020), which was just above the up-list threshold of 650 birds, had an estimated effective population size of 108 birds. The JBLM-YTC (68 birds) had an estimated  $N_e$  of 10 birds. The ‘extinction vortex’ scenario described in the literature that results from inbreeding and reduced fitness in small populations (Gilpin and Soule 1986), and is aggravated by habitat and demographic stressors, may describe these populations. Extinction of the JBLM-YTC population within a decade or so is very likely unless they can be increased very soon, but habitat condition may no longer be adequate to long sustain a population. Population modeling based on trends of lek counts indicated that “extinction is probable for both the Moses Coulee [Douglas County] and Yakima [JBLM-YTC] population” (Garton et al. 2015). The hope of reintroductions in the future is tempered by the recent failure of the sage-grouse reintroduction project by the Yakama Nation, and the probable extirpation of the Lincoln County population.

## HABITAT STATUS

Sage-grouse in Washington inhabit large remnants of shrub-steppe on public land, areas where a matrix of private land contains a high percentage of shrub-steppe fragments, and lands enrolled in Farm Bill conservation programs (Conservation Reserve Program [CRP] and State Acres For Wildlife [SAFE]). Within this matrix, sage-grouse make use of some cropland for leks and foraging. Larger areas not converted to cropland were typically grazed by livestock, and some of these remaining shrub-steppe areas provide winter habitat, but little perennial grass or forb cover needed for nesting and brood-rearing, a legacy of historical heavy grazing. The largest areas of shrub-steppe vegetation on public lands are affected by multiple factors that have degraded their habitat value for sage-grouse as well. The current condition and situation of the Sage-grouse Management Units (SMU) with, and adjacent to extant populations (Fig. 4), are briefly described below.

***Moses Coulee & Mansfield Plateau SMUs/Moses Coulee PAC.*** The Moses Coulee population centered in Douglas County occupies a 461,583 ac Habitat Concentration Area (HCA) that is a mosaic of cropland, CRP, and patches of high-quality shrub-steppe. Washington Department of Natural Resources (DNR) owns significant portions of the Mansfield Plateau (12%) and Moses Coulee (8%) sage-grouse units, with much of that leased for grazing, crops, or in CRP; Bureau of Land Management (BLM) and WDFW also own small portions. The CRP program has been essential for providing habitat for sage-grouse (Schroeder and Vander Haegen 2006, 2011). The Sage-grouse and Sharp-tailed Grouse SAFE program (now 'Shrub-steppe' SAFE) has specific planting requirements and was expected to boost grouse populations, but the tilling of older CRP starting in 2010 precipitated a decline in grouse numbers from which the populations had not yet recovered. In 2020, the Pearl Hill Fire burned ~223,000 ac including much of the Mansfield Plateau unit indicating a loss of nesting, brood-rearing, and wintering habitat in the areas surrounding ~50% of the active leks in Douglas County. Other recent fires include the 2012 Barker Canyon (>17,000 ac) and Leahy fires (73,000 ac) that also impacted portions of the Mansfield Plateau. The Barker Canyon Fire affected sagebrush cover, but the native grasses and forbs have recovered well. The 1,350 ac Foster Creek Fire in 2012 included private lands, BLM and WDFW; most of the ~725 ac of WDFW land had suitable sage-grouse habitat.

***Joint Base Lewis-McChord Yakima Training Center SMU/ PAC.*** JBLM-YTC is an active U.S. Army training facility, and ongoing programs are required to maintain vegetation and restore impacts from training impacts and fires. Habitat modeling in 2014 identified ~92,117 ac (~30% of the installation) that likely had 15–40% sagebrush canopy cover (Lannoye and White 2014b). However, much of that habitat is located in the northern portion of the installation that receives little to no sage-grouse use, likely because this area is naturally separated from the southern portion of JBLM-YTC by several tall east-west ridges. Additionally, noise from Interstate 90 (I-90) and large power transmission lines, including the 230 kilovolt Vantage-Pomona Heights transmission line that is being constructed (BLM 2017, PacifiCorp n.d.), reduce the quality of JBLM-YTC's northern areas for sage-grouse.

Several large fires have occurred since the 2014 analysis, that contributed to further habitat loss and degradation. Fire is a constant threat on JBLM-YTC, particularly when live-fire training occurs during the driest months (June–October). In 2016, the Range 12 fire burned the southern part of JBLM-YTC and much of the shrub-steppe between the installation and the Hanford unit, totaling ~173,000 ac. Additional fires have originated from adjacent interstates (I-90, I-82), including the Boylston Fire that originated from I-90 and burned >44,000 ac in the northern portion of JBLM-YTC in 2018. As of 2018, 55% of YTC had burned at least once in the past 30 years. As of February 2020, JBLM-YTC had an estimated




Figure 4. Land cover and Sage-grouse Management Units, Habitat Concentration Areas (Robb and Schroeder 2012), and the 2020 fires.

58,090 ac (18% of the installation) of high-quality sage-grouse habitat with 15–40% sagebrush canopy cover and an additional 158,841 ac of habitat that may achieve sagebrush canopy cover of 15–40% through restoration and/or natural succession. In 2020, the lightning-caused Taylor Pond Fire burned ~17,500 ac on the YTC; ~2600 ac were areas that had not previously burned in the last 30 years. Sage-grouse on JBLM-YTC often use areas with <15% sagebrush cover because these areas often have healthy bunchgrass communities and other non-sagebrush shrubs (e.g., yellow rabbitbrush [*Chrysothamnus viscidiflorus*]) that provide cover. These areas provide seasonal habitat for the species, but lack sagebrush for winter forage. A telemetry study during 2012-2018 identified ~117,128 ac of occupied area on YTC (34% of total area) and 4,400 ac on adjacent private lands; the population core area was 23,722 ac (96 km<sup>2</sup>). The Sage-grouse Protection Area on YTC, which encompasses 78,600 ac, has temporal and spatial protections, but 53% of the total population-level home range and 34% of the population core area were outside the Sage-grouse Protection Area.

**Crab Creek SMU/PAC.** Substantial shrub-steppe habitat has remained in the Lincoln County portion of the Crab Creek SMU where ‘channeled scablands’ formed by the ice age floods contain thin or rocky soil that is poorly suited to cropland. Many of the areas with deeper soils were converted to wheat, and many were later enrolled in CRP contracts. The combined WDFW (Swanson Lakes Wildlife Area) and BLM (Twin Lakes and Telford management areas) ownership totals 53,000 ac. The relatively large blocks of suitable shrub-steppe vegetation and >3,000 ac of restored habitat, along with management changes, had improved the potential for sage-grouse since the birds were extirpated in the 1980s. Many miles of fences had been removed or marked to reduce collision mortalities of birds, and 5.7 miles of power distribution lines were removed or buried.

In 2020, the Whitney Fire burned >130,000 ac, including almost the entire Swanson Lake Wildlife Area and almost all the sage-grouse habitat that had been occupied since the reintroduction (Fig. 4). During the 10-year period from 2003–2012, approximately 29,000 ac (28%) of the Crab Creek Habitat Concentration Area in Lincoln County had burned in 6 major wildfires, including the 2012 Apache Pass Fire which affected occupied sage-grouse habitat.

**Dry Falls SMU.** This unit is very important for any future connectivity between Crab Creek and Moses Coulee units, particularly the area from Banks Lake, south to Ephrata (Robb and Schroeder 2012); a few sage-grouse are known to have moved through this area. Connectivity value is compromised by Banks Lake and two 500 kilovolt transmission lines radiating from Grand Coulee Dam, and the impact of the 2017 Spartan Fire (8,700 ac).

**Colockum SMU.** This unit is very important for potential connectivity between Moses Coulee and the JBLM-YTC populations (Robb and Schroeder 2012). The Colockum, Quilomene, and Whiskey Dick Wildlife Areas comprise 2/3 of this unit, but topography is rather rugged and connectivity value is compromised by I-90, multiple power transmission lines south of the unit on JBLM-YTC, the Wild Horse wind turbines, and the effects of the 2018 Milepost 22 Fire (7,600 ac).

**Toppenish Ridge SMU/Yakama PAC.** The Toppenish Ridge SMU, on the Yakama Indian Reservation, contains substantial areas of shrub-steppe (Jamison and Livingston 2004). The major management issues are feral horses and wildfires. The Yakama Nation has been engaged in efforts to reduce the potential for large wildfires, and to reduce feral horse numbers. Fences exclude horses from 19,500 ac and an additional 18,000 ac are being fenced. Additional funding was being pursued to enlarge the original enclosure to protect an additional 30,000 ac (D. Blodgett III, pers. comm.).

**Umtanum Ridge, Ahtanum Ridge, Hanford and Rattlesnake Hills SMUs.** These units are important for potential population expansion and connectivity between the JBLM-YTC unit and the Toppenish Ridge unit, if the decline in the JBLM-YTC population can be reversed. The Ahtanum Ridge unit was impacted

by the Ahtanum Ridge Fire (~6,000 ac) in 2020. The Umtanum Ridge was affected by the Pipeline Fire (6,500 ac) in 2019, and the Evans Canyon Fire in 2020 (76,000 ac). The Rattlesnake Hills would connect the JBLM-YTC with the Hanford SMU, but it was impacted by the Range 12 fire (176,000 ac), and the Hanford SMU has had fires repeatedly dating back to the 1980s; the low precipitation in the lower Columbia Basin makes vegetation recovery slow and restoration on the Hanford difficult.

**Bridgeport Point SMU.** The Bridgeport Point unit was affected by the Cold Springs Fire (190,000 ac) in 2020.

## FACTORS AFFECTING GREATER SAGE-GROUSE IN WASHINGTON

### Adequacy of Regulatory Mechanisms

**Federal regulation.** Sage-grouse are not directly protected by federal regulations at this time, but as a state-listed species, it is considered when federal actions would negatively affect them. From 2001-2015, the sage-grouse population in Washington was considered the Columbia Basin Distinct Population Segment (DPS) and was a candidate for listing under the Endangered Species Act. This raised the priority of sage-grouse conservation in the state and there was a consistent focus on habitat protection on federal lands, including JBLM-YTC and BLM, and for funding of conservation actions by USFWS. BLM has been, and continues to be, an important partner in sage-grouse recovery, by funding research, restoring habitat, and other conservation work. JBLM-YTC has been proactive in efforts to accommodate the needs of sage-grouse with their Army training mission. The recent decision that the population did not qualify as a DPS (USFWS 2015), however, has affected funding priority for sage-grouse work, and a first draft of the update to their Integrated Natural Resource Plan (INRMP) included reduced sage-grouse protections. Sage-grouse in Washington are the most genetically unique of any subpopulation in North America (Oh et al. 2019). The next most unique subpopulation is the Bi-state population on the border of California and Nevada, which is considered a DPS by the U.S. Fish and Wildlife Service (USFWS 2010, 2019).

**State and local regulations.** Loss of sage-grouse habitat is often caused by conversion to cropland or development. On non-federal lands, the Growth Management Act (GMA) is Washington's primary regulatory tool to protect rare and threatened species from development impacts. Local governments are required to: 1) create and implement development regulations that protect state-listed species and their habitat; 2) adopt zoning ordinances that ensure areas outside of urban growth areas remain rural in character, and; 3) ensure that development does not occur on natural resource lands designated for long-term agricultural use. However, rural densities allowed (e.g. ~1 dwelling/20 ac) by zoning meet the needs of most species, but likely exceed the tolerance of sage-grouse. The state rule implementing GMA (WAC 365-190-130) requires that wildlife habitat conservation areas (FWHCA - a type of critical area) must be designated and counties and cities should consult current information on priority habitats and species (PHS) identified by WDFW.

PHS management recommendations (Schroeder et al. 2003) are not regulatory, but they are often adopted through county regulations. Known or discovered locations of sage-grouse and habitat trigger the process of avoiding, minimizing, and mitigating impacts. Though the specific nature of these protections varies across the counties, the inclusion of sage-grouse and shrub-steppe habitat provides a mechanism for minimizing disturbance from construction and development activities. Although land use regulations generally provide some protection for wildlife and occupied habitat, they typically do not adequately protect habitat that is not occupied, thus they are poorly suited for the recovery of species that require large landscapes.

## Continued Habitat Loss, Fragmentation or Degradation

Sage-grouse are generally a species of undeveloped shrub-steppe landscapes, therefore conserving large contiguous patches of intact habitat is important (Connelly et al. 2011b). Moreover, Connelly et al. (2011b:83) cautioned that “*failure to protect what is left and fix what is broken will likely result in extirpation of many, if not most, populations of Greater Sage-Grouse.*”

A range-wide analysis of sage-grouse data found a strong negative effect of development (urban, suburban areas, and interstate and state highways) within 18 km (11.2 mi) of leks; most active leks had no developed lands within 5 km (3.1 mi; Johnson et al. 2011). Compared to where they remain, the portion of sage-grouse range where they were extirpated contained almost 27 times the human density, almost 3 times more area in agriculture, was 60% closer to highways, and had 25% higher density of roads than occupied range (Wisdom et al. 2011). The Moses Coulee and Mansfield Plateau SMUs have potential to be influenced by development because of the amount of private land and its location near significant population centers and Banks Lake, which is attractive for recreation. Development on the JBLM -YTC, including development of training ranges and other facilities, has undoubtedly had some cumulative effects, and Army training affects habitat quality through sagebrush mortality and disturbance to understory vegetation, which requires ongoing rehabilitation and restoration.

Another long-term concern with the potential for major impact on sage-grouse in Washington is that Federal Farm Bill programs (e.g. CRP/SAFE) are vulnerable to changes by Congress. Farm Bill Programs have been essential in providing additional habitat and in buffering patches of remnant natural habitat. The Sage & Sharp-tailed grouse SAFE program has 72,941 ac enrolled, and Shrub-steppe SAFE has 19,530 ac enrolled in Washington (these were combined in the 2018 Farm Bill as ‘Shrub-steppe SAFE’). The Douglas County sage-grouse population largely depends on these programs and has benefitted from a waiver of the 25% cap of cropland acres in a county; but the waiver was left out of the 2018 Farm Bill. It is also uncertain if the SAFE program will continue after the current contracts, and during 2025–2027, 61,449 ac under contracts in Douglas County will expire.

***Climate change and wildfire.*** Habitat changes as a result of climate changes affected sage-grouse abundance over the last 4,000 years (Wolfe and Broughton 2016), but the incidence of drought and wind events that increases wildfire size and frequency are of more immediate concern. Wildfires are the most immediate threat to sage-grouse in Washington (Stinson et al. 2004). Wildfires impact significant amounts of shrub-steppe annually (Coates et al. 2015). High severity fires eliminate sagebrush, and it can take > 10 years to recover adequate sagebrush cover to be suitable for sage-grouse, assuming enough nearby sagebrush survived to provide a seed source. Drought can greatly influence the risk of catastrophic fire, and some of the ignitions are due to human activities that are not closely regulated such as target shooting and burning of weeds. Gaps in fire district coverage can also lead to time delays in suppressing fires when they are small and more easily controlled. The emphasis on protecting infrastructure and directing fire resources to developed areas allows fires to expand. Efforts within Washington and rangewide to reduce the size of wildfires include establishing fuel breaks; their effectiveness may depend on sustained funding to maintain them over the long-term so that they remain effective, and firebreaks would be ineffective during extremely windy conditions, as in September 2020.

***Electrical transmission lines and renewable energy projects.*** Transmission lines have been negatively correlated with sage-grouse persistence and movements (Connelly et al. 2004, Beck et al. 2006, Wisdom et al. 2011, Schroeder and Vander Haegen 2014, Shirk et al. 2015, Gibson et al. 2018). Major transmission lines have a substantial footprint on the JBLM-YTC and in Douglas County; two radio-collared sage-grouse that moved from Lincoln County to Douglas County were found dead near transmission lines and were probably collision casualties. The impact of wind turbines is not clear; sage-grouse generally avoid tall structures, but in 2006, a female from the JBLM-YTC nested (unsuccessfully)

near a turbine of the Wild Horse project in the Colockum management unit. Recent proposals for several large solar energy projects in eastern Washington have the potential to further reduce and fragment shrub-steppe habitat, depending on location, and would likely need additional powerlines; WDFW has not yet received formal proposals.

***Livestock grazing and management.*** Livestock grazing is the most widespread land use occurring in sage-grouse range. Livestock grazing is compatible with sage-grouse where the habitat characteristics needed for breeding and wintering can be consistently maintained (Connelly et al. 2000a, 2011b; Wambolt et al. 2002, Crawford et al. 2004). The effects of livestock on sage-grouse habitat depend on stocking level, season of use, utilization levels, history of the site, and drought. The most immediate impact of grazing can be reduction of grass cover at sage-grouse nest sites, which can result in high rates of nest predation (Gregg et al. 1994, Hockett 2002, Rebolz 2007), although rotational grazing in Montana, and late season grazing in Wyoming did not have this effect (Monroe et al. 2017, Smith et al. 2017). Collisions with fencing constructed to manage livestock can cause direct mortality to sage-grouse (Stevens et al. 2012). Water developments can result in the degradation of important brood-rearing habitat by concentrating livestock and they may facilitate the spread of West Nile Virus by providing mosquito breeding sites (Walker and Naugle 2011). Raven occurrence is also associated with cattle (Coates et al. 2016). Sage-grouse population declines may be correlated with drought (Johnson et al. 2011), which can increase the negative effects of a grazing regime that might otherwise be sustainable. Though the range of sage-grouse still shows some effects of excessive historical grazing, in general, livestock management has improved, and ranching on private lands is less detrimental for sage-grouse than alternative land uses such as development of ranchettes or conversion to cropland. The Douglas County General Conservation Plan and the Sage-grouse Initiative (below in *Management Activities*) provide incentives for landowners to incur fewer impacts to sage-grouse on working ranch and farm lands.

Livestock grazing is currently not permitted on most of WDFW-managed lands within the Sage-grouse Management Units; there are seven grazing leases representing a small percentage of the acres in the Units, and only one of these leases is in occupied habitat. Grazing permits on WDFW managed land for periods of more than two weeks require livestock grazing management plans that include monitoring and schedules for evaluation. DNR owns more than 150,000 ac of land in sage-grouse PACs in Douglas, Lincoln, and Grant Counties, and 80,000 ac are leased for grazing. On public lands, grazing is monitored to ensure that the appropriate standards are met. On state lands, this is required by Ecosystem Standards for State-Owned Agricultural and Grazing Land (RCW 79.13.600, 79.13.610, and 77.12.204). BLM primarily implements deferred rotation grazing systems with conservative stocking rates (J. Lowe, pers. comm.). The rotations are set up to seasonally avoid grazing in areas that are most likely to support nesting birds during the breeding season. An assessment of sage-grouse habitat on the Twin Lakes allotment showed BLM land under grazing management is meeting breeding habitat requirements for nesting habitat indicators and is not significantly different than un-grazed control transects (BLM 2014a).

#### Other Factors Affecting Sage-grouse

***Predation.*** Predation is the most important proximate cause of mortality for sage-grouse, and the rate of predation is affected by the quality of habitat (Connelly et al. 2011a,b). Losses to predation are sustainable in large populations but have a more significant impact on small populations in fragmented habitat. Hagen (2011) suggested that areas with higher predator populations in human altered landscapes may be population sinks. He noted that short-term reduction of predators may be warranted during translocation projects, because translocated birds often suffer higher than normal rates of mortality (Hagen 2011). Habitat changes and human-associated food and water sources (e.g. roadkill, agriculture, landfills, stock ponds) and nesting and perching structures have generally increased the abundance of some important predators of sage-grouse eggs, chicks, and adults, particularly Common Ravens, Coyotes,

and Great Horned Owls (*Bubo virginianus*; Schroeder and Baydack 2001, Coates and Delehanty 2010, Stinson and Schroeder 2012, Schroeder et al. 2019). The population of ravens has increased ~400% in western North America in the past 40 years (Howe et al. 2014, Sauer et al. 2017, O’Neil et al. 2018). Coates et al. (2020) reported that ~64% of sage-grouse breeding areas in the Great Basin were adversely impacted by high raven density

**West Nile Virus.** West Nile Virus (WNV), a disease new to North America, has caused high mortality in sage-grouse populations in some locations (Naugle et al. 2005, Walker and Naugle 2011). It is transmitted between mosquitoes and birds; many infected birds die within 4–8 days, but if they survive, the antibodies may confer long-lasting protection from reinfection (Kilpatrick et al. 2007). In 2009, 4 mosquito samples from the JBLM-YTC tested positive for WNV, but effects of the disease on Washington grouse populations are unknown.

## MANAGEMENT ACTIVITIES

Stinson and Schroeder (2014) previously described conservation actions to address tasks in the state recovery plan for sage-grouse; updates of several activities are briefly described in annual reports (Schroeder et al. 2019) and mentioned below.

**Lincoln County reintroduction.** During 2008–2015, 277 sage-grouse from Oregon were released in Lincoln County (Schroeder et al. 2019). The movements, productivity, habitat use, and survival of these birds were monitored closely as part of a graduate research project (Stonehouse 2013, Stonehouse et al. 2015). Sage-grouse numbers were down after extreme conditions in winter 2016–2017 and the population has struggled since then. Additional releases to reinforce numbers were planned, but the Whitney Fire has likely eliminated the original location from consideration.

**JBLM-YTC augmentations and demographic, habitat, and predator studies.** A population augmentation effort was conducted to address genetic concerns associated with small population size of the JBLM-YTC population. During 2004–2006, 61 birds from Nevada or Oregon were released. Genetic analysis to determine if the augmentation was successful was inconclusive, and the augmentation effort resumed in 2014–16, with 36 females from Idaho (n=18) and Nevada (n=18). Ebenhoch et al. (2019) evaluated vital rates of resident vs. translocated birds and White et al. (unpublished data) assessed the spatial distribution and demographics of resident sage-grouse from 2012–2018. Sage-grouse habitat models, and predator assessments and management plans were also completed (Vernadero Group 2012, White and Lannoye 2014, Lannoye and White 2014a,b, Harris Environmental 2015).

**Yakama Nation reintroduction.** The Yakama Nation attempted to re-establish a population on the Yakama Reservation in the Toppenish Ridge SMU. A total of 155 sage-grouse from Oregon, Nevada, and Wyoming were released from 2005–2006, and 2013–2014. However, this incipient population struggled; a single male was present on the lek in 2018, and none were present in 2019.

**Wildfire suppression and prevention.** The threat of wildfires is being addressed in several ways. The JBLM-YTC has done substantial planning and established the capacity for aggressive fire suppression. Although initially effective upon implementation from 2010-2013, lack of consistent application of wildland fire management actions starting in 2014 has resulted in significant impacts to habitat both on and off the installation. The Lincoln County Conservation District initiated a fuel breaks project in 2015, and BLM began a project in 2019, mowing 9 miles; to date ~15-20 miles have been established. Swanson Lakes WLA staff are working with the District and BLM to develop a maintenance plan so that the firebreaks will remain effective. There is also proposed legislation that would remedy the gaps on the landscape where primary responsibility for fire suppression is not clear.



**Habitat restoration.** Restoration projects have had varied success at reestablishing sage-grouse habitat due to differences in precipitation levels, soil quality, invasion by non-native species, and the length of time required (Arkle et al. 2014). JBLM-YTC has ongoing sagebrush restoration to address areas affected by past wildfires and chronic training impacts; however, given the time required for sage-brush regrowth, the efforts are not meeting the need given the rate and scale that wildland fire impacts are occurring. Many other habitat restoration projects in sage-grouse management units have been completed or are in various stages of completion on WDFW and BLM lands (Stinson and Schroeder 2014, Schroeder et al. 2019). Since 1995, >3,000 ac of former cropland had been restored in Lincoln County, though it isn't clear how much will recover from the recent fire. The funding required to restore suitable habitat conditions at the needed scale limits the amount and pace of restoration work that can occur.

**Fence marking and fence and perch structure removal.** Fence collisions can be a major source of mortality for sage-grouse, although making fences more visible can dramatically reduce collision risk (Stevens et al. 2012, Van Lanen et al. 2017). Many miles of fencing in SMUs have been marked by BLM, WDFW, JBLM -YTC, conservation districts, partner organizations and volunteers, and many miles of unneeded fences have been removed. In Lincoln County, 7.7 miles of power distribution line and telephone lines were removed eliminating 126 poles, 2 miles were installed below-ground, perch deterrents have been installed on 68 powerpoles on the central distribution line in the reintroduction area, and other structures used as predator perches (e.g. a wind mill) have also been taken down.

**Sage-grouse Initiative (SGI).** The Natural Resource Conservation Service (NRCS), in partnership with Pheasants Forever, initiated the Sage-grouse Initiative (SGI) in 2010. Ranchers in sage-grouse habitat areas of central and eastern Washington may be eligible to receive financial assistance to help protect habitat and improve range conditions for their livestock for existing grazed ranch lands and expired CRP lands as well as restore marginal crop land back to native habitat. SGI uses farm bill dollars to implement conservation practices that benefit both their daily operations as well as sage-grouse and other wildlife species. SGI typically funds infrastructure such as pipelines, troughs, wells and fencing to develop rest/deferred-rotational grazing systems on private lands, and requires marking of fences in sage-grouse areas. Other common practices include: escape ramps; native seedings; obstruction removal; pollinator plots; cover crops; shrub plantings; and riparian restoration. Since 2010, 120,547 ac of private/public land have been enrolled in 135 SGI contracts in Washington with an overall investment over 6 million dollars, primarily in Douglas County. In 2020, the Chelan-Douglas Land Trust submitted application to NRCS for protecting approximately 6,724 acres of rangeland and dry cropland from development in southern Douglas County that was eligible for special funding through SGI. The easement represents the first SGI-funded Agricultural Land Easement (ALE) that NRCS completed in Washington, and the single largest ALE in the state. Over the next several years, NRCS hopes to work more closely with WDFW to help with connectivity between sage-grouse populations within the state.

**Douglas County General Conservation Plan.** The Foster Creek Conservation District (2014) developed a Multiple Species General Conservation Plan (GCP) for Douglas County that includes sage-grouse as a covered species. A GCP is a programmatic Habitat Conservation Plan under which multiple Section 10 permits (of the federal Endangered Species Act) can be granted for "incidental take" for otherwise lawful activities. The GCP describes a process for applicants (private agriculture landowners) to develop voluntary site-specific farm plans that will result in improved habitat for covered species while gaining long-term assurances for their agriculture operations. The GCP has grazing guidelines for developing grazing management plans that would promote better habitat and improve plant productivity and vigor, seed production, photosynthesis, recovery, and re-growth.

**Coordination and cooperation.** Sage-grouse in Washington are the focus of an interagency 'tech team', and a larger working group, that help coordinate conservation actions and identify priorities for

conservation actions. The tech team recently assisted with the development of a guidance document for grouse translocations, that is being used as the foundation for a Western Association of Fish and Wildlife Agencies (WAFWA) whitepaper.

## Research

***Landscape connectivity.*** Re-establishing connections between populations that are now isolated is vital for the long-term viability of sage-grouse populations in Washington. Robb and Schroeder (2012) modeled habitat concentration areas and movement corridors for sage-grouse to help prioritize protection and restoration of key linkage habitat. Shirk et al. (2015) used data from telemetry, genetics, and leks to evaluate the expert opinion models used in the analysis to predict rates of movement, gene flow, and lek persistence.

***Sage-grouse male movements and post-fire habitat use.*** Movements of sage-grouse were analyzed before and after the Apache Pass Fire in Lincoln County (Wells et al. 2016). Thornton and Olsoy (2018) analyzed the movements and resource use by translocated and resident male sage-grouse in Lincoln County.

***Survival and habitat use by translocated sage-grouse.*** Ebenhoch (2017) and Ebenhoch et al. (2019) compared movements, survival and reproduction of translocated and resident sage-grouse on JBLM-YTC. Stonehouse et al. (2015) analyzed habitat use by the sage-grouse translocated to Lincoln County.

## CONCLUSION AND RECOMMENDATION

Greater Sage-grouse are now limited in distribution to two main areas centered in Douglas County and on JBLM-YTC; the small number of birds in Lincoln County will likely be extirpated by 2021. Due to the sage-grouse polygynous mating system, the effective sizes of Washington's populations were not sustainable. The spring 2020 total estimate was 775 birds, up slightly from 688 in 2019, but the devastating fires in September 2020 may result in a ~50% population decline. The full impact may not be known for 1–2 years. Efforts are underway by various partner agencies to reduce the size of wildfires, but they still pose the greatest threat to habitat and any efforts will not be able to prevent large fires during extreme conditions as existed in September 2020.

The Douglas County sage-grouse population is unique in that Farm Bill conservation programs have been essential for maintaining it, but recent changes to Farm Bill programs create great uncertainty going forward. Without the Farm Bill programs and other conservation efforts, sage-grouse will likely decline to extinction in Washington.

Sage-grouse will likely decline well below the level indicated (<650 birds) in the state recovery plan for up-listing (Stinson et al. 2004). That objective, which was roughly derived from the 50/500 effective population 'rule' and the estimate of an  $N/N_e$  ratio of 0.156, assumed the populations were connected, but they appear to be functionally isolated. Conservation science has increasingly emphasized the negative effects of fragmentation and small population size for genetic health and fitness, particularly given the effect of the polygynous mating system on effective population size. The  $N_e$  of the Douglas County, JBLM-YTC, and Lincoln County populations were 108, 10, and 2 birds, respectively, before the September fires. The Lincoln County population may now be effectively extinct, and the JBLM-YTC population has continued a steady decline.

Concurrent with the troubling decline, genomic analysis has indicated that Washington's population is more distinct than the Bi-state population in California and Nevada which has been proposed for listing as a threatened DPS under the Endangered Species Act (USFWS 2019). Oh et al. (2019) suggested that Washington's highly differentiated population may possess important adaptations and "may warrant recognition and protection as a genetically distinct conservation unit." Macdonald et al. (2017) suggested that peripheral populations of species may be a valuable resource for translocation to augment vulnerable populations facing rapid ecological and climatic changes elsewhere. Washington's sage-grouse populations would first need to recover from the current population bottleneck before any such translocation would be considered, and they continue to lose genetic diversity as long as populations remain small.

For these reasons, we recommend that the Greater Sage-grouse be up-listed to endangered in Washington.

## REFERENCES CITED

The references cited in the Periodic *Status Review for the Greater Sage-grouse* are categorized for their level of peer review pursuant to section 34.05.271 RCW, which is the codification of Substitute House Bill 2661 that passed the Washington Legislature in 2014. A key to the review categories under section 34.05.271 RCW is provided in Table A.

**Table A. Key to 34.05.271 RCW Categories:**

Category Code	34.05.271(1)(c) RCW
i	(i) Independent peer review: review is overseen by an independent third party.
ii	(ii) Internal peer review: review by staff internal to the department of fish and wildlife.
iii	(iii) External peer review: review by persons that are external to and selected by the Department of Fish and Wildlife.
iv	(iv) Open review: documented open public review process that is not limited to invited organizations or individuals.
v	(v) Legal and policy document: documents related to the legal framework for the significant agency action including but not limited to: (A) federal and state statutes; (B) court and hearings board decisions; (C) federal and state administrative rules and regulations; and (D) policy and regulatory documents adopted by local governments.
vi	(vi) Data from primary research, monitoring activities, or other sources, but that has not been incorporated as part of documents reviewed under the processes described in (c)(i), (ii), (iii), and (iv) of this subsection.
vii	(vii) Records of the best professional judgment of Department of Fish and Wildlife employees or other individuals.
viii	(viii) Other: Sources of information that do not fit into one of the categories identified in this subsection (1)(c).

Reference	Category
Arkle, R.S., D.S. Pilliod, S.E. Hanser, M.L. Brooks, J.C. Chambers, J.B. Grace, K.C. Knutson, D.A. Pyke, J.L. Welty, and T.A. Wirth. 2014. Quantifying restoration effectiveness using multi-scale habitat models: implications for sage-grouse in the Great Basin. <i>Ecosphere</i> 5(3):31. <a href="http://dx.doi.org/10.1890/ES13-00278.1">http://dx.doi.org/10.1890/ES13-00278.1</a>	i
Beck, J.L., K.P. Reese, J. W. Connelly, and M. B. Lucia. 2006. Movement and survival of juvenile greater sage-grouse in southwestern Idaho. <i>Wildlife Society Bulletin</i> 34:1070-1078.	i
Bergerud, A.T. 1988. Population ecology of North American grouse. Pp 578-648 in A. T. Bergerud and M. W. Gratson, eds. <i>Adaptive strategies and population ecology of northern grouse</i> . University of Minnesota Press., Minneapolis.	i
BLM (Bureau of Land Management). 2014a. Habitat Assessment for Greater Sage-grouse in the Upper Crab Creek Habitat Concentration Area. Unpublished report. Bureau of Land Management, Spokane District, Border Field Office. Spokane, WA. 109 p.	viii
BLM (Bureau of Land Management). 2017. Record of Decision for the Vantage to Pomona Heights 230 kV Transmission Line Project. Case File: WAOR 65753. USDI BLM, Spokane, WA.	v
Coates, P.S. and D.J. Delehanty 2010. Nest predation of greater sage-grouse in relation to microhabitat factors and predators. <i>Journal of Wildlife Management</i> 74(2): 240–248.	i
Coates, P.S., M.A. Ricca, B.G. Prochazka, K.E. Doherty, M.L. Brooks, and M.L. Casazza. 2015. Long-term effects of wildfire on greater sage-grouse—Integrating population and ecosystem concepts for management in the Great Basin: U.S. Geological Survey Open-File Report 2015–1165, 42 p., <a href="http://dx.doi.org/10.3133/ofr20151165">http://dx.doi.org/10.3133/ofr20151165</a> .	viii

Reference	Category
Coates, P.S., B.E. Brussee, K.B. Howe, K.B. Gustafson, M.L. Casazza, and D.J. Delehanty. 2016. Landscape characteristics and livestock presence influence common ravens: relevance to greater sage-grouse conservation. <i>Ecosphere</i> 7(2): 1-20.	i
Coates, P. S., S. T. O'Neil, B. E. Brussee, M. A. Ricca, P. J. Jackson, J. B. Dinkins, K. B. Howe, A. M. Moser, L. J. Foster, and D. J. Delehanty. 2020. Broad-scale impacts of an invasive native predator on a sensitive native prey species within the shifting avian community of the North American Great Basin. <i>Biological Conservation</i> 243 (108409):1-10.	i
Connelly, J.W., K.P. Reese, R.A.Fischer, and W.L.Wakkinen. 2000a. Response of a sage grouse breeding population to fire in southeastern Idaho. <i>Wildlife Society Bulletin</i> 28(1):90-96.	i
Connelly, J.W., M.A. Schroeder, A.R. Sands, C.E. Braun. 2000b. Guidelines to manage sage grouse populations and their habitats. <i>Wildlife Society Bulletin</i> 28:967-985.	i
Connelly, J.W., S.T. Knick, M.A. Schroeder, and S. J. Stiver. 2004. Conservation assessment of greater sage-grouse and sagebrush habitats. Unpublished Report, Western Association of Fish and Wildlife Agencies. Cheyenne, WY. 610 pp.	i
Connelly, J. W., E.T. Rinkes, and C.E. Braun. 2011a. Characteristics of Greater Sage-grouse Habitats: a landscape species at micro- and macroscales. Pp. 69-83, in S. T. Nick and J. W. Connelly (eds.). Greater Sage-grouse: ecology and conservation of a landscape species and its habitats. <i>Studies in Avian Biology</i> vol.38. University of California Press, Berkeley, California. 646 pp.	
Connelly, J.W., C.A. Hagen, and M.A. Schroeder. 2011b. Characteristics and Dynamics of Greater Sage-grouse Populations. Pp. 53-67, in S.T. Nick and J.W. Connelly (eds.). Greater Sage-grouse: ecology and conservation of a landscape species and its habitats. <i>Studies in Avian Biology</i> vol.38. University of California Press, Berkeley, California. 646 pp.	i
Crawford, J.A., R.A. Olson, N.E. West, J.C. Mosley, M.A. Schroeder, T. D. Whitson, R. F.. Miller, M. A. Gregg, and C. S. Boyd. 2004. Ecology and management of sage-grouse and sage-grouse habitat. <i>Journal of Range Management</i> 57: 2-19.	i
Ebenhoch, K.G. 2017. Comparing population vital rates of resident and translocated Greater Sage-grouse on the Yakima Training Center, Yakima, WA. M.S. thesis. Washington State University, Pullman, Washington. 61 pp.	i
Ebenhoch, K., D. Thornton, L. Shipley, J.A. Manning, and K. White. 2019. Effects of post-release movements on survival of translocated Sage-Grouse. <i>Journal of Wildlife Management</i> 83(6):1314-1325.	i
Foster, L.J., K.M. Dugger, C.A. Hagen, and D.A. Budeau. 2019. Greater Sage-grouse vital rates after Wildfire. <i>Journal of Wildlife Management</i> 83(1):121–134.	i
Foster Creek Conservation District. 2014. Multiple species general conservation plan for Douglas County, Washington. Foster Creek Conservation District, Waterville, Washington. 379 pp	viii
Frankham, R. 1995. Effective population size/adult population size ratios in wildlife: a review. <i>Genetical Research</i> 66:95–107.	i
Garton, E.O., A.G.Wells, J. A. Baumgardt, and J.W. Connelly. 2015. Greater sage-grouse population dynamics and probability of persistence. Final Report to Pew Charitable Trusts. – <a href="http://www.pewtrusts.org/~media/assets/2015/04/garton-et-al-2015-greater-sagegrouse-population-dynamics-and-persistence-31815.pdf">http://www.pewtrusts.org/~media/assets/2015/04/garton-et-al-2015-greater-sagegrouse-population-dynamics-and-persistence-31815.pdf</a> .	viii
Gibson, D., E. J. Blomberg, M. T. Atamian, S. P. Espinosa, and J. S. Sediger. 2018. Effects of power lines on habitat use and demography of Greater Sage-Grouse ( <i>Centrocercus urophasianus</i> ). <i>Wildlife Monographs</i> 200:1–41.	i
Gilpin, M. E., and M. E. Soule. 1986. Minimum viable populations: processes of extinction. Pages 19-34 in M. Soule, editor. <i>Conservation Biology: The Science of Scarcity and Diversity</i> . Sinauer Associates, Sunderland, MA.	i
Gregg, M.A., and J.A. Crawford. 2009. Survival of Greater Sage-grouse Chicks and Broods in the Northern Great Basin. <i>Journal of Wildlife Management</i> 73(6):904–913.	i
Gregg, M.A., J.A. Crawford, M.S. Drut, and A.K. DeLong. 1994. Vegetational cover and predation of sage-grouse nests in Oregon. <i>Journal Wildlife Management</i> 58:162-166.	i
Hagen, C.A. 2011. Predation on Greater Sage-grouse: facts, process, and effects. Pp. 95-100, in S. T. Nick and J. W. Connelly (eds.). Greater Sage-grouse: ecology and conservation of a landscape	i

Reference	Category
species and its habitats. Studies in Avian Biology vol.38. University of California Press, Berkeley, California. 646 pp.	
Hagen, C.A., J.W. Connelly, and M.A. Schroeder. 2007. A meta-analysis of greater sage-grouse <i>Centrocercus urophasianus</i> nesting and brood-rearing habitats. Wildlife Biology 13(suppl. 1):42-50.	i
Harris Environmental Group. 2015. Yakima Training Center's Greater Sage-grouse predator assessment and management plan (phase III) Final Report. Contract #W912DW-14-D-1013.	viii
Hockett, G.A. 2002. Livestock impacts on the herbaceous components of sage grouse habitat: a review. Intermountain Journal of Sciences 8:105-114.	i
Howe, K.B., P.S. Coates, and D.J. Delehanty. 2014. Selection of anthropogenic features and vegetation characteristics by nesting Common Ravens in the sagebrush ecosystem. Condor 116:35-49.	i
Jamison, B.E., and M.F. Livingston. 2004. Sage-grouse evaluation in the shrub steppe ecosystem of the Yakama Reservation, Washington. Project Final Report, Interagency Agreement No. GP00065800. Yakama Nation Wildlife Resource Management, Toppenish, Washington. 343 pp.	viii
Johnson, D.H., M.J. Holloran, J.W. Connelly, S.E. Hanser, C.L. Amundson, and S.T. Knick. 2011. Influences of environmental and anthropogenic features on Greater Sage-grouse populations, 1997-2007. Pp. 407-450, in S.T. Nick and J.W. Connelly (eds.). Greater Sage-grouse: ecology and conservation of a landscape species and its habitats. Studies in Avian Biology vol.38. University of California Press, Berkeley, California. 646 pp.	i
Kelsey, R.G., J.R. Stephens, F. Shafizadeh. 1982. The chemical constituents of sagebrush foliage and their isolation. J Range Manage. 35(5):617-622.	i
Kilpatrick, A.M., S.L. LaDeau, and P.P. Marra. 2007. Ecology of West Nile virus transmission and its impact on birds in the western hemisphere. Auk 124(4):1121-1136.	i
Lannoye, J., and K. White. 2014a. Yakima Training Center's 2014 Sage-grouse predator assessment. Stell Environmental.	viii
Lannoye, J., and K. White. 2014b. Yakima Training Center's 2014 Sage-grouse habitat model and assessment. Stell Environmental. 104 pp.	viii
Macdonald, S.L., J. Llewelyn, C. Moritz, and B.L. Phillips. 2017. Peripheral isolates as sources of adaptive diversity under climate change. Frontiers in Ecology and Evolution 5:88	i
Monroe, A.P., C.L. Aldridge, T.J. Assal, K.E. Veblen, D.A. Pyke, and M.L. Casazza. 2017. Patterns in greater sage-grouse population dynamics correspond with public grazing records at broad scales: Ecological Applications 27(4): 1096-1107.	i
Moynahan, B.J., M.S. Lindberg, J.J. Rotella, and J.W. Thomas. 2007. Factors affecting nest survival of Greater Sage-Grouse in northcentral Montana. Journal Wildlife Management 71:1773-1783.	i
Naugle, D.E., C.L. Aldridge, B.L. Walker, K.E. Doherty, M.R. Matchett, J. McIntosh, T.E. Cornish, M.S. Boyce. 2005. West Nile virus and sage-grouse: What more have we learned? Wildlife Society Bulletin 33(2): 616-623.	i
Oh, K.P., C.L. Aldridge, J.S. Forbet, C.Y. Dadabay, and S.J. Oyler-McCance. 2019. Conservation genomics in the sagebrush sea: population divergence, demographic history, and local adaptation in sage-grouse ( <i>Centrocercus</i> spp.). Genome Biology and Evolution 11(7):2023-2034.	i
O'Neil, S.T., P.S. Coates, B.E. Brussee, P.J. Jackson, K.B. Howe, A.M. Moser, L. J. Foster, and D. J. Delehanty. 2018. Broad-scale occurrence of a subsidized avian predator: reducing impacts of ravens on sage-grouse and other sensitive prey. J. Applied Ecology 00:1-12.	i
PacifiCorp. n.d. <a href="https://www.pacificorp.com/transmission/transmission-projects/vantage-to-pomona-heights.html">https://www.pacificorp.com/transmission/transmission-projects/vantage-to-pomona-heights.html</a> .	viii
Rebholz, J. L. 2007. Influence of habitat characteristics on Greater Sage-grouse reproductive success in the Montana Mountains, Nevada. M. S. Thesis, Oregon State University, Corvallis. 64 pp.	viii
Ritchie, M.E., M.L. Wolfe, and R. Danvir. 1994. Predation of artificial sage grouse nests in treated and untreated sagebrush. Great Basin Naturalist 54:122-129.	i
Robb, L. and M.A. Schroeder. 2012. Appendix A.2 Habitat Connectivity for Greater Sage-grouse ( <i>Centrocercus urophasianus</i> ) in the Columbia Plateau Ecoregion. In, Washington Wildlife Habitat Connectivity Working Group. Washington Connected Landscapes Project: Analysis of the Columbia Plateau Ecoregion. Washington's Department of Fish and Wildlife, and	iii

Reference	Category
Department of Transportation, Olympia, WA. Available from <a href="http://waconnected.org/columbia-plateau-ecoregion/">http://waconnected.org/columbia-plateau-ecoregion/</a>	
Sauer, J.R., D.K. Niven, J.E. Hines, D.J. Ziolkowski, Jr, K.L. Pardieck, J.E. Fallon, and W.A. Link. 2017. The North American Breeding Bird Survey, Results and Analysis 1966 - 2015. Version 2.07.2017 USGS Patuxent Wildlife Research Center, Laurel, MD.	viii
Schroeder, M.A. 1997. Unusually high reproductive effort by Sage Grouse in a fragmented habitat in north-central Washington. <i>Condor</i> 99:933-941.	i
Schroeder, M.A. 2000. Population dynamics of greater and Gunnison sage-grouse: a review. Job Progress Report. Upland Bird Research. Washington Department of Fish and Wildlife.	vii
Schroeder, M.A., and R. K. Baydack. 2001. Predation and the management of prairie grouse. <i>Wildlife Society Bulletin</i> 29:24-32.	i
Schroeder, M.A., and W.M. Vander Haegen. 2006. Use of Conservation Reserve Program fields by Greater Sage-grouse and other shrubsteppe-associated wildlife in Washington state. Technical report prepared for U.S. Department of Agriculture Farm Service Agency. Washington Department of Fish and Wildlife, Olympia, Washington.	vii
Schroeder, M.A., and M. Vander Haegen. 2011. Response of Greater Sage-grouse to the Conservation Reserve Program in Washington State. Pages 517-529 in S. T. Knick and J. W. Connelly, (eds.). Greater Sage-grouse: ecology and conservation of a landscape species and its habitats. <i>Studies in Avian Biology</i> No. 38.	i
Schroeder, M.A., and W.M. Vander Haegen. 2014. Monitoring of Greater Sage-grouse and other breeding birds on the Winthrop Wind Power Project Site. Final Report. Washington Department of Fish and Wildlife, Olympia, WA. 26 pp.	vii
Schroeder, M.A., J.R. Young, and C.E. Braun. 1999. Sage grouse ( <i>Centrocercus urophasianus</i> ). In A. Poole and F. Gill, (eds.). <i>The Birds of North America</i> , No. 425. The Academy of Natural Sciences, Philadelphia, Pennsylvania, and the American Ornithologists' Union, Washington, D.C.	i
Schroeder, M.A., M. Atamian, J. Lowe, K. Thorburn, M. Finch, J. Anderson, K. White, E. Braaten, and D.W. Stinson. 2019. Recovery of Greater Sage-grouse in Washington: progress report and proposal. Washington Department of Fish and Wildlife, Olympia, Washington.	vii
Schroeder, M.A., D. Stinson, and M. Tirhi. 2003. Greater Sage-grouse ( <i>Centrocercus urophasianus</i> ). <i>Priority Habitat and Species Management Recommendations Vol. IV: Birds</i> . 19 pp.	iii
Shirk, A J., M.A. Schroeder, L A. Robb. 2015. Empirical validation of landscape resistance models: insights from Greater Sage-grouse ( <i>Centrocercus urophasianus</i> ). <i>Landscape Ecology</i> 15 May 2015. <a href="http://dx.doi.org/10.1007/s10980-015-0214-4">http://dx.doi.org/10.1007/s10980-015-0214-4</a>	i
Smith, J.T., J.D. Tack, L.I. Berkeley, M. Szczypinski, and D.E. Naugle. 2017. Effects of rotational grazing management on nesting Greater Sage-grouse. <i>J. Wildlife Management</i> 82(1):103-112.	i
Stevens, B.S., K.P. Reese, J.W. Connelly, and D.D. Musil. 2012. Greater Sage-grouse and fences: does marking reduce collisions? <i>Wildlife Society Bulletin</i> 36(2):297-303.	i
Stinson, D.W., and M.A. Schroeder. 2012. Washington State Recovery Plan for the Columbian Sharp-tailed Grouse. Washington Department of Fish and Wildlife, Olympia. 159+x pp.	iii, iv
Stinson, C. M., and M. A. Schroeder. 2014. Sage-grouse Conservation in Washington: 2013. 48 pp.	ii
Stinson, D.W., D.W. Hays, and M.A. Schroeder. 2004. Washington State recovery plan for the Greater Sage-grouse. Washington Department of Fish and Wildlife, Olympia, Washington. 109 pp.	iii, iv
Stonehouse, K.F. 2013. Habitat selection by sympatric translocated greater sage-grouse and Columbian sharp-tailed grouse in eastern Washington. M.S. thesis, Washington State University, Pullman, Washington.	viii
Stonehouse, K.F., L.A. Shipley, J. Lowe, M.T. Atamian, M E. Swanson, and M.A. Schroeder. 2015. Habitat selection and use by sympatric, translocated Greater Sage-grouse and Columbian sharp-tailed Grouse. <i>Journal of Wildlife Management</i> 79(8):1308-1326.	i
Sveum, C.M., J.A. Crawford, and W. D. Edge. 1998. Use and selection of brood-rearing habitat by sage-grouse in south central Washington. <i>Great Basin Naturalist</i> 58:344-351.	i
Thornton, D., and P. Olsoy. 2018. Characterizing movement patterns of sage-grouse through high-resolution GPS telemetry. 2016-2017 Annual Report.	vi
USFWS (U.S. Fish and Wildlife Service). 2010. <i>Endangered and Threatened Wildlife and Plants</i> ; 12-	v

Reference	Category
Month Finding for Petitions to List the Greater Sage-grouse ( <i>Centrocercus urophasianus</i> ) as Threatened or Endangered. Federal Register Vol. 75, No. 55 (March 23):13910-14012.	
USFWS (U.S. Fish and Wildlife Service). 2015. Endangered and Threatened Wildlife and Plants; 12-Month Finding on a Petition to List Greater Sage-grouse ( <i>Centrocercus urophasianus</i> ) as an Endangered or Threatened. Federal Register Vol. 80, No. 191 (October 2):59858-59942.	v
USFWS (U.S. Fish and Wildlife Service). 2019. Endangered and Threatened Wildlife and Plants; Threatened status for the Bi-State Distinct Population Segment of Greater Sage-grouse and designation of critical habitat: proposed rule. Federal Register Vol. 84, No. 71 (April 12):14909-14910.	v
Van Lanen, N.J., A.W. Green, T.R. Gorman, L.A. Quattrini, and D.C. Pavlacky, Jr. 2017. Evaluating efficacy of fence markers in reducing greater sage-grouse collisions with fencing: Biological Conservation 213:70–83.	i
Vernadero Group. 2012. Final 2012 Sage Grouse habitat assessment and predator survey at Joint Base Lewis McChord Yakima Training Center. Prepared for US Army Environmental Command, by Vernadero Group, Inc., Phoenix, Arizona.	viii
Walker, B.L., and D.E. Naugle. 2011. West Nile Virus ecology in sagebrush habitat and impacts on Greater Sage-grouse populations. Pp. 127-142, in S. T. Nick and J. W. Connelly (eds.). Greater Sage-grouse: ecology and conservation of a landscape species and its habitats. Studies in Avian Biology vol.38. University of California Press, Berkeley, California. 646 pp.	i
Wambolt, C.L., A.J. Harp, B.L. Welch, N. Shaw, J.W. Connelly, K.P. Reese, C.E. Braun, D.A. Klebenow, E.D. McArthur, J.G. Thompson, L.A. Torell, J.A. Tanaka. 2002. Conservation of greater sage-grouse on public lands in the western U.S.: implications of recovery and management policies. PACWPL-Policy Paper SG- 02-02. Policy Analysis Center for Western Public Lands, Caldwell, ID. 41 pp..	viii
Welch, B.L. 2005. Big sagebrush: A sea fragmented into lakes, ponds, and puddles. Gen. Tech Rep. RMRS-GTR-144. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 210 p.	i
Wells, A.G., J. Lowe, L. Shipley, and M. Atamian. 2016. The influence of fire on patterns of habitat selection by sage-grouse in the channeled scablands of Lincoln County, Washington. Final Report.	viii
White, K. and J. Lannoye. 2014. Yakima Training Center’s Sage-grouse predator management plan (phase II). Stell Environmental.	viii
Wisdom, M.J., C. W. Meinke, S.T. Knick, and M.A. Schroeder. 2011. Factors associated with extirpation of sage-grouse. Pp. 451-472, in S. T. Nick and J. W. Connelly (eds.). Greater Sage-grouse: ecology and conservation of a landscape species and its habitats. Studies in Avian Biology vol.38. University of California Press, Berkeley, California. 646 pp.	i
Wolfe, A.L., and J.M. Broughton. 2016. Bonneville Basin Avifaunal Change at the Pleistocene/Holocene Transition: Evidence from Homestead Cave. Ch. 14, p 371–419, in Charles G. Oviatt, John F. Shroder (editors) Developments in Earth Surface Processes, Vol. 20. ISSN 0928-2025, ISBN 9780444635907, <a href="http://dx.doi.org/10.1016/B978-0-444-63590-7.00014-7">http://dx.doi.org/10.1016/B978-0-444-63590-7.00014-7</a>	i



## PERSONAL COMMUNICATIONS

David Blodgett III, Wildlife Biologist  
Wildlife Resource Management  
Yakama Nation,  
Toppenish, Washington

Grant Casady, Associate Professor  
Director of Environmental Studies  
Whitworth University

Jason Lowe, Biologist  
USDI Bureau of Land Management  
Spokane District Office,  
Spokane, Washington

Ellie Mangelinckx  
Whitetail Environmental, LLC  
U.S. Army -Yakima Training Center, Environmental Division  
970 Firing Center Rd, Bldg. 810  
Yakima, Washington

Kevin White  
Whitetail Environmental, LLC  
U.S. Army -Yakima Training Center, Environmental Division  
970 Firing Center Rd, Bldg. 810  
Yakima, Washington

## APPENDIX A. PUBLIC COMMENTS

WDFW received public comments during the 90-day public review period for the draft *Status Review for the Greater Sage-grouse*. WDFW received 1,262 individual comment letters from citizens (most were copied form letters), organizations, and Douglas County; 1,259 response letters indicated support for WDFW's status recommendation to uplist the Greater Sage-grouse to an endangered species in Washington and for continued recovery actions.



# WASHINGTON STATE STATUS REPORTS, PERIODIC STATUS REVIEWS, RECOVERY PLANS, AND CONSERVATION PLANS

## Periodic Status Reviews

2021	Steller Sea Lion
2021	Greater Sage-grouse
2021	Gray Whale
2021	Humpback Whale
2020	Mazama Pocket Gopher
2019	Tufted Puffin
2019	Oregon Silverspot
2018	Grizzly Bear
2018	Sea Otter
2018	Pygmy Rabbit
2017	Fisher
2017	Blue, Fin, Sei, North Pacific Right, and Sperm Whales
2017	Woodland Caribou
2017	Sandhill Crane
2017	Western Pond Turtle
2017	Green and Loggerhead Sea Turtles
2017	Leatherback Sea Turtle
2016	American White Pelican
2016	Canada Lynx
2016	Marbled Murrelet
2016	Peregrine Falcon
2016	Bald Eagle
2016	Taylor's Checkerspot
2016	Columbian White-tailed Deer
2016	Streaked Horned Lark
2016	Killer Whale
2016	Western Gray Squirrel
2016	Northern Spotted Owl
2016	Snowy Plover

## Conservation Plans

2013	Bats
------	------

## Recent Status Reports

2021	Oregon Vesper Sparrow
2019	Pinto Abalone
2017	Yellow-billed Cuckoo
2015	Tufted Puffin
2007	Bald Eagle
2005	Mazama Pocket Gopher, Streaked Horned Lark, and Taylor's Checkerspot
2005	Aleutian Canada Goose
1999	Northern Leopard Frog
1999	Mardon Skipper
1999	Olympic Mudminnow
1998	Margined Sculpin
1998	Pygmy Whitefish
1997	Gray Whale
1997	Olive Ridley Sea Turtle
1997	Oregon Spotted Frog

## Recovery Plans

2020	Mazama Pocket Gopher
2019	Tufted Puffin
2012	Columbian Sharp-tailed Grouse
2011	Gray Wolf
2011	Pygmy Rabbit: Addendum
2007	Western Gray Squirrel
2006	Fisher
2004	Sea Otter
2004	Greater Sage-Grouse
2003	Pygmy Rabbit: Addendum
2002	Sandhill Crane
2001	Lynx
1999	Western Pond Turtle
1996	Ferruginous Hawk
1995	Pygmy Rabbit
1995	Snowy Plover

Status reports and plans are available on the WDFW website at:  
<http://wdfw.wa.gov/publications/search.php>



