

Tucannon River Spring Chinook Salmon Hatchery Evaluation Program 2021 Annual Report



by Michael P. Gallinat
and Dane E. Kiefel



*Washington Department of
Fish and Wildlife
Fish Program*

Tucannon River Spring Chinook Salmon Hatchery Evaluation Program

2021 Annual Report

by

Michael P. Gallinat
Dane E. Kiefel

Washington Department of Fish and Wildlife
Fish Program/Science Division
600 Capitol Way North
Olympia, Washington 98501-1091

Prepared for:

U.S. Fish and Wildlife Service
Lower Snake River Compensation Plan Office
1387 S. Vinnell Way, Suite 343
Boise, Idaho 83709
Cooperative Agreement: F220C00018-00

August 2022

Acknowledgments

The Tucannon River Spring Chinook Salmon Hatchery Evaluation Program is the result of efforts by many individuals within the Washington Department of Fish and Wildlife (WDFW) and from other agencies.

We would like to express our gratitude to Ace Trump, Lyons Ferry Hatchery Complex Manager and Hatchery Specialists Rianna Earl, Derek Gloyn, Doug Maxey, and Dan Pounds for their cooperation with hatchery sampling, providing information regarding hatchery operations and hatchery records, and their input on evaluation and research activities. We also thank all additional hatchery personnel who provide the day-to-day care of the spring Chinook and for their assistance with hatchery spawning, sampling, and record keeping.

We thank Andrew Claiborne for providing scale ages and Kelly Britt for providing information on fish health during the year. Special thanks go to David Bramwell for help formatting this report.

We thank the staff of the Snake River Lab; in particular, Joe Bumgarner, Jenna Fortier, Todd Miller, Lance Ross and seasonal workers Carson Alessio, Caleb Bak, Jeffery Christopherson, and Leah Spoolstra who helped collect the information presented in this report. We also sincerely thank Steve Richards and Shawna Meehan for their assistance with PIT tagging.

We thank Joe Bumgarner, Alf Haukenes, Rod Engle, and Laurie Peterson for reviewing the draft report.

The United States Fish and Wildlife Service through the Lower Snake River Compensation Plan Office funds the supplementation program. A grant through the Bonneville Power Administration provided funding for a portion of the hatchery program PIT tags.

Abstract

Lyons Ferry Hatchery (LFH) and Tucannon Fish Hatchery (TFH) were built/modified under the Lower Snake River Fish and Wildlife Compensation Plan. One objective of the Plan is to compensate for the estimated annual loss of 1,152 Tucannon River spring Chinook caused by hydroelectric projects on the Snake River. This report summarizes activities of the Washington Department of Fish and Wildlife Lower Snake River Hatchery Evaluation Program for Tucannon River spring Chinook for the period May 2021 to April 2022.

A total of 140 salmon were captured in the TFH trap in 2021 (75 natural adults, 8 natural jacks, 35 hatchery adults, and 22 hatchery jacks). Of these, 115 fish (83 natural, 32 hatchery) were collected for broodstock and 25 adipose clipped strays were killed outright. During 2021, three (2.6%) salmon collected for broodstock died prior to spawning.

Spawning of supplementation fish occurred once a week between 31 August and 28 September, with peak eggtake occurring on 14 September. A total of 166,237 eggs were collected from 38 natural and 9 hatchery-origin female Chinook. Egg mortality to eye-up was 11.9% (19,883 eggs) which left 146,354 live eggs. An additional 0.4% (647) loss of sac-fry left 145,707 BY 2021 fish for production.

Weekly spawning ground surveys began 26 August and were completed by 1 October 2021. A total of 35 redds and 15 carcasses (5 natural, 10 hatchery) were found. Four redds (11% of the total) were counted above the adult trap, even though fish were not intentionally passed upstream. All hatchery origin fish recovered during spawning ground surveys were strays and strays accounted for 41.4% of the run after expansions. Based on redd counts, carcasses recovered, and broodstock collection, the estimated return to the river for 2021 was 215 spring Chinook (101 natural adults, 7 natural jacks and 85 hatchery-origin adults, 22 hatchery jacks).

A total of 42,046 BY20 smolts were direct stream released at TFH on 11 April and 19,974 BY20 smolts were released at the mouth of the Tucannon River on 22 April for a total of 62,020 BY20 smolts released.

Evaluation staff operated a downstream migrant trap to provide juvenile outmigration estimates. During the 2020/2021 emigration, we estimated that 174 natural spring Chinook (BY 2019) smolts emigrated from 1 October 2020 to 6 July 2021 from the Tucannon River.

Smolt-to-adult return rates (SAR) for natural origin salmon are eight times higher on average (based on geometric means) than hatchery origin salmon. However, hatchery salmon survive three times greater than natural salmon from parent to adult progeny over the length of the project. Managers are currently discussing alternative hatchery rearing and release strategies (e.g., hatchery releases below Bonneville Dam, barge transportation, captive broodstock, etc.) in an attempt to increase hatchery fish survival and preserve this stock.

Table of Contents

| | |
|--|----|
| List of Tables | ii |
| List of Figures | iv |
| List of Appendices | v |
| Introduction..... | 1 |
| Program Objectives..... | 1 |
| ESA Permits..... | 2 |
| Facility Descriptions | 2 |
| Tucannon River Watershed Characteristics..... | 3 |
| Adult Salmon Evaluation..... | 5 |
| Broodstock Trapping | 5 |
| Broodstock Mortality | 7 |
| Broodstock Spawning | 8 |
| Broodstock BKD Screening and Virology Testing | 9 |
| Natural Spawning..... | 10 |
| Historical Trends in Natural Spawning..... | 11 |
| Stream Nutrient Enrichment | 14 |
| Genetic Sampling..... | 15 |
| Age Composition, Length Comparisons, and Fecundity | 15 |
| Arrival and Spawn Timing Trends | 20 |
| Total Run-Size | 22 |
| Spawning Escapement | 24 |
| Coded-Wire Tag Sampling | 26 |
| Stray Salmon into the Tucannon River..... | 27 |
| Adult PIT Tag Returns..... | 29 |
| Juvenile Salmon Evaluation..... | 31 |
| Hatchery Rearing, Marking, and Release | 31 |
| Smolt Trapping | 32 |
| Smolt Migration to Lower Monumental and McNary Dams..... | 35 |
| Survival Rates | 37 |
| Fishery Contribution and Out-of-Basin Straying..... | 46 |
| Adjusted Hatchery SAS | 48 |
| Alternative Rearing and Release Strategies..... | 49 |
| Conclusions and Recommendations | 54 |
| Literature Cited..... | 57 |

List of Tables

| | | |
|-----------|--|----|
| Table 1. | Description of five strata within the Tucannon River..... | 4 |
| Table 2. | Numbers of spring Chinook salmon captured at the TFH trap, trap mortalities, strays or jacks killed outright, fish collected for broodstock, and passed upstream or held for adult outplanting for natural spawning from 1986-2021 | 6 |
| Table 3. | Numbers of pre-spawning mortalities and percent of fish collected for broodstock at TFH and held at TFH (1985-1991 and 2019) or LFH (1992-2018 and 2020-2021) | 7 |
| Table 4. | Number of fish spawned or killed outright (K.O.), estimated egg collection, and egg mortality of natural and hatchery origin Tucannon River spring Chinook salmon at LFH in 2021 | 8 |
| Table 5. | Enzyme Linked Immunosorbent Assay (ELISA) values for hatchery spawned Tucannon River spring Chinook females, 2021 | 9 |
| Table 6. | Numbers and general locations of salmon redds and carcasses (includes pre-spawn mortalities) recovered on the Tucannon River spawning grounds, 2021 (the Tucannon Hatchery adult trap is located at rkm 59)..... | 10 |
| Table 7. | Number of spring Chinook salmon redds and redds/km (in parenthesis) by stratum and year, and the number and percent of redds above and below the TFH adult trap in the Tucannon River, 1985-2021 | 13 |
| Table 8. | Summary of Chinook salmon carcass distribution on the Tucannon River on 8 and 10 January 2022 | 15 |
| Table 9. | Average number of eggs/female (n, SD) by age group of Tucannon River natural and hatchery origin broodstock, 1990-2021 (partial spawned females are excluded) | 19 |
| Table 10. | Peak dates of arrival of natural and hatchery salmon to the TFH adult trap and peak (date) and duration (number of days) for spawning in the hatchery and river, 1986-2021..... | 21 |
| Table 11. | Estimated spring Chinook salmon run to the Tucannon River and recovered pre-spawn mortalities (PSM), 1985-2021 | 23 |
| Table 12. | Estimated spawning escapement and the calculation methodology used for the 1985 to 2021 run years..... | 25 |
| Table 13. | Coded-wire tag codes of hatchery salmon sampled at LFH and the Tucannon River, 2021..... | 26 |
| Table 14. | Spring Chinook salmon (natural and hatchery) sampled from the Tucannon River, 2021..... | 26 |
| Table 15. | Tucannon River PIT tag array detections of spring Chinook originally tagged at locations other than the Tucannon River during 2021 | 28 |
| Table 16. | Genetic stock assignment results of adults PIT tagged at Lower Granite Dam and entering the Tucannon River based on Genetic Stock Identification (GSI) and Parentage Based Tagging (PBT) during 2021 | 28 |

| | |
|---|----|
| Table 17. Number of Tucannon River spring Chinook juvenile fish PIT tagged by origin and calendar year and adult returns detected (%) in the Columbia River System by origin | 29 |
| Table 18. Number and origin of PIT tagged Tucannon River spring Chinook returns that overshot the Tucannon River (includes fish that were last detected returning downstream towards the Tucannon River) and also detected at Lower Granite Dam (LGR) that stayed above LGR Dam | 30 |
| Table 19. Sample size (N), mean length (mm), coefficient of variation (CV), condition factor (K), mean weight (g), and precocity of 2020 BY juveniles sampled at TFH..... | 31 |
| Table 20. Spring Chinook salmon released into the Tucannon River at TFH and the mouth of the Tucannon River, 2022 release year..... | 31 |
| Table 21. Median and mean travel time and outmigration speed of hatchery-origin Tucannon River spring Chinook to Lower Monumental and McNary Dams in 2021 | 35 |
| Table 22. Estimates of <i>natural in-river produced</i> Tucannon spring Chinook salmon (both hatchery and natural origin parents) abundance by life stage for 1985-2021 broods. | 38 |
| Table 23. Estimates of Tucannon spring Chinook salmon abundance (<i>spawned and reared in the hatchery</i>) by life stage for 1985-2021 broods..... | 39 |
| Table 24. Percent survival by brood year for juvenile salmon and the multiplicative advantage of hatchery-reared salmon over naturally-reared salmon in the Tucannon River | 40 |
| Table 25. Adult returns and SARs of natural salmon to the Tucannon River for brood years 1985-2018 | 41 |
| Table 26. Adult returns and SARs of hatchery salmon to the Tucannon River for brood years 1985-2018 | 42 |
| Table 27. Progeny-to-parent survival estimates of Tucannon River spring Chinook salmon from 1985 through 2017 brood years (2017 brood year incomplete)..... | 44 |
| Table 28. Summary of Tucannon River spring Chinook recovered outside of the Tucannon River and represent possible strays to other areas (2005-2017 brood years) | 47 |
| Table 29. Hatchery SAS adjusted for recoveries from outside the Tucannon River subbasin as reported in the RMIS database, 1985-2016 brood years..... | 48 |

List of Figures

| | | |
|------------|--|----|
| Figure 1. | Location of the Tucannon River, and Lyons Ferry and Tucannon Hatcheries within the Snake River basin..... | 3 |
| Figure 2. | Historical Below Low and Low, and Moderate and High ELISA values for Tucannon River spring Chinook salmon female broodstock for the 1998 to 2021 return years ... | 9 |
| Figure 3. | Spring Chinook redd density (redds/km) in the Tucannon River, 1986-2021..... | 11 |
| Figure 4. | The proportion of redds above Marengo that were either above the adult trap/weir or below the adult trap/weir with trend lines, 1985-1993 and 2000-2014..... | 12 |
| Figure 5. | Historical (1985-2020), and 2021 age composition (run year) for spring Chinook in the Tucannon River..... | 16 |
| Figure 6. | Weighted mean age of natural and hatchery origin males (NM, HM) and natural and hatchery origin females (NF, HF) for the 1985 to 2016 brood years for spring Chinook in the Tucannon River..... | 17 |
| Figure 7. | Mean post-orbital to hypural-plate (POH) length comparisons between age-4 natural and hatchery-origin males (NM and HM) and natural and hatchery-origin females (NF and HF) with 95% confidence intervals for the years 1985-2021..... | 18 |
| Figure 8. | Cumulative run timing by date at the Tucannon Fish Hatchery adult trap on the Tucannon River for both adult and jack natural and hatchery origin Tucannon River spring Chinook salmon, 1996-2021..... | 22 |
| Figure 9. | Emigration timing of natural spring Chinook salmon captured during smolt trap operations (rkm 3) on the Tucannon River for the 2020-21 migration year..... | 32 |
| Figure 10. | Length frequency distribution of sampled natural spring Chinook salmon captured in the Tucannon River smolt trap, 2020/2021 season..... | 33 |
| Figure 11. | The cumulative timing to Lower Monumental Dam for natural and hatchery origin Tucannon River spring Chinook emigrants from 2021 compared to the 2006-2020 average..... | 36 |
| Figure 12. | Return per spawner (with replacement line) for the 1985-2017 brood years (2017 incomplete brood year)..... | 43 |
| Figure 13. | Tucannon River spring Chinook natural origin returns with the moving ten-year geometric mean (black line) for the 1985-2021 run years..... | 45 |
| Figure 14. | Total escapement for Tucannon River spring Chinook salmon for the 1985-2021 run years..... | 47 |
| Figure 15. | Tucannon River hatchery-origin spring Chinook survival with 95% confidence interval from release at Curl Lake to the Tucannon River instream antenna arrays (TFH = Tucannon Fish Hatchery; UTR = Upper Tucannon River; MTR = Middle Tucannon River; LTR = Lower Tucannon River) and Lower Monumental Dam (LMO) for the 2012 to 2020 migration years..... | 51 |

List of Appendices

| | | |
|-------------|--|----|
| Appendix A: | Annual Section 10 Permit #18024 Takes for 2021, and NEOR/SEWA Terms and Conditions Biological Opinion Reporting Requirements | 61 |
| Appendix B: | Spring Chinook Captured, Transported to Lyons Ferry Hatchery, or Returned to the River at the Tucannon Hatchery Trap in 2021 | 65 |
| Appendix C: | Age Composition by Brood Year for Tucannon River Spring Chinook Salmon .. | 67 |
| Appendix D: | Total Estimated Run-Size of Tucannon River Spring Chinook Salmon (1985-2021) | 69 |
| Appendix E: | Stray Hatchery-Origin Spring Chinook Salmon in the Tucannon River (1990-2021) | 71 |
| Appendix F: | Final PIT Tag Detections of Returning Tucannon River Spring Chinook, 2015 to 2021 Calendar Years | 75 |
| Appendix G: | Historical Hatchery Releases (1987-2022 Release Years) | 81 |
| Appendix H: | Numbers of Fish Species Captured by Month in the Tucannon River Smolt Trap during the 2021 Outmigration..... | 85 |
| Appendix I: | Proportionate Natural Influence (PNI) for the Tucannon Spring Chinook Population (1985-2021) | 87 |
| Appendix J: | Recoveries of Coded-Wire Tagged Salmon Released into the Tucannon River for the 1985-2017 Brood Years | 89 |

Introduction

Program Objectives

Legislation under the Water Resources Act of 1976 authorized the establishment of the Lower Snake River Compensation Plan (LSRCP) to help mitigate for the losses of salmon and steelhead runs due to construction and operation of the Snake River dams and authorized hatchery construction and production in Washington, Idaho, and Oregon as a mitigation tool (USACE 1975). In Washington, Lyons Ferry Hatchery (LFH) was constructed, and Tucannon Fish Hatchery (TFH) was modified. Under the original mitigation negotiations, local fish and wildlife agencies determined through a series of conversion rates of McNary Dam counts that 2,400 spring Chinook (2% of passage at McNary Dam) annually escaped into the Tucannon River. The agencies also estimated a 48% cumulative loss rate to juvenile downstream migrants passing through the four lower Snake River dams. As such, 1,152¹ lost adult Tucannon River origin spring Chinook needed to be compensated for above the project area, with the expectation that the other 1,248 (52%) would continue to come from natural production. An additional 4,608 were originally assumed to have been harvested in downriver fisheries or in the ocean and was an additional objective of the plan. The agencies also determined through other survival studies at the time that a smolt-to-adult survival rate (SAR) to the project area of 0.87% was a reasonable expectation for spring and summer Chinook salmon. Based on an assumed 0.87% above project area SAR and the 1,152 above project area mitigation goal it was determined that 132,000 smolts needed to be released annually. In 1984, Washington Department of Fish and Wildlife² (WDFW) began to evaluate the success of these two hatcheries in meeting the mitigation goal and identifying factors that would improve performance of the hatchery fish.

In an attempt to increase adult returns and come closer to achieving the LSRCP mitigation goal, the co-managers agreed to increase the conventional supplementation program goal to 225,000 yearling smolts annually beginning with the 2006 brood year. Size at release was increased to 38 g fish [12 fish/lb (fpp)] beginning with the 2011 brood year. In theory, both actions should have increased adult hatchery salmon returns back to the river, however, it does not appear that these actions will produce enough adult returns to reach the LSRCP adult mitigation goal (1,152).

Because of this, WDFW and the LSRCP, along with the co-managers, have initiated an additional hatchery spring Chinook program in SE Washington. A program using Carson stock spring Chinook salmon has been implemented in the Touchet River, with eyed eggs shipped to

¹ The project area escapement is 1,152. It was also assumed that four times that number (4,608 fish) would be harvested below the project area. Here “project area” is defined as above Ice Harbor Dam.

² Formerly Washington Department of Fisheries.

LFH beginning with the 2018 brood year with the first smolt releases occurring in 2020. Moving forward, adult returns from the Tucannon and Touchet programs will be used to measure contribution towards the LSRCP spring Chinook mitigation goal (1,152) for Washington.

This report summarizes work performed by the WDFW Tucannon Spring Chinook Evaluation Program from May 2021 through April 2022.

ESA Permits

The Tucannon River spring Chinook population was originally listed as “endangered” under the Endangered Species Act (ESA) on April 22, 1992 (FR 57 No. 78: 14653). The listing status was changed to “threatened” in 1995 (April 17, 1995; FR 60 No. 73: 19342). The listing was reviewed again in 1999 (FR 64 (57): 14517-14528) with the population remaining listed as “threatened” as part of the Snake River Spring/Summer Chinook Salmon evolutionary significant unit (ESU). The WDFW was originally issued a Section 10 Permit (#848 – broodstock collection and monitoring) which expired in March 1998. Permits #1126 and #1129 were issued in 1998 to allow continued take for this program, but those permits have since expired. A Hatchery and Genetic Management Plan (HGMP) was originally submitted as the application for a new Section 4 (d) Permit for this program in 2005. An updated HGMP requesting ESA Section 10 permit coverage was submitted in 2011 and was approved in 2016 (Permit #18024). This annual report summarizes all work performed by WDFW’s LSRCP Tucannon Spring Chinook Salmon Evaluation Program during 2021. Numbers of direct and indirect takes of listed Snake River spring Chinook (Tucannon River stock) for the 2021 calendar year are presented in Appendix A (Tables 1-2), along with information required for the NEOR/SEWA Biological Opinion reporting.

Facility Descriptions

Lyons Ferry Hatchery is located on the Snake River (rkm 90) at its confluence with the Palouse River and has eight deep wells that produce nearly constant 11° C water (Figure 1). It is used for adult broodstock holding and spawning, and early life incubation and rearing.

Tucannon Fish Hatchery, located at rkm 59 on the Tucannon River, has an adult collection trap on site (Figure 1). Adults returning to TFH are transported to LFH and held until spawning. Juveniles are reared at TFH through the winter on a combination of well, spring, and river water. River water is the primary source, which allows for a more natural winter temperature profile. Curl Lake Acclimation Pond was not used during 2021/2022 due to low numbers of fish and plans for a release strategy comparison study.

Tucannon River Watershed Characteristics

The Tucannon River empties into the Snake River between Little Goose and Lower Monumental Dams approximately 622 rkm from the mouth of the Columbia River (Figure 1). Stream elevation rises from 150 m at the mouth to 1,640 m at the headwaters (Bugert et al. 1990). Total watershed area is approximately 1,295 km². Local habitat problems related to logging, road building, recreation, and agriculture/livestock grazing have limited the production potential of spring Chinook in the Tucannon River. Land use in the Tucannon watershed is approximately 36% grazed rangeland, 33% dry cropland, 23% forest, 6% WDFW, and 2% other use (Tucannon Subbasin Summary 2001). Five unique strata have been distinguished by predominant land use, habitat, and landmarks (Figure 1; Table 1) and are referenced throughout this report.




Figure 1. Location of the Tucannon River, and Lyons Ferry and Tucannon Hatcheries within the Snake River basin.

Table 1. Description of five strata within the Tucannon River.

| Strata | Land Ownership/Usage | Spring Chinook Habitat^a | River Kilometer^b |
|---------------|--------------------------------|---|------------------------------------|
| Lower | Private/Agriculture & Ranching | Not-Usable (temperature limited) | 0.0-20.1 |
| Marengo | Private/Agriculture & Ranching | Marginal (temperature limited) | 20.1-39.9 |
| Hartsock | Private/Agriculture & Ranching | Fair to Good | 39.9-55.5 |
| HMA | State & Federal/Recreational | Good to Excellent | 55.5-74.5 |
| Wilderness | Federal/Recreational | Excellent | 74.5-86.3 |

^a Strata were based on water temperature, habitat, and landowner use.

^b Rkm descriptions: 0.0–mouth at the Snake River; 20.1-Territorial Rd.; 39.9–Marengo Br.; 55.5-HMA Boundary Fence; 74.5-Panjab Br.; 86.3-Rucherts Camp.

Adult Salmon Evaluation

Broodstock Trapping

The allowed collection goal for broodstock is 170 adult salmon, depending upon size and fecundity, collected from throughout the duration of the run to meet the smolt production/release goal of 225,000. The proportion of natural origin fish incorporated into the broodstock is based on the estimated run size and the Tucannon Spring Chinook Salmon HGMP sliding scale. Additional jack salmon may be collected up to their proportion of the run with an upper limit of 10% used in broodstock, if needed. Returning Tucannon stock hatchery salmon were identified by coded-wire tag (CWT) in the snout, with no adipose fin clips. Adipose clipped fish captured at the trap are killed outright as strays.

The TFH adult trap began operation in February (for steelhead) with the first spring Chinook captured on 24 May (Appendix B). State and Tribal Fisheries Managers decided to collect all Tucannon River returns to be used for broodstock due to the expected low run size. This was also done to circumvent potential high in-river pre-spawn mortality that has been observed in preceding years.

The trap was operated through 30 September. A total of 140 fish entered the trap (76 natural adults, 7 natural jacks, 35 hatchery adults, and 22 hatchery jacks) and 115 fish (76 natural adults, 7 natural jacks, 22 hatchery adults, 10 hatchery jacks) were collected for broodstock (Table 2, Appendix B). Twenty-five adipose clipped strays were killed outright at the adult trap (Table 2, Appendix B). Adults collected for broodstock were injected with erythromycin at 20 mg/kg. Antibiotic injections for broodstock were repeated 30 days prior to spawning. Broodstock received formalin drip treatments during holding at 167 ppm every other day to control fungus.

Table 2. Numbers of spring Chinook salmon captured at the TFH trap, trap mortalities, strays or jacks killed outright, fish collected for broodstock, and passed upstream or held for adult outplanting for natural spawning from 1986-2021.

| Year | Captured at Trap | | Trap Mortalities | | Killed Outright ^a | Broodstock Collected | | Passed Upstream | | Held for Outplanting | |
|-------------------|------------------|----------|------------------|----------|------------------------------|----------------------|----------|-----------------|----------------|----------------------|------------------|
| | Natural | Hatchery | Natural | Hatchery | Hatchery | Natural | Hatchery | Natural | Hatchery | Natural | Hatchery |
| 1986 | 247 | 0 | 0 | 0 | 0 | 116 | 0 | 131 | 0 | 0 | 0 |
| 1987 | 209 | 0 | 0 | 0 | 0 | 101 | 0 | 108 | 0 | 0 | 0 |
| 1988 | 267 | 9 | 0 | 0 | 0 | 116 | 9 | 151 | 0 | 0 | 0 |
| 1989 | 156 | 102 | 0 | 0 | 0 | 67 | 102 | 89 | 0 | 0 | 0 |
| 1990 | 252 | 216 | 0 | 1 | 0 | 60 | 75 | 192 | 140 | 0 | 0 |
| 1991 | 109 | 202 | 0 | 0 | 0 | 41 | 89 | 68 | 113 | 0 | 0 |
| 1992 | 242 | 305 | 8 | 3 | 0 | 47 | 50 | 187 | 252 | 0 | 0 |
| 1993 | 191 | 257 | 0 | 0 | 0 | 50 | 47 | 141 | 210 | 0 | 0 |
| 1994 | 36 | 34 | 0 | 0 | 0 | 36 | 34 | 0 | 0 | 0 | 0 |
| 1995 | 10 | 33 | 0 | 0 | 0 | 10 | 33 | 0 | 0 | 0 | 0 |
| 1996 | 76 | 59 | 1 | 4 | 0 | 35 | 45 | 40 | 10 | 0 | 0 |
| 1997 | 99 | 160 | 0 | 0 | 0 | 43 | 54 | 56 | 106 | 0 | 0 |
| 1998 ^b | 50 | 43 | 0 | 0 | 0 | 48 | 41 | 1 | 1 | 0 | 0 |
| 1999 ^c | 4 | 139 | 0 | 1 | 0 | 4 | 135 | 0 | 0 | 0 | 0 |
| 2000 | 25 | 180 | 0 | 0 | 17 | 12 | 69 | 13 | 94 | 0 | 0 |
| 2001 | 405 | 276 | 0 | 0 | 0 | 52 | 54 | 353 | 222 | 0 | 0 |
| 2002 | 168 | 610 | 0 | 0 | 0 | 42 | 65 | 126 | 545 | 0 | 0 |
| 2003 | 84 | 151 | 0 | 0 | 0 | 42 | 35 | 42 | 116 | 0 | 0 |
| 2004 | 311 | 155 | 0 | 0 | 0 | 51 | 41 | 260 | 114 | 0 | 0 |
| 2005 | 131 | 114 | 0 | 0 | 3 | 49 | 51 | 82 | 60 | 0 | 0 |
| 2006 | 61 | 78 | 0 | 1 | 2 | 36 | 53 | 25 | 22 | 0 | 0 |
| 2007 | 112 | 112 | 0 | 0 | 6 | 54 | 34 | 58 | 72 | 0 | 0 |
| 2008 | 114 | 386 | 0 | 0 | 1 | 42 | 92 | 72 | 293 | 0 | 0 |
| 2009 | 390 | 835 | 0 | 0 | 7 | 89 | 88 | 301 | 740 | 0 | 0 |
| 2010 | 774 | 796 | 0 | 0 | 9 | 86 | 87 | 688 | 700 | 0 | 0 |
| 2011 | 400 | 383 | 0 | 0 | 6 | 89 | 77 | 311 | 300 | 0 | 0 |
| 2012 | 240 | 301 | 0 | 0 | 6 | 93 | 77 | 147 | 218 | 0 | 0 |
| 2013 | 271 | 268 | 0 | 0 | 2 | 98 | 60 | 173 | 206 | 0 | 0 |
| 2014 ^d | 343 | 215 | 0 | 0 | 0 | 86 | 41 | 257 | 174 | 0 | 0 |
| 2015 | 285 | 594 | 0 | 0 | 32 | 101 | 30 | 126 | 348 | 58 | 184 |
| 2016 | 127 | 468 | 0 | 0 | 114 | 55 | 71 | 6 | 19 | 66 | 264 |
| 2017 | 26 | 237 | 0 | 0 | 15 | 18 | 93 | 0 | 0 | 8 ^e | 129 ^e |
| 2018 | 73 | 358 | 0 | 0 | 38 | 37 | 123 | 15 | 3 | 21 | 194 |
| 2019 | 39 | 144 | 0 | 0 | 1 | 36 | 113 | 1 | 1 | 2 | 29 |
| 2020 | 38 | 18 | 0 | 0 | 2 | 38 | 15 | 0 | 1 ^f | 0 | 0 |
| 2021 | 83 | 57 | 0 | 0 | 25 | 83 | 32 | 0 | 0 | 0 | 0 |

^a Fish identified as strays at the adult trap are killed outright. Some hatchery jacks were killed outright in 2016.

^b Two males (one natural, one hatchery) captured were transported back downstream to spawn in the river.

^c Three hatchery males that were captured were transported back downstream to spawn in the river.

^d Ninety-four natural origin fish were collected for broodstock; however, eight natural origin females were returned to the river for natural spawning leaving a total of 86 natural origin fish collected for broodstock.

^e None of the fish held for adult outplanting in 2017 were outplanted. All of the fish held for adult outplanting were < 61 cm in fork length (jack size) and were either used to supplement broodstock (natural jacks) or were killed outright.

^f This fish was mistakenly passed downstream.

Broodstock Mortality

Three (2.6%) of the 115 salmon collected for broodstock died prior to spawning in 2021 (Table 3). Pre-spawn mortality of both natural and hatchery origin fish averaged over 10% from 2017-2020. Higher losses in 2017 and 2018 may have been the result of halting prophylactic antibiotic injections (2017) and utilization of a different antibiotic (2018). High pre-spawn mortality was experienced when fish were held at TFH (1986-1991 and 2019), likely due to higher water temperatures (Table 3).

Table 3. Numbers of pre-spawning mortalities and percent of fish collected for broodstock at TFH and held at TFH (1985-1991 and 2019) or LFH (1992-2018 and 2020-2021).

| Year | Natural | | | % of collected | Hatchery | | | % of collected |
|------|---------|--------|------|----------------|----------|--------|------|----------------|
| | Male | Female | Jack | | Male | Female | Jack | |
| 1985 | 3 | 10 | 0 | 59.1 | — | — | — | — |
| 1986 | 15 | 10 | 0 | 21.6 | — | — | — | — |
| 1987 | 10 | 8 | 0 | 17.8 | — | — | — | — |
| 1988 | 7 | 22 | 0 | 25.0 | — | — | 9 | 100.0 |
| 1989 | 8 | 3 | 1 | 17.9 | 5 | 8 | 22 | 34.3 |
| 1990 | 12 | 6 | 0 | 30.0 | 14 | 22 | 3 | 52.0 |
| 1991 | 0 | 0 | 1 | 2.4 | 8 | 17 | 32 | 64.0 |
| 1992 | 0 | 4 | 0 | 8.2 | 2 | 0 | 0 | 4.0 |
| 1993 | 1 | 2 | 0 | 6.0 | 2 | 1 | 0 | 6.4 |
| 1994 | 1 | 0 | 0 | 2.8 | 0 | 0 | 0 | 0.0 |
| 1995 | 1 | 0 | 0 | 10.0 | 0 | 0 | 3 | 9.1 |
| 1996 | 0 | 2 | 0 | 5.7 | 2 | 1 | 0 | 6.7 |
| 1997 | 0 | 4 | 0 | 9.3 | 2 | 2 | 0 | 7.4 |
| 1998 | 1 | 2 | 0 | 6.3 | 0 | 0 | 0 | 0.0 |
| 1999 | 0 | 0 | 0 | 0.0 | 3 | 1 | 1 | 3.8 |
| 2000 | 0 | 0 | 0 | 0.0 | 1 | 2 | 0 | 3.7 |
| 2001 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0.0 |
| 2002 | 0 | 0 | 0 | 0.0 | 1 | 1 | 0 | 3.1 |
| 2003 | 0 | 1 | 0 | 2.4 | 0 | 0 | 1 | 2.9 |
| 2004 | 0 | 3 | 0 | 5.9 | 0 | 0 | 1 | 2.4 |
| 2005 | 2 | 0 | 0 | 4.1 | 1 | 2 | 0 | 5.9 |
| 2006 | 0 | 0 | 0 | 0.0 | 1 | 0 | 0 | 1.9 |
| 2007 | 0 | 2 | 1 | 5.6 | 0 | 2 | 0 | 5.9 |
| 2008 | 1 | 1 | 0 | 4.8 | 0 | 0 | 1 | 1.1 |
| 2009 | 0 | 0 | 0 | 0.0 | 0 | 2 | 0 | 2.3 |
| 2010 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0.0 |
| 2011 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0.0 |
| 2012 | 0 | 0 | 0 | 0.0 | 1 | 2 | 0 | 3.9 |
| 2013 | 2 | 3 | 0 | 5.1 | 0 | 2 | 0 | 3.3 |
| 2014 | 0 | 1 | 0 | 1.2 | 0 | 0 | 0 | 0.0 |
| 2015 | 0 | 1 | 0 | 1.0 | 0 | 1 | 0 | 3.3 |
| 2016 | 0 | 1 | 0 | 1.8 | 2 | 0 | 0 | 2.8 |
| 2017 | 0 | 2 | 0 | 12.5 | 4 | 8 | 0 | 12.6 |
| 2018 | 2 | 2 | 0 | 10.8 | 12 | 4 | 0 | 13.0 |
| 2019 | 3 | 9 | 0 | 33.3 | 14 | 38 | 0 | 46.0 |
| 2020 | 2 | 3 | 0 | 13.2 | 0 | 0 | 0 | 0.0 |
| 2021 | 0 | 1 | 0 | 1.2 | 0 | 1 | 1 | 6.3 |

Broodstock Spawning

Spawning at LFH was conducted once a week from 31 August to 28 September, with the peak eggtake occurring on 14 September. During the spawning process, the eggs of two females were split in half and fertilized by two males following a 2 x 2 factorial spawning matrix approach. Factorial mating can have substantial advantages in increasing the effective number of breeders (Busack and Knudsen 2007). The priority order of crosses was Natural x Hatchery, Natural x Natural, and Hatchery x Hatchery, depending upon availability and origin of ripe fish on each spawning date.

A total of 166,237 eggs were collected from 47 spawned females (Table 4). Eggs were initially disinfected, and water hardened for one hour in an iodophor (buffered iodine) solution (100 ppm). The eggs were incubated in vertical tray incubators. Fungus on the incubating eggs was controlled with formalin applied every-other day at 1,667 ppm for 15 minutes. Mortality to eye-up was 11.9%, which left 146,354 live eggs. An additional 0.4% (647) loss of eggs and sac-fry left 145,707 fish for production.

Table 4. Number of fish spawned or killed outright (K.O.), estimated egg collection, and egg mortality of natural and hatchery origin Tucannon River spring Chinook salmon at LFH in 2021. (Numbers in parentheses were live spawned).

| Spawn Date | Natural Origin | | | | | | Eggs Taken |
|---------------|-----------------|----------------|---------|----------------|---------|----------------|----------------|
| | Males | | Jacks | | Females | | |
| | Spawned | K.O. | Spawned | K.O. | Spawned | K.O. | |
| 8/31 | 1 (3) | | | | | 2 | 6,565 |
| 9/07 | (12) | | | 1 | | 10 | 38,476 |
| 9/14 | 2 (16) | | | | | 15 | 56,152 |
| 9/21 | 6 (9) | | | | | 11 | 39,461 |
| 9/28 | 28 | | | 6 | | | |
| Totals | 37 | | | 7 | | 38 | 140,654 |
| Egg Mortality | | | | | | | 17,937 |
| Spawn Date | Hatchery Origin | | | | | | Eggs Taken |
| | Males | | Jacks | | Females | | |
| | Spawned | K.O. | Spawned | K.O. | Spawned | K.O. | |
| 8/31 | | 1 ^a | | 1 ^a | | 5 ^a | |
| 9/07 | 2 | 1 ^a | | 1 ^a | | 3 | 8,187 |
| 9/14 | 1 (1) | | | | | 3 | 9,431 |
| 9/21 | 1 | | | 2 ^a | | 2 | 5,740 |
| 9/28 | 1 | | | 5 ^b | | 1 | 2,225 |
| Totals | 5 | 2 | | 9 | | 9 | 25,583 |
| Egg Mortality | | | | | | | 1,946 |

^a Hatchery strays that were not used for spawning.

^b Two of the five were stray hatchery jacks. The other three were Tucannon hatchery jacks not used for spawning.

Broodstock BKD Screening and Virology Testing

Broodstock females were screened for Bacterial Kidney Disease (BKD), caused by the bacterium *Renibacterium salmoninarum*, using Enzyme Linked Immunosorbent Assay (ELISA). None of the spawned females had high values in 2021 (Table 5). High values from 2017 were believed to be directly related to the decision to suspend antibiotic injections during that year (Figure 2). During 2020, the Fish Health Specialist switched back to erythromycin injections from tulathromycin (Draxxin³) injections since it was considered to be more effective against BKD. Spawned females were also examined for viruses and sampling showed no evidence of virus in the samples tested.

Table 5. Enzyme Linked Immunosorbent Assay (ELISA) values for hatchery spawned Tucannon River spring Chinook females, 2021.

| ELISA Value | ELISA O.D. | Number of Females | Percent (%) |
|--------------|---------------|-------------------|--------------|
| Below Low | < 0.099 | 47 | 100.0 |
| Low | 0.099 – 0.198 | 0 | 0.0 |
| Moderate | 0.199 – 0.450 | 0 | 0.0 |
| High | > 0.450 | 0 | 0.0 |
| Total | | 47 | 100.0 |




Figure 2. Historical Below Low and Low, and Moderate and High ELISA values for Tucannon River spring Chinook salmon female broodstock for the 1998 to 2021 return years.

³ The use of trade names does not imply endorsement by the Washington Department of Fish and Wildlife.

Natural Spawning

Pre-spawn mortality surveys were conducted from 16 June until 9 July when access to the river was closed due to the Green Ridge forest fire. The pre-spawn mortality surveys covered from Bridge 14 (rkm 51.5) to the Tucannon Fish Hatchery Intake (rkm 59.2). No salmon carcasses were observed during pre-spawn mortality surveys.

Weekly spawning ground surveys began on 26 August below the Hatchery Intake and were completed by 1 October. On 8 September, permission was granted to survey upstream of the Hatchery Intake, but only up to Panjab Bridge (rkm 74.5). The river upstream of Panjab Bridge was not surveyed during 2021 (Table 6). A total of 35 redds were counted during surveys, with four redds (11%) observed above the adult trap (Table 6), even though fish were not passed upstream during 2021. Fifteen carcasses were recovered during surveys (5 natural origin, 10 hatchery origin) and all of the recovered hatchery origin carcasses were strays. A cumulative 179 river kilometers were walked from Panjab Bridge down to the mouth during 2021.

Table 6. Numbers and general locations of salmon redds and carcasses (includes pre-spawn mortalities) recovered on the Tucannon River spawning grounds, 2021 (the Tucannon Hatchery adult trap is located at rkm 59).

| Stratum | Rkm ^a | Number of redds | Carcasses Recovered | |
|---------------------------------------|------------------|-----------------|---------------------|-----------|
| | | | Natural | Hatchery |
| Wilderness | 84-86 | --- | --- | --- |
| | 78-84 | --- | --- | --- |
| | 75-78 | --- | --- | --- |
| HMA | 73-75 | 0 | 0 | 0 |
| | 68-73 | 0 | 0 | 0 |
| | 66-68 | 0 | 0 | 0 |
| | 62-66 | 2 | 0 | 0 |
| | 59-62 | 2 | 1 | 1 |
| -----Tucannon Fish Hatchery Trap----- | | | | |
| Hartsock | 56-59 | 13 | 1 | 5 |
| | 52-56 | 2 | 1 | 1 |
| | 47-52 | 4 | 0 | 2 |
| | 43-47 | 3 | 1 | 1 |
| Marengo | 40-43 | 4 | 0 | 0 |
| | 34-40 | 1 | 0 | 0 |
| | 28-34 | 0 | 0 | 0 |
| Below Marengo | 0-28 | 4 | 1 | 0 |
| Totals | 0-86 | 35 | 5 | 10 |

^a Rkm descriptions: 86-Rucherts Camp; 84-Sheep Cr.; 78-Lady Bug Flat CG; 75-Panjab Br.; 73-Cow Camp Bridge; 68-Camp Wooten Br.; 66-Curl Lake; 62-Beaver/Watson Lakes Br.; 59-Tucannon Hatchery Intake/Adult Trap; 56-Cummings Creek Br.; 52-Br. 14; 47-Br. 12; 43-Br. 10; 40-Marengo Br.; 34-King Grade Br.; 28-Enrich Br. (Brines Rd.).

Historical Trends in Natural Spawning

Examining historical traits in natural spawning (1985-present), redd density has varied greatly with run size over the years with a high of 8.3 redds/km in 2010 to a low of 0.1 redds/km during 1995 (Figure 3; Table 7). Since the program's inception in 1985, the proportion of the total number of redds occurring below the adult trap has increased (Figure 4; Table 7). This is likely the result of a combination of fish that were unwilling to enter the TFH fish ladder/trap and an emphasis on broodstock collection that was intended to reduce the risk of extinction.




Figure 3. Spring Chinook redd density (redds/km) in the Tucannon River, 1986-2021.




Figure 4. The proportion of redds above Marengo that were either above the adult trap/weir or below the adult trap/weir with trend lines, 1985-1993 and 2000-2014. (Note: The 1994-1999 data was removed from the graph due to management actions at the trap, and after 2014 since adult outplants began in 2015 and due to the collection of all fish for broodstock in recent years.).

Table 7. Number of spring Chinook salmon redds and redds/km (in parentheses) by stratum and year, and the number and percent of redds above and below the TFH adult trap in the Tucannon River, 1985-2021.

| Year | Strata ^a | | | | Total Redds ^b | TFH Adult Trap ^b | | | |
|-------------------|---------------------|------------|-----------|----------|-----------------------------|-----------------------------|------|-------|-------|
| | Wilderness | HMA | Hartsock | Marengo | | Above | % | Below | % |
| 1985 ^c | 101 (9.2) | 165 (8.7) | 50 (3.1) | – | 316 | – | – | – | – |
| 1986 | 53 (4.5) | 117 (6.2) | 29 (1.9) | 0 (0.0) | 200 | 163 | 81.5 | 37 | 18.5 |
| 1987 | 15 (1.3) | 140 (7.4) | 30 (1.9) | – | 185 | 149 | 80.5 | 36 | 19.5 |
| 1988 | 18 (1.5) | 79 (4.2) | 20 (1.3) | – | 117 | 90 | 76.9 | 27 | 23.1 |
| 1989 | 29 (2.5) | 54 (2.8) | 23 (1.5) | – | 106 | 74 | 69.8 | 32 | 30.2 |
| 1990 | 20 (1.7) | 94 (4.9) | 64 (4.1) | 2 (0.3) | 180 | 96 | 53.3 | 84 | 46.7 |
| 1991 | 3 (0.3) | 67 (2.9) | 18 (1.1) | 2 (0.3) | 90 | 40 | 44.4 | 50 | 55.6 |
| 1992 | 17 (1.4) | 151 (7.9) | 31 (2.0) | 1 (0.2) | 200 | 130 | 65.0 | 70 | 35.0 |
| 1993 | 34 (3.4) | 123 (6.5) | 34 (2.2) | 1 (0.2) | 192 | 131 | 68.2 | 61 | 31.8 |
| 1994 | 1 (0.1) | 10 (0.5) | 28 (1.8) | 5 (0.9) | 44 | 2 | 4.5 | 42 | 95.5 |
| 1995 | 0 (0.0) | 2 (0.1) | 3 (0.2) | 0 (0.0) | 5 | 0 | 0.0 | 5 | 100.0 |
| 1996 | 1 (0.1) | 33 (1.7) | 34 (2.2) | 1 (0.2) | 69 | 11 | 16.2 | 58 | 83.8 |
| 1997 | 2 (0.2) | 43 (2.3) | 27 (1.7) | 1 (0.2) | 73 | 30 | 41.1 | 43 | 58.9 |
| 1998 | 0 (0.0) | 3 (0.2) | 20 (1.3) | 3 (0.5) | 26 | 3 | 11.5 | 23 | 88.5 |
| 1999 | 1 (0.1) | 34 (1.8) | 6 (0.4) | 0 (0.0) | 41 | 3 | 7.3 | 38 | 92.7 |
| 2000 | 4 (0.4) | 68 (3.6) | 20 (1.3) | 0 (0.0) | 92 | 45 | 48.9 | 47 | 51.1 |
| 2001 | 22 (2.0) | 194 (10.2) | 80 (5.0) | 1 (0.1) | 297 | 166 | 55.9 | 131 | 44.1 |
| 2002 | 29 (2.6) | 214 (11.3) | 45 (2.8) | 11 (0.9) | 299 | 200 | 66.9 | 99 | 33.1 |
| 2003 | 3 (0.3) | 89 (4.7) | 26 (1.6) | 0 (0.0) | 118 | 61 | 51.7 | 57 | 48.3 |
| 2004 | 24 (2.2) | 119 (6.3) | 17 (1.1) | 0 (0.0) | 160 | 112 | 70.0 | 48 | 30.0 |
| 2005 | 4 (0.4) | 71 (3.7) | 27 (1.7) | 5 (0.4) | 107 | 46 | 43.0 | 61 | 57.0 |
| 2006 | 2 (0.2) | 81 (4.3) | 17 (1.1) | 1 (0.1) | 109 | 58 | 53.2 | 51 | 46.8 |
| 2007 | 2 (0.2) | 63 (3.3) | 16 (1.0) | 0 (0.0) | 81 | 32 | 39.5 | 49 | 60.5 |
| 2008 | 30 (2.7) | 146 (7.7) | 22 (1.4) | 1 (0.1) | 199 | 141 | 70.9 | 58 | 29.1 |
| 2009 | 67 (6.1) | 329 (17.3) | 52 (3.3) | 3 (0.3) | 451 | 292 | 64.7 | 159 | 35.3 |
| 2010 | 83 (7.5) | 289 (15.2) | 106 (6.6) | 3 (0.3) | 481 | 297 | 61.7 | 184 | 38.3 |
| 2011 | 35 (3.2) | 196 (10.3) | 53 (3.3) | 6 (0.5) | 297 | 165 | 55.6 | 132 | 44.4 |
| 2012 | 11 (1.0) | 132 (6.9) | 23 (1.4) | 0 (0.0) | 169 | 84 | 49.7 | 85 | 50.3 |
| 2013 | 3 (0.3) | 42 (2.2) | 15 (0.9) | 0 (0.0) | 64 | 25 | 39.1 | 39 | 60.9 |
| 2014 | 26 (2.4) | 70 (3.7) | 25 (1.6) | 1 (0.1) | 124 | 83 | 66.9 | 41 | 33.1 |
| 2015 | 56 (5.1) | 91 (4.8) | 33 (2.1) | 4 (0.3) | 191 | 120 | 62.8 | 71 | 37.2 |
| 2016 | 37 (3.4) | 79 (4.2) | 31 (1.9) | 3 (0.3) | 154 | 83 | 53.9 | 71 | 46.1 |
| 2017 | 8 (0.7) | 47 (2.5) | 15 (0.9) | 0 (0.0) | 70 | 29 | 41.4 | 41 | 58.6 |
| 2018 | 31 (2.8) | 64 (3.4) | 13 (0.8) | 0 (0.0) | 109 | 77 | 70.6 | 32 | 29.4 |
| 2019 | 0 (0.0) | 9 (0.5) | 1 (0.1) | 0 (0.0) | 11 | 3 | 27.3 | 8 | 72.7 |
| 2020 | 0 (0.0) | 10 (0.5) | 2 (0.1) | 2 (0.2) | 14 | 2 | 14.3 | 12 | 85.7 |
| 2021 | – | 17 (0.9) | 13 (0.8) | 1 (0.1) | 35 | 4 | 11.4 | 31 | 88.6 |

Note: – indicates the river was not surveyed in that section during that year.

^a Excludes redds found below the Marengo stratum.

^b Includes all redds counted during redd surveys.

^c The 1985 redd counts were revised to account for all redds during the spawning season (WDFW 2017).

Stream Nutrient Enrichment

The majority of hatchery broodstock carcasses have traditionally been buried on-site at LFH after spawning. However, declines in salmonid abundance during the last century have resulted in decreased deposition of marine-derived nutrients and pose a significant restraint in the recovery of threatened and endangered Pacific salmon (Nehlsen et al. 1991; Scheuerell et al. 2005). The importance of marine derived nutrients to salmon recovery efforts has prompted local volunteer groups and state, federal, and tribal agencies to add supplemental nutrients into freshwater habitats, especially in salmon depleted habitats (Kohler et al. 2012). Stream nutrient enrichment efforts in the Tucannon River had been sporadic during the history of the hatchery program. However, except for 2019 due to a broken freezer, stream nutrient enrichment has been occurring on an annual basis since 2016.

A total of 1,221 fall Chinook salmon carcasses were available for stream nutrient enrichment in the Tucannon River as the result of hatchery spawning at LFH during 2021. Virology testing did not show presence of viral pathogens, so the WDFW's "Protocols for the Nutrient Enrichment of the Tucannon River to Increase Production of Salmon and Steelhead" were followed and the fall Chinook carcasses, due to their relative abundance, were used as a surrogate for spring Chinook carcasses. Department and Confederated Tribes of the Umatilla Indian Reservation (CTUIR) employees and volunteers from the Tri-State Steelheader's Regional Fisheries Enhancement Group distributed the carcasses between the Tucannon Fish Hatchery Bridge (rkm 58.2) and Highway 12 Bridge (rkm 22.0) on 8 and 10 January 2022 (Table 8). Carcasses were distributed to benefit both spring Chinook and steelhead based on redd locations, the expected downstream movement of juveniles, and stream drift of aquatic macroinvertebrates.

Standard tracking of salmon/steelhead production and growth will occur and characteristics of juveniles (numbers, size, weight, condition factor, etc.) captured at the Tucannon River smolt trap will be monitored to examine the relationship between stream nutrient enrichment and juvenile production.

Table 8. Summary of Chinook salmon carcass distribution on the Tucannon River on 8 and 10 January 2022.

| Location Name | River kilometer | Number of Carcasses |
|--------------------------|------------------------|----------------------------|
| Hatchery Bridge | 58.2 | 150 |
| Cummings Creek Bridge | 55.9 | 150 |
| Bridge 14 | 51.5 | 130 |
| Bridge 13 | 48.9 | 100 |
| Bridge 12 | 47.1 | 100 |
| Bridge 11 | 44.0 | 100 |
| Bridge 10 | 43.3 | 76 |
| Marengo Bridge | 39.9 | 115 |
| King Grade Bridge | 34.1 | 100 |
| Brines Rd./Enrich Bridge | 28.0 | 100 |
| Near Highway 12 Bridge | 22.0 | 100 |
| Totals | | 1,221 |

Genetic Sampling

During 2021, we collected 127 DNA samples (tissue samples) from hatchery broodstock and carcasses collected from the spawning grounds (87 natural origin and 40 hatchery origin). These samples were sent to the WDFW genetics lab in Olympia, Washington for storage. Genetic samples from the broodstock (spawned fish only) were also collected and sent to the Idaho Department of Fish and Game for parentage-based tagging analysis for Snake River Basin spring/summer Chinook populations. Genotypes, allele frequencies, and tissue samples from some of the previous sampling years are available from WDFW's Genetics Laboratory.

Age Composition, Length Comparisons, and Fecundity

We determine the age composition of each year's returning adults from scale samples of natural origin fish, and both scales and CWTs from hatchery-origin fish collected for broodstock and from carcasses collected during spawning ground surveys. This enables us to compare ages of natural and hatchery-reared fish, and to examine trends and variability in age structure. The recovery of jack salmon from the river is low and jacks are typically not collected for broodstock, so their representation is biased low compared to observations from the adult trap.

Overall, hatchery origin fish return at a younger age than natural origin fish and have fewer age-5 fish in the population compared to the historical age composition (Figure 5). This difference is likely due to larger size-at-release of hatchery origin fish that can lead to higher proportions of early maturing fish (hatchery origin smolts are generally 40-50 mm greater in length than natural smolts). The mean age (weighted) of males and females for both hatchery and natural

origin fish (Figure 6) has fluctuated over time, with hatchery males and females generally having a lower mean age than natural origin fish due to their larger size at release. Females are typically closer in mean age than males (Figure 6). The age composition by brood year for natural and hatchery origin fish is found in Appendix C.




Figure 5. Historical (1985-2020), and 2021 age composition (run year) for spring Chinook in the Tucannon River.




Figure 6. Weighted mean age of natural and hatchery origin males (NM, HM) and natural and hatchery origin females (NF, HF) for the 1985 to 2016 brood years for spring Chinook in the Tucannon River.

Another metric monitored on returning adult natural and hatchery origin fish is size at age, measured as the mean post-orbital to hypural-plate (POH) length. We examined size at age for age-4 adult returns using multiple comparison analysis from 1985-2021 and found that age-4 male and female natural origin spring Chinook salmon had significantly larger POH length ($P < 0.001$) than age-4 male and female hatchery-origin spring Chinook salmon (Figure 7).




Figure 7. Mean post-orbital to hypural-plate (POH) length comparisons between age-4 natural and hatchery-origin males (NM and HM) and natural and hatchery-origin females (NF and HF) with 95% confidence intervals for the years 1985-2021.

To estimate fecundities (number of eggs/female) from the 2021 return year, dead eggs were counted for each female and a subsample of 100 live eyed eggs was weighed. The total mass of live eggs was also weighed and divided by the average weight per egg to yield total number of live eggs. This estimate was decreased by 4% to compensate for adherence of water on the eggs (WDFW Snake River Lab, unpublished data). Fecundities of natural and hatchery origin fish from the Tucannon River program have been documented since 1990 (Table 9). We performed an analysis of variance to determine if there were differences in mean fecundities of hatchery and natural origin fish. The significance level for all statistical tests was 0.05. Natural origin females were significantly more fecund than hatchery origin fish for both age-4 ($P < 0.001$) and age-5 fish ($P < 0.001$). These data correspond with data collected by Gallinat and Chang (2013) that examined the effects of hatchery rearing on selected phenotypic traits of female Tucannon River spring Chinook salmon. They found that hatchery origin females had significantly lower fecundity than natural origin fish after correcting for body size.

Table 9. Average number of eggs/female (n, SD) by age group of Tucannon River natural and hatchery origin broodstock, 1990-2021 (partial spawned females are excluded).

| Year | Age 4 | | | | Age 5 | | | |
|-------------|--------------|--------------|--------------|-------------|--------------|---------------|--------------|--------------|
| | Natural | | Hatchery | | Natural | | Hatchery | |
| 1990 | 3,691 | (13, 577.3) | 2,795 | (18, 708.0) | 4,383 | (8, 772.4) | No Fish | |
| 1991 | 3,140 | (5, 363.3) | 2,649 | (9, 600.8) | 4,252 | (11, 776.0) | 3,052 | (1, 000.0) |
| 1992 | 3,736 | (16, 588.3) | 3,286 | (25, 645.1) | 4,800 | (2, 992.8) | 3,545 | (1, 000.0) |
| 1993 | 3,267 | (4, 457.9) | 3,456 | (5, 615.4) | 4,470 | (2, 831.6) | 4,129 | (1, 000.0) |
| 1994 | 3,688 | (13, 733.9) | 3,280 | (11, 630.3) | 4,848 | (8, 945.8) | 3,352 | (10, 705.9) |
| 1995 | No Fish | | 3,584 | (14, 766.4) | 5,284 | (6, 1,361.2) | 3,889 | (1, 000.0) |
| 1996 | 3,510 | (17, 534.3) | 2,853 | (18, 502.3) | 3,617 | (1, 000.0) | No Fish | |
| 1997 | 3,487 | (15, 443.1) | 3,290 | (24, 923.2) | 4,326 | (3, 290.8) | No Fish | |
| 1998 | 4,204 | (1, 000.0) | 2,779 | (7, 405.5) | 4,017 | (28, 680.5) | 3,333 | (6, 585.2) |
| 1999 | No Fish | | 3,121 | (34, 445.4) | No Fish | | 3,850 | (1, 000.0) |
| 2000 | 4,144 | (2, 1,571.2) | 3,320 | (34, 553.6) | 3,618 | (1, 000.0) | 4,208 | (1, 000.0) |
| 2001 | 3,612 | (27, 518.1) | 3,225 | (24, 705.4) | No Fish | | 3,585 | (2, 1,191.5) |
| 2002 | 3,584 | (14, 740.7) | 3,368 | (24, 563.7) | 4,774 | (7, 429.1) | No Fish | |
| 2003 | 3,342 | (10, 778.0) | 2,723 | (2, 151.3) | 4,428 | (7, 966.3) | 3,984 | (17, 795.9) |
| 2004 | 3,376 | (26, 700.5) | 2,628 | (17, 397.8) | 5,191 | (1, 000.0) | 2,151 | (1, 000.0) |
| 2005 | 3,399 | (18, 545.9) | 2,903 | (22, 654.2) | 4,734 | (7, 1,025.0) | No Fish | |
| 2006 | 2,857 | (17, 559.1) | 2,590 | (26, 589.8) | 3,397 | (1, 000.0) | 4,319 | (1, 000.0) |
| 2007 | 3,450 | (14, 721.1) | 2,679 | (6, 422.7) | 4,310 | (12, 1,158.0) | 3,440 | (2, 997.7) |
| 2008 | 3,698 | (16, 618.9) | 3,018 | (40, 501.3) | 4,285 | (1, 000.0) | 4,430 | (1, 000.0) |
| 2009 | 3,469 | (34, 628.9) | 3,267 | (52, 641.3) | 4,601 | (6, 753.6) | No Fish | |
| 2010 | 3,579 | (38, 594.8) | 3,195 | (44, 640.9) | No Fish | | No Fish | |
| 2011 | 3,513 | (18, 613.0) | 3,061 | (30, 615.1) | 4,709 | (27, 755.2) | 3,954 | (11, 731.3) |
| 2012 | 2,998 | (40, 618.1) | 2,539 | (45, 462.5) | 4,371 | (5, 478.0) | 3,105 | (2, 356.4) |
| 2013 | 3,479 | (34, 574.8) | 3,145 | (28, 592.9) | 4,702 | (12, 931.5) | 3,746 | (2, 185.3) |
| 2014 | 3,622 | (34, 501.3) | 3,280 | (26, 545.6) | 4,575 | (3, 807.3) | 3,558 | (1, 000.0) |
| 2015 | 3,683 | (47, 629.5) | 3,468 | (20, 671.8) | 4,755 | (8, 818.0) | No Fish | |
| 2016 | 3,456 | (19, 676.1) | 3,133 | (36, 652.7) | 4,096 | (12, 891.2) | 3,514 | (5, 508.6) |
| 2017 | 3,393 | (8, 453.9) | 3,034 | (50, 586.0) | No Fish | | No Fish | |
| 2018 | 2,977 | (9, 573.1) | 2,860 | (64, 522.2) | No Fish | | No Fish | |
| 2019 | 3,420 | (7, 672.9) | 2,841 | (35, 587.0) | No Fish | | No Fish | |
| 2020 | 3,296 | (13, 412.6) | 2,445 | (7, 673.7) | 4,098 | (2, 101.8) | No Fish | |
| 2021 | 3,701 | (38, 600.8) | 2,834 | (8, 549.2) | No Fish | | No Fish | |
| Mean | 3,481 | | 3,045 | | 4,463 | | 3,689 | |
| SD | 632.2 | | 641.5 | | 856.3 | | 725.2 | |

Arrival and Spawn Timing Trends

We monitor peak arrival and spawn timing to determine whether the hatchery program has caused a shift to these features (Table 10). Peak arrival dates were based on the greatest number of fish trapped on a single day. Peak spawn in the hatchery was determined by the day when the most females were spawned. Peak spawning in the river was determined by the highest weekly redd count.

Peak arrival to the TFH adult trap for both natural and hatchery origin fish during 2021 was a little later than the historical average (Table 10). Peak spawning date in the hatchery was 14 September for both hatchery and natural origin fish and was similar to the historical mean (Table 10). The duration of spawning in the hatchery (28 days) was also close to the historical mean. Spawning in the river peaked on 9 September. The duration of active spawning in the Tucannon River (35 days) was also within the range found from previous years.

Natural origin adults typically arrive earlier at the adult trap and at a slightly faster rate than hatchery origin adults (Figure 8). On average, about half of the total run of hatchery origin adults typically arrive at the adult trap by 7 June (Figure 8). After the end of June, the hatchery adults tend to arrive at the adult trap at a slightly faster rate than natural origin fish. Jacks typically arrive later than the adults, with natural jacks arriving the latest of all (Figure 8).

Table 10. Peak dates of arrival of natural and hatchery salmon to the TFH adult trap and peak (date) and duration (number of days) for spawning in the hatchery and river, 1986-2021.

| Year | Peak Arrival at Trap | | Spawning in Hatchery | | | Spawning in River | |
|-------------------|----------------------|-------------|----------------------|-------------|-----------------|-------------------|------------------|
| | Natural | Hatchery | Natural | Hatchery | Duration | Combined | Duration |
| 1986 | 5/27 | – | 9/17 | – | 31 | 9/16 | 36 |
| 1987 | 5/15 | – | 9/15 | – | 29 | 9/23 | 35 |
| 1988 | 5/24 | – | 9/07 | – | 22 | 9/17 | 35 |
| 1989 | 6/06 | 6/12 | 9/15 | 9/12 | 29 | 9/13 | 36 |
| 1990 | 5/22 | 5/23 | 9/04 | 9/11 | 36 | 9/12 | 42 |
| 1991 | 6/11 | 6/04 | 9/10 | 9/10 | 29 | 9/18 | 35 |
| 1992 | 5/18 | 5/21 | 9/15 | 9/08 | 28 | 9/09 | 44 |
| 1993 | 5/31 | 5/27 | 9/13 | 9/07 | 30 | 9/08 | 52 |
| 1994 | 5/25 | 5/27 | 9/13 | 9/13 | 22 | 9/15 | 29 |
| 1995 ^a | – | 6/08 | 9/13 | 9/13 | 30 | 9/12 | 21 |
| 1996 | 6/06 | 6/20 | 9/17 | 9/10 | 21 | 9/18 | 35 |
| 1997 | 6/15 | 6/17 | 9/09 | 9/16 | 30 | 9/17 | 50 |
| 1998 | 6/03 | 6/16 | 9/08 | 9/16 | 36 | 9/17 | 16 |
| 1999 ^a | – | 6/16 | 9/07 | 9/14 | 22 | 9/16 | 23 |
| 2000 | 6/06 | 5/22 | – | 9/05 | 22 | 9/13 | 30 |
| 2001 | 5/23 | 5/23 | 9/11 | 9/04 | 20 | 9/12 | 35 |
| 2002 | 5/29 | 5/29 | 9/10 | 9/03 | 22 | 9/11 | 42 |
| 2003 | 5/25 | 5/25 | 9/09 | 9/02 | 36 | 9/12 | 37 |
| 2004 | 6/04 | 6/02 | 9/14 | 9/07 | 29 | 9/08 | 30 |
| 2005 | 6/01 | 5/31 | 9/06 | 9/06 | 28 | 9/14 | 28 |
| 2006 | 6/12 | 6/09 | 9/12 | 9/12 | 28 | 9/08 | --- ^b |
| 2007 | 6/04 | 6/04 | 9/18 | 9/04 | 22 | 9/12 | 30 |
| 2008 | 6/16 | 6/20 | 9/09 | 9/16 | 21 | 9/11 | 34 |
| 2009 | 6/01 | 6/15 | 9/15 | 9/08 | 29 | 9/10 | 37 |
| 2010 | 6/04 | 6/03 | 9/14 | 9/08 | 14 ^c | 9/10 | 33 |
| 2011 | 6/08 | 6/23 | 9/06 | 9/06 | 22 | 9/16 | 33 |
| 2012 | 5/30 | 6/02 | 9/11 | 9/18 | 22 | 9/12 | 36 |
| 2013 | 6/06 | 6/06 | 9/10 | 9/10 | 29 | 9/11 | 42 |
| 2014 | 5/27 | 6/04 | 9/09 | 9/09 | 22 ^c | 9/11 | 35 |
| 2015 | 5/18 | 5/20 | 9/15 | 9/08 | 29 | 9/09 | 44 |
| 2016 | 5/19 | 6/06 | 9/13 | 9/06 | 22 | 9/07 | 36 |
| 2017 | 6/06 | 6/18 | 9/12 | 9/12 | 29 | 9/11 | 26 |
| 2018 | 5/29 | 6/15 | 9/11 | 9/11 | 22 | 9/12 | 42 |
| 2019 | 5/31 | 6/04 | 9/10 | 9/10 | 22 | 9/11 | 38 |
| 2020 | 6/05 | 6/11 | 9/08 | 9/08 | 22 | 9/09 | 36 |
| Mean | 6/01 | 6/06 | 9/12 | 9/10 | 26 | 9/13 | 35 |
| 2021 | 6/05 | 6/11 | 9/14 | 9/14 | 28 | 9/09 | 35 |

^a Too few natural salmon were trapped in 1995 and 1999 to determine peak arrival.

^b Access restrictions during the Columbia Complex Forest Fire prohibited spawning ground surveys during the beginning of spawning.

^c Unspawned females determined to be in excess of eggtake goals were returned to the river for natural spawning which may have truncated duration of spawning in the hatchery.




Figure 8. Cumulative run timing by date at the Tucannon Fish Hatchery adult trap on the Tucannon River for both adult and jack natural and hatchery origin Tucannon River spring Chinook salmon, 1996-2021.

Total Run-Size

During 2021, no fish were passed above the trap. However, some adults were able to bypass the TFH trap/intake dam and we counted four redds above the trap. We calculated the number of fish above the trap by using the fish/redd estimate (2.15 – from the spawning escapement calculation in the next section) for an estimate of eight. We also multiplied the fish/redd estimate by the number of redds below the trap (31) for a total of 67 fish below the trap.

The run-size estimate for 2021 was calculated by adding the estimated number of fish upstream of the TFH adult trap (8), the estimated fish below the weir (67), adipose clipped strays killed at the trap (25), and the number of broodstock collected (115) (Table 11). Run-size for 2021 was estimated to be 215 fish (101 natural adults, 7 natural jacks, and 85 hatchery adults, 22 hatchery-origin jacks). Historical breakdowns are provided in Appendix D.

Table 11. Estimated spring Chinook salmon run to the Tucannon River and recovered pre-spawn mortalities (PSM), 1985-2021.

| Year ^a | Total Redds | Fish/Redd Ratio ^b | Potential Spawners | Broodstock Collected | Trap/Holding Mortalities ^c | Total Run-Size | River PSM ^d | Percent Natural |
|-------------------|-------------|------------------------------|--------------------|----------------------|---------------------------------------|----------------|------------------------|-----------------|
| 1985 ^e | 316 | 2.60 | 822 | 22 | 0 | 844 | 0 | 100 |
| 1986 | 200 | 2.60 | 520 | 116 | 0 | 636 | 0 | 100 |
| 1987 | 185 | 2.60 | 481 | 101 | 0 | 582 | 0 | 100 |
| 1988 | 117 | 2.60 | 304 | 125 | 0 | 429 | 0 | 96 |
| 1989 | 106 | 2.60 | 276 | 169 | 0 | 445 | 0 | 76 |
| 1990 | 180 | 3.39 | 610 | 135 | 1 | 746 | 7 | 66 |
| 1991 | 90 | 4.33 | 390 | 130 | 0 | 520 | 8 | 50 |
| 1992 | 200 | 2.82 | 564 | 97 | 11 | 672 | 81 | 58 |
| 1993 | 192 | 2.27 | 436 | 97 | 0 | 533 | 56 | 57 |
| 1994 | 44 | 1.59 | 70 | 70 | 0 | 140 | 0 | 70 |
| 1995 | 5 | 2.20 | 11 | 43 | 0 | 54 | 0 | 39 |
| 1996 | 69 | 2.00 | 138 | 80 | 5 | 223 | 29 | 64 |
| 1997 | 73 | 2.00 | 146 | 97 | 0 | 243 | 108 | 50 |
| 1998 | 26 | 1.94 | 51 | 89 | 0 | 140 | 4 | 61 |
| 1999 | 41 | 2.60 | 107 | 136 | 1 | 244 | 1 | 1 |
| 2000 | 92 | 2.60 | 239 | 81 | 17 | 337 | 2 | 24 |
| 2001 | 297 | 3.00 | 891 | 106 | 0 | 997 | 12 | 71 |
| 2002 | 299 | 3.00 | 897 | 107 | 0 | 1,004 | 1 | 35 |
| 2003 | 118 | 3.10 | 366 | 77 | 0 | 443 | 1 | 56 |
| 2004 | 160 | 3.00 | 480 | 92 | 0 | 572 | 1 | 70 |
| 2005 | 107 | 3.10 | 332 | 100 | 3 | 435 | 0 | 69 |
| 2006 | 109 | 1.60 | 174 | 89 | 3 | 266 | 0 | 57 |
| 2007 | 81 | 3.10 | 250 | 88 | 6 | 344 | 0 | 58 |
| 2008 | 199 | 4.10 | 1,056 | 134 | 1 | 1,191 | 0 | 45 |
| 2009 | 451 | 3.70 | 1,676 | 177 | 7 | 1,860 | 2 | 40 |
| 2010 | 481 | 4.87 | 2,341 | 173 | 9 | 2,523 | 2 | 57 |
| 2011 | 297 | 3.79 | 1,128 | 166 | 6 | 1,300 | 0 | 58 |
| 2012 | 169 | 6.30 | 1,059 | 170 | 6 | 1,235 | 4 | 66 |
| 2013 | 64 | 14.96 | 955 | 158 | 2 | 1,115 | 2 | 67 |
| 2014 | 124 | 7.70 | 959 | 127 | 0 | 1,086 | 18 | 83 |
| 2015 | 191 | 6.10 ^f | 1,604 | 131 | 42 | 1,777 | 28 | 41 |
| 2016 | 154 | 3.87 ^f | 478 | 126 | 148 | 752 | 6 | 30 |
| 2017 | 70 | 3.55 | 249 | 111 | 152 | 512 | 1 | 13 |
| 2018 | 109 | 2.02 ^f | 335 | 160 | 50 | 545 | 0 | 15 |
| 2019 | 11 | 1.97 | 22 | 170 ^g | 11 | 203 | 2 | 22 |
| 2020 | 14 | 1.84 | 26 | 53 | 2 | 81 | 1 | 68 |
| 2021 | 35 | 2.15 | 75 | 115 | 25 | 215 | 0 | 50 |

^a In 1994, 1995, 1998 and 1999, fish were not passed upstream, and in 1996 and 1997, high pre-spawning mortality occurred in fish passed above the trap, therefore; fish/redd ratio was based on the sex ratio of broodstock collected.

^b From 1985-1989 the TFH trap was temporary, thereby underestimating total fish passed upstream of the trap. The 1985-1989 fish/redd ratios were calculated from the 1990-1993 average, excluding 1991 because of a large jack run.

^c This total includes stray fish that are killed at the trap and pre-spawn mortalities of fish held at LFH for adult outplanting. During 2016, jacks were killed outright at the adult trap and are included in this total. During 2017, jacks were killed at LFH.

^d Effort in looking for pre-spawn mortalities has varied from year to year with more effort expended during years with poor conditions or large runs.

^e The 1985 redd counts were revised on the SASI database to account for all redds during the spawning season (WDFW 2017).

^f The fish/redd ratio was not used to estimate the number of fish below the adult trap due to survival differences between outplanted fish and fish that were passed upstream.

^g This total includes 149 adults kept for spawning and 21 jacks that were held and then outplanted but not recovered.

Spawning Escapement

To calculate spawning escapement, we assume one redd per female (Murdoch et al. 2009) and multiply the number of redds by the sex ratio (e.g., 1.02 males: 1 female = 2.02 fish/redd) of the pre-spawning population that was collected at the adult trap (i.e., no carcass collection bias issues). This should provide a more accurate expansion method than simply applying a constant value based on assumptions, or data from other studies, since it incorporates the natural variability that occurs in most populations (Murdoch et al. 2010). Because spawner distribution of hatchery and natural origin fish may be different, we expanded the natural and hatchery fish by reach [Wilderness, HMA (above trap), HMA (below trap), Hartsock, Marengo, and below Marengo] based on carcass recoveries. The total for all reaches equals the spawning escapement.

Sex ratio from the adult trap was only available from 2000 to present. For 1985 to 1999, we used corrected carcass data based on the methodology of Murdoch et al. (2010). For years when the corrected carcass data produced clear outliers, or produced spawning escapements greater than the run escapement, we used data cited by Meekin (1967) that cited an average of 2.20 adults/redd and proportionately adjusted that figure up during years with high jack returns. The spawning escapement for 2021 was 75 fish (25 natural-origin, 50 hatchery-origin) based on 2.15 fish per redd. The estimated spawning escapement for 1985 to 2021 is found in Table 12.

Table 12. Estimated spawning escapement and the calculation methodology used for the 1985 to 2021 run years.

| Run Year | Number of Redds | Spawning Escapement | Natural:Hatchery Ratio | Fish/Redd | Methodology |
|-------------------|------------------------|----------------------------|-------------------------------|------------------|-------------------------|
| 1985 ^a | 316 | 695 | 1.000:0.000 | 2.20 | Meekin (1967) |
| 1986 | 200 | 440 | 1.000:0.000 | 2.20 | Meekin (1967) |
| 1987 | 185 | 407 | 1.000:0.000 | 2.20 | Meekin (1967) |
| 1988 | 117 | 257 | 1.000:0.000 | 2.20 | Meekin (1967) |
| 1989 | 106 | 276 | 0.988:0.012 | 2.60 | Meekin (1967) |
| 1990 | 180 | 572 | 0.785:0.215 | 3.18 | Corrected Carcasses |
| 1991 | 90 | 291 | 0.677:0.323 | 3.23 | Corrected Carcasses |
| 1992 | 200 | 476 | 0.641:0.359 | 2.38 | Corrected Carcasses |
| 1993 | 192 | 397 | 0.617:0.383 | 2.07 | Corrected Carcasses |
| 1994 | 44 | 97 | 1.000:0.000 | 2.20 | Meekin (1967) |
| 1995 | 5 | 27 | 1.000:0.000 | 5.30 | Corrected Carcasses |
| 1996 | 69 | 152 | 0.767:0.233 | 2.20 | Meekin (1967) |
| 1997 | 73 | 105 | 0.644:0.356 | 1.44 | Corrected Carcasses |
| 1998 | 26 | 60 | 0.739:0.261 | 2.30 | Meekin (1967) |
| 1999 | 41 | 160 | 0.023:0.977 | 3.91 | Corrected Carcasses |
| 2000 | 92 | 201 | 0.307:0.693 | 2.18 | Sex ratio at Adult Trap |
| 2001 | 297 | 766 | 0.801:0.199 | 2.58 | Sex ratio at Adult Trap |
| 2002 | 299 | 568 | 0.395:0.605 | 1.90 | Sex ratio at Adult Trap |
| 2003 | 118 | 329 | 0.742:0.258 | 2.79 | Sex ratio at Adult Trap |
| 2004 | 160 | 346 | 0.826:0.174 | 2.16 | Sex ratio at Adult Trap |
| 2005 | 107 | 264 | 0.804:0.196 | 2.47 | Sex ratio at Adult Trap |
| 2006 | 109 | 202 | 0.759:0.241 | 1.85 | Sex ratio at Adult Trap |
| 2007 | 81 | 211 | 0.776:0.224 | 2.60 | Sex ratio at Adult Trap |
| 2008 | 199 | 796 | 0.610:0.390 | 4.00 | Sex ratio at Adult Trap |
| 2009 | 451 | 1191 | 0.507:0.493 | 2.64 | Sex ratio at Adult Trap |
| 2010 | 481 | 938 | 0.578:0.422 | 1.95 | Sex ratio at Adult Trap |
| 2011 | 297 | 849 | 0.703:0.297 | 2.86 | Sex ratio at Adult Trap |
| 2012 | 169 | 335 | 0.698:0.302 | 1.98 | Sex ratio at Adult Trap |
| 2013 | 64 | 170 | 0.697:0.303 | 2.66 | Sex ratio at Adult Trap |
| 2014 | 124 | 294 | 0.726:0.274 | 2.37 | Sex ratio at Adult Trap |
| 2015 | 191 | 523 | 0.330:0.670 | 2.74 | Sex ratio at Adult Trap |
| 2016 | 154 | 340 | 0.336:0.664 | 2.21 | Sex ratio at Adult Trap |
| 2017 | 70 | 249 | 0.195:0.805 | 3.55 | Sex ratio at Adult Trap |
| 2018 | 109 | 220 | 0.134:0.866 | 2.02 | Sex ratio at Adult Trap |
| 2019 | 11 | 22 | 0.364:0.636 | 1.97 | Sex ratio at Adult Trap |
| 2020 | 14 | 26 | 0.667:0.333 | 1.84 | Sex ratio at Adult Trap |
| 2021 | 35 | 75 | 0.333:0.667 | 2.15 | Sex ratio at Adult Trap |

^a The 1985 redd counts were revised on the SASI database to account for all redds during the spawning season (WDFW 2017).

Coded-Wire Tag Sampling

Broodstock collection, pre-spawn mortalities, and carcasses recovered during spawning ground surveys provide representatives of the annual run that can be sampled for CWT study groups (Table 13). In 2021, based on the estimated escapement of hatchery and natural origin fish to the river, we sampled approximately 72% of the run (Table 14).

Table 13. Coded-wire tag codes of hatchery salmon sampled at LFH and the Tucannon River, 2021.

| CWT Code | Broodstock Collected | | | Recovered in Tucannon River | | | Totals |
|---------------|----------------------|-----------------|-----------|-----------------------------|---------------------|-----------|-----------|
| | Pre-spawn Mortality | Killed Outright | Spawned | Dead in Trap ^a | Pre-spawn Mortality | Spawned | |
| 63-74-21 | | 3 | | | | | 3 |
| 63-73-96 | 1 | | 13 | | | | 14 |
| -Strays- | | | | | | | |
| 09-13-38 | | 1 | | | | | 1 |
| 09-13-40 | | | | 2 | | | 2 |
| 09-12-20 | | 1 | | | | 1 | 2 |
| 09-12-21 | | | | 1 | | | 1 |
| 09-12-23 | | | | 3 | | 2 | 5 |
| 09 | 1 | 8 | | | | 1 | 10 |
| Ad/No Wire | | 3 | | 19 | | 6 | 28 |
| Lost | | | 1 | | | | 1 |
| Totals | 2 | 16 | 14 | 25 | | 10 | 67 |

^a Adipose clipped strays are killed outright at the trap.

Table 14. Spring Chinook salmon (natural and hatchery) sampled from the Tucannon River, 2021.

| | 2021 | | |
|---|---------|----------|-------|
| | Natural | Hatchery | Total |
| <i>Total escapement to river</i> | 108 | 107 | 215 |
| Broodstock collected | 83 | 32 | 115 |
| Fish dead in adult trap ^a | 0 | 25 | 25 |
| Total hatchery sample | 83 | 57 | 140 |
| <i>Total fish left in river</i> | 25 | 50 | 75 |
| In-river pre-spawn mortalities observed | 0 | 0 | 0 |
| Spawned carcasses recovered | 5 | 10 | 15 |
| Total river sample | 5 | 10 | 15 |
| Carcasses sampled | 88 | 67 | 155 |

^a Adipose clipped strays are killed outright at the trap.

Stray Salmon into the Tucannon River

Spring Chinook from other river systems (strays) are periodically recovered in the Tucannon River, although they generally have been at a low proportion of the total run (Bumgarner et al. 2000). However, Umatilla River hatchery strays accounted for 8 and 12% of the total Tucannon River run in 1999 and 2000, respectively (Gallinat et al. 2001). Increased strays, particularly from the Umatilla River, was a concern since it exceeded the 5% stray proportion of hatchery fish deemed acceptable by NOAA Fisheries, and was contrary to fish management intent for the Tucannon River. In addition, the Oregon Department of Fish and Wildlife (ODFW) and the Confederated Tribes of the Umatilla Indian Reservation (CTUIR) did not mark a portion of Umatilla River origin spring Chinook with an RV or LV fin clip (65-70% of releases), or CWT for the 1997-1999 brood years. Because of that action, some stray fish that returned from those brood years were physically indistinguishable from natural origin Tucannon River spring Chinook. Scale samples were collected from adults in those brood years to determine hatchery-origin fish based on scale pattern analysis. However, we are unable to differentiate between unmarked Tucannon fish and unmarked strays based on scale patterns. Beginning with the 2000 BY, Umatilla River hatchery-origin spring Chinook were 100% marked (adipose clipped), however, the implementation of a “stepping stone” hatchery management protocol for the Umatilla Hatchery Program has resulted in a portion of Umatilla Hatchery releases being unclipped (but 100% CWT) beginning with the 2009 BY. Unfortunately, because of this, they are externally identical to Tucannon fish. As such, this hinders our ability to selectively remove stray hatchery fish during broodstock collection, or from fish passed upstream at the TFH adult trap. We will continue to monitor the Tucannon River and emphasize the need for external marks and CWTs for Umatilla River releases.

A total of 49 strays were recovered during 2021, comprised of 25 AD clip/no wire strays killed outright at the adult trap, 14 collected for broodstock (inadvertently), and ten recovered during spawning ground surveys (Appendix E). After expansions, strays accounted for an estimated 41.4% of the total 2021 run, which is the highest proportion of strays in the population since sampling began (Appendix E).

An added concern for the future is the implementation of WDFW’s new hatchery program for the Touchet River using Carson stock spring Chinook that began to return in 2021 (BY18). Potential straying from this hatchery program into the Tucannon River would be additive to the current stray rates being observed in the Tucannon River and could lead to outbreeding depression. All juveniles from the Touchet River spring Chinook program are 100% AD-clipped, with ~34% of the production receiving CWT and ~6% receiving PIT tags to monitor potential straying into the Tucannon River.

The increased use of passive integrated transponder (PIT) tags by fish and wildlife agencies and the utilization of in-stream PIT tag arrays in the Tucannon River have permitted us to identify the origin of some spring Chinook PIT tagged from other locations during 2021. Eight fish originally PIT tagged at locations other than the Tucannon River were detected in the Tucannon River (Table 15). The majority of the fish (4) were of unknown origin that were tagged as adults at Lower Granite Dam and eventually returned back downstream and entered the Tucannon River (Table 15). Two of these fish were identified as Tucannon stock and the other two were strays (Hells Canyon and South Fork Salmon River) based on genetic stock assignment results (Table 16). The other fish tagged as an adult at Bonneville Dam had a 09 Agency only (ODFW) CWT. Three hatchery origin fish originally tagged as juveniles from the Umatilla River, Yakima River, and Clearwater River were also detected in the Tucannon River during 2021 (Table 15).

Table 15. Tucannon River PIT tag array detections of spring Chinook originally tagged at locations other than the Tucannon River during 2021.

| PIT Tag | Origin | Tag Date | Life Stage At Tagging | Tag Release Location | Detection Date | Tucannon Site ^a |
|----------------|----------------|----------|-----------------------|----------------------|----------------|----------------------------|
| 3DD.0077963772 | H | 11/21/19 | Juvenile | Imeques AP, ODFW | 6/12/21 | TFH |
| 3DD.003D31BA97 | H | 3/25/20 | Juvenile | Roza Dam, Yakima | 10/23/21 | MTR |
| 3DD.003D669E64 | H | 9/23/20 | Juvenile | Clearwater, IDFG | 7/8/21 | LTR |
| 3DD.003D493B4B | W | 5/10/21 | Adult | Lower Granite Dam | 5/31/21 | TFH |
| 3DD.003D493B40 | W | 5/25/21 | Adult | Lower Granite Dam | 6/14/21 | UTR |
| 3DD.003D82A136 | H ^b | 5/26/21 | Adult | Bonneville Dam | 6/21/21 | TFH |
| 3DD.00775F0BF1 | W | 6/3/21 | Adult | Lower Granite Dam | 6/19/21 | UTR |
| 3DD.003D493972 | W | 6/30/21 | Adult | Lower Granite Dam | 8/20/21 | TFH |

^a PIT tag array locations are as follows: LTR – Lower Tucannon River (rkm 2.2), MTR – Middle Tucannon River (rkm 17.8), UTR – Upper Tucannon River (rkm 44.4), TFH – Tucannon Fish Hatchery (rkm 59.2).

^b This fish had a 09 (Oregon) agency only coded-wire tag.

Table 16. Genetic stock assignment results of adults PIT tagged at Lower Granite Dam and entering the Tucannon River based on Genetic Stock Identification (GSI) and Parentage Based Tagging (PBT) during 2021. (Data from Jesse McCane, PSMFC.)

| PIT Tag | Assigned Name | Genetic Stock | Stock Probability |
|----------------|-----------------|---------------|-------------------|
| 3DD.003D493B4B | OtsLGRU21S 0063 | HELLSC | 0.627 |
| 3DD.003D493B40 | OtsLGRU21S 0816 | TUCANO | 0.830 |
| 3DD.00775F0BF1 | OtsLGRU21S 1134 | TUCANO | 0.680 |
| 3DD.003D493972 | OtsLGRU21S 2050 | SFSALM | 0.563 |

Adult PIT Tag Returns

Five hundred ninety-five Tucannon River spring Chinook originally PIT tagged as juveniles have been detected returning to the Columbia River System (Table 17).

Table 17. Number of Tucannon River spring Chinook juvenile fish PIT tagged by origin and calendar year and adult returns detected (%) in the Columbia River System by origin.

| Tag Year | PIT Tagged Hatchery | PIT Tagged Natural | PIT Tagged Captive Brood | Detected H Adult Returns | Detected N Adult Returns | Detected CB Adult Returns |
|---------------|---------------------|--------------------|--------------------------|--------------------------|--------------------------|---------------------------|
| 1995 | 1,292 | --- | --- | 1 (0.08%) | --- | --- |
| 1996 | 1,923 | --- | --- | 0 | --- | --- |
| 1997 | 1,984 | --- | --- | 2 (0.10%) | --- | --- |
| 1998 | 1,999 | --- | --- | 0 | --- | --- |
| 1999 | 335 | 374 | --- | 2 (0.60%) | 5 (1.34%) | --- |
| 2000 | --- | --- | --- | --- | --- | --- |
| 2001 | 301 | 158 | --- | 0 | 0 | --- |
| 2002 | 318 | 321 | --- | 1 (0.31%) | 3 (0.93%) | --- |
| 2003 | 1,010 | --- | 1,007 | 3 (0.30%) | --- | 0 |
| 2004 | 1,012 | --- | 1,029 | 0 | --- | 0 |
| 2005 | 993 | 93 | 993 | 0 | 1 (1.08%) | 0 |
| 2006 | 1,001 | 70 | 1,002 | 1 (0.10%) | 1 (1.43%) | 0 |
| 2007 | 1,308 | 504 | 1,000 | 3 (0.23%) | 10 (1.98%) | 4 (0.40%) |
| 2008 | 4,989 | 1,915 | 997 | 47 (0.94%) | 47 (2.45%) | 6 (0.60%) |
| 2009 | 4,987 | 1,232 | --- | 13 (0.26%) | 17 (1.38%) | --- |
| 2010 | 15,000 | 2,800 | --- | 85 (0.57%) | 17 (0.61%) | --- |
| 2011 | 24,976 | 5,267 | --- | 38 (0.15%) | 23 (0.44%) | --- |
| 2012 | 22,982 | 3,889 | --- | 26 (0.11%) | 22 (0.57%) | --- |
| 2013 | 14,987 | 4,026 | --- | 32 (0.21%) | 41 (1.02%) | --- |
| 2014 | 14,969 | 660 | --- | 35 (0.23%) | 0 | --- |
| 2015 | 14,962 | 368 | --- | 25 (0.17%) | 1 (0.27%) | --- |
| 2016 | 14,983 | 1,429 | --- | 51 (0.34%) | 4 (0.28%) | --- |
| 2017 | 14,984 | 870 | --- | 16 (0.11%) | 1 (0.11%) | --- |
| 2018 | 14,937 | 366 | --- | 4 (0.03%) | 1 (0.27%) | --- |
| 2019 | 14,220 | 1,499 | --- | 1 (0.01%) | 5 (0.33%) | --- |
| 2020 | 14,987 | 911 | --- | 0 | 0 | --- |
| Totals | 205,439 | 26,752 | 6,028 | 386 (0.19%) | 199 (0.74%) | 10 (0.17%) |

From the detected returns, 142 (24%) of the returning PIT tagged spring Chinook were detected upstream of the Tucannon River (Table 18; Appendix F). Forty-three of these fish (7%) had their last detections at or above Lower Granite Dam (Table 18; Appendix F). The overshoot rate has generally decreased over time, and it is unknown whether this is related to changes in smolt release methods (from direct release to acclimation ponds with volitional release), changes in hydropower operations and river flows, changes in the proportion barged downstream, increases in tagging numbers/sample size, or greater detection capabilities in the Tucannon River (Table

18). This does not appear to be a hatchery effect as both natural and hatchery origin fish overshoot the Tucannon River (Table 18). Non-direct homing behavior has been documented for adult Chinook in the Columbia River System (Keefer et al. 2008a), and similar percentages of natural origin spring Chinook from the John Day River have been documented bypassing that river (Jim Ruzycki, ODFW, personal communication). However, more research into these events should be conducted to examine whether they are natural straying occurrences, or if it is related to hydropower operations. The installation of PIT tag arrays in the Tucannon River [Lower Tucannon River (LTR) at rkm 2.2 - 2005, Middle Tucannon River (MTR) at rkm 17.8 and Upper Tucannon River (UTR) at rkm 44.4 - 2011, and Tucannon Fish Hatchery (TFH) at rkm 59.2 – 2012] have enabled us to document that the majority of the Tucannon spring Chinook that overshoot are able to make it back (about 70%) to the Tucannon River (Table 18). Returning spring Chinook overshooting the Tucannon River continues to be a concern, especially if they are unable to return to the Tucannon River, or if they return in a more compromised state (e.g., injuries from additional dam crossings, added energy expenditure), and may partially explain why this population has been slow to respond to recovery and supplementation actions.

Table 18. Number and origin of PIT tagged Tucannon River spring Chinook returns that overshoot the Tucannon River (includes fish that were last detected returning downstream towards the Tucannon River) and also detected at Lower Granite Dam (LGR) that stayed above LGR Dam. Years with installed in-stream PIT tag arrays (2005 – 2019) are included for comparison.

| Tag Years | # Adult Detections Bonneville | Initial # Adults Above Tucannon R. | Initial Overshoot Rate | Percent Natural | Percent Hatchery | # Adults Above LGR | Percent Natural | Percent Hatchery | Overshoot Rate (%) |
|------------------|-------------------------------------|--|------------------------------|--------------------|---------------------|-----------------------|--------------------|---------------------|-----------------------|
| 1995-1999 | 10 | 8 | 80.0 | 37.5 | 62.5 | 8 | 37.5 | 62.5 | 80.0 |
| 2000-2004 | 7 | 2 | 28.6 | 50.0 | 50.0 | 2 | 50.0 | 50.0 | 28.6 |
| 2005-2009 | 150 | 20 | 13.3 | 35.0 | 65.0 | 14 | 42.9 | 57.1 | 9.3 |
| 2010-2014 | 319 | 80 | 25.1 | 37.5 | 62.5 | 12 | 41.7 | 58.3 | 3.8 |
| 2015-2019 | 109 | 32 | 29.4 | 3.1 | 96.9 | 7 | 0.0 | 100.0 | 6.4 |
| Totals | 595 | 142 | 23.9% | 29.6% | 70.4% | 43 | 34.9% | 65.1% | 7.2% |
| 2005-2019 | 578 | 132 | 22.8% | | | 33 | | | 5.7% |

Juvenile Salmon Evaluation

Hatchery Rearing, Marking, and Release

The BY20 supplementation juveniles (62,762) were tagged with CWT (63/79/43) at LFH from 15 March to 16 March 2021. The fish were transferred to TFH on 21 October 2021 for overwinter rearing. A total of 2,160 fish were sampled at TFH for precocity (external observation only) and mark/tag quality and 264 were sampled for length and weight statistics (Table 19). Twenty thousand fish designated for experimental release at the mouth of the Tucannon River were PIT tagged on 8-9 March 2022 and 20,000 fish from the release at TFH group (control) were PIT tagged on 9-10 March 2022. Detections of PIT tags will be used to compare outmigration survival and adult return estimates between the two groups.

Brood year 2020 fish from both groups were sampled just prior to release by WDFW evaluations staff (Table 19). The target release size was 38 g fish (12 fpp). Mortalities were scanned for PIT tags and 19,897 PIT tagged fish were released at TFH on 11 April 2022 and 19,667 PIT tagged fish were released at the mouth of the Tucannon River on 22 April 2022. We used PIT tag detections at the MTR and LTR antenna arrays to gauge timing of the release near the mouth. An estimated total of 62,020 BY20 smolts were released (42,046 release at TFH and 19,974 released near the mouth of the Tucannon River). Estimated numbers and size of fish released are provided in Table 20. Historical release numbers are found in Appendix G.

Table 19. Sample size (N), mean length (mm), coefficient of variation (CV), condition factor (K), mean weight (g), and precocity of 2020 BY juveniles sampled at TFH.

| Date | Group | N | Mean | | | Mean Wt. (g) | % Precocity ^a |
|---------|---------------|-----|-------------|------|------|--------------|--------------------------|
| | | | Length (mm) | CV | K | | |
| 1/19/22 | Combined | 264 | 136.4 | 12.2 | 1.28 | 33.7 | 1.53 |
| 4/11/22 | TFH Release | 305 | 149.4 | 10.7 | 1.17 | 40.2 | 0.66 |
| 4/11/22 | Mouth Release | 293 | 145.2 | 10.6 | 1.18 | 37.1 | 1.02 |

^a Based on external observations.

Table 20. Spring Chinook salmon released into the Tucannon River at TFH and the mouth of the Tucannon River, 2022 release year.

| Release Date | Release Location | CWT Code | Total Release | Number CWT | Size | |
|--------------|------------------|----------|---------------|------------|------------|----------|
| | | | | | Total (kg) | Mean (g) |
| 4/11 | TFH | 63/79/43 | 42,046 | 41,287 | 1,689 | 40.2 |
| 4/22 | Mouth | 63/79/43 | 19,974 | 19,613 | 743 | 37.1 |

Smolt Trapping

Evaluation staff operated a 1.5 m rotary screw trap at rkm 3 on the Tucannon River beginning on 1 October 2020 to estimate numbers of migrating juvenile natural spring Chinook. The smolt trap was pulled for the season on 6 July 2021. Numbers of each fish species captured by month during the 2021 outmigration can be found in Appendix H. Only 13 natural origin spring Chinook were captured during 2020/2021. The main outmigration of natural origin spring Chinook for the 2020/2021 outmigration occurred during the spring, with a limited outmigration during the fall and winter months (Figure 9). Prior years have shown increased outmigration in the fall and winter from larger adult returns (Gallinat and Ross 2014, Gallinat and Ross 2015), although even in those years, the majority of the outmigration occurred in the spring.




Figure 9. Emigration timing of natural spring Chinook salmon captured during smolt trap operations (rkm 3) on the Tucannon River for the 2020-21 migration year.

Natural spring Chinook emigrating from the Tucannon River (BY 2019) averaged 109 mm (Figure 10), with a CV of 10.1%. This is in comparison to a mean length of 139 mm for hatchery-origin fish (BY 2019) sampled at TFH (Gallinat and Kiefel 2021).




Figure 10. Length frequency distribution of sampled natural spring Chinook salmon captured in the Tucannon River smolt trap, 2020/2021 season.

Each week we attempted to determine trap efficiency by clipping a portion of the caudal fin on a representative subsample of captured migrants and releasing them approximately one kilometer upstream. The percent of marked fish recaptured was used as an estimate of weekly trapping efficiency. In previous reports we attempted to relate trap efficiency to abiotic factors such as stream flow or staff gauge level based on similar juvenile outmigration studies (Groot and Margolis 1991; Seiler et al. 1999; Cheng and Gallinat 2004). We found no significant relationships.

To estimate potential juvenile migrants passing when the trap was not operated for short intervals (≤ 5 days), such as periods when freshets washed out large amounts of debris from the river, we calculated the mean number of fish trapped for three days before and three days after non-trapping periods. The mean number of fish trapped daily was then divided by the estimated trap efficiency to calculate fish passage. The estimated number of fish passing each day was then applied to each day the trap was not operated.

We estimated outmigration based on the approach of Steinhorst et al. (2004). This involved using a Bailey-modified Lincoln-Peterson estimation with 95% bootstrap confidence intervals by running the Gauss Run-Time computer program (version 7.0). Bootstrap iterations numbered 1,000. The program allows for the division of the out-migration trapping season into strata with

similar capture efficiencies as long as at least seven marked recaptures occurred. Strata with less than seven recaptures were grouped with either the preceding or following stratum, depending upon similarity in trapping/flow conditions. Where river conditions were similar, we used our best judgment to group the strata.

Several assumptions are required to attain unbiased estimates of smolt production. How well the assumptions are met will determine the accuracy and precision of the estimates. Some of these assumptions are:

- Survival from release to the trap was 100%.
- All marked fish are identified and correctly enumerated.
- Fish do not lose their marks.
- All fish in the tag release group emigrate (i.e., do not residualize in the area of release).
- Marked fish are caught at the same rate as unmarked fish.

Due to the low numbers of captured spring Chinook in 2021, we were unable to estimate an outmigration number using our standard methods described above. Instead, we used the average trapping efficiency (7.49%) for spring Chinook for the last five years. We therefore estimate that only 174 natural origin spring Chinook (2019 BY) emigrated from 1 October 2020 to 6 July 2021.

Smolt Migration to Lower Monumental and McNary Dams

With the use of PIT tags, we monitored the migration travel time and speed of juvenile spring Chinook from the Tucannon River (both hatchery and natural origin) to Lower Monumental and McNary Dams for the 2021 outmigration (Table 21). Hatchery fish were PIT tagged on 8 March 2021, about two weeks prior to release, while natural origin fish were PIT tagged at the Tucannon River smolt trap (rkm 3), described earlier.

Hatchery fish were direct stream released at the TFH outlet (rkm 58) on 24 March 2021. Natural origin spring Chinook were released immediately following PIT tagging at the smolt trap, so the release date/time provided in the PTAGIS tagging files have been used for travel time/speed calculations. A total of 13 natural origin fish were PIT tagged at the smolt trap and released between 13 November 2020 to 20 May 2021 with only one of those fish detected at Lower Monumental Dam after five days of travel (a travel rate of 14 km/day). In contrast, it took about thirty days for hatchery fish to reach Lower Monumental dam at an average rate of around 5 km/day (Table 21). Historically, natural-origin fish have faster migration time to the dams than hatchery-origin fish (Figure 11).

From 2007 to 2017, barge transportation at Lower Monumental Dam typically began between 1-12 May (PTAGIS website 2020). From 2018 on, transportation has begun around 23 April. For 2018 and 2019, spring Chinook were released later (last week of April, first week of May) per the request of the co-managers to allow for greater potential transportation at Lower Monumental Dam. However, those BYs (2016 and 2017) have had poor adult returns.

Table 21. Median and mean travel time and outmigration speed of hatchery-origin Tucannon River spring Chinook to Lower Monumental and McNary Dams in 2021.

| Release Dates | Sample Size | Median Travel Days | Mean Travel Days | Mean Travel Days (S.D.) | Median Travel Speed (km/day) | Mean Travel Speed (km/day) | Mean Travel Speed S.D. |
|---|-------------|--------------------|------------------|-------------------------|------------------------------|----------------------------|------------------------|
| Hatchery-origin – Lower Monumental Dam | | | | | | | |
| 3/24/21 | 91 | 31.0 | 29.4 | 7.8 | 4.0 | 4.8 | 2.5 |
| Hatchery-origin – McNary Dam | | | | | | | |
| 3/24/21 | 87 | 35.0 | 32.9 | 9.3 | 5.1 | 6.0 | 2.2 |




Figure 11. The cumulative timing to Lower Monumental Dam for natural and hatchery origin Tucannon River spring Chinook emigrants from 2021 compared to the 2006-2020 average.

Survival Rates

Point estimates of population sizes have been calculated for various life stages (Tables 22 and 23) of natural and hatchery-origin spring Chinook from spawning ground and juvenile mid-summer population surveys, smolt trapping, and fecundity estimates. Survivals between life stages have been calculated for both natural and hatchery salmon to assist in the evaluation of the hatchery program. These survival estimates provide insight as to where efforts should be directed to improve not only the survival of fish produced within the hatchery, but fish in the river as well.

As expected, juvenile (egg-parr-smolt) survival rates for hatchery fish are considerably higher than for naturally reared salmon (Table 24) because they have been protected in the hatchery. However, SARs to the Tucannon River of natural salmon were eight times higher (based on geometric means) than for hatchery-reared salmon (Tables 25 and 26). With the exception of the 2006 brood year, hatchery SARs (mean 0.23%; geometric mean 0.15%) documented from the 1985-2016 broods have been well below the original LSRCP survival assumption of 0.87% (which was used to size the original hatchery program of 132,000 smolts). Hatchery SARs for Tucannon River salmon need to substantially improve in order to help meet the mitigation goal of 1,152 hatchery adult salmon. The target size at release was increased to 38 g fish (12 fpp) beginning with the 2011 brood year in an attempt to improve poor smolt-to-adult return survival rates.

Table 22. Estimates of natural in-river produced Tucannon spring Chinook salmon (both hatchery and natural origin parents) abundance by life stage for 1985-2021 broods.

| Brood Year | Females in River | | Mean Fecundity ^a | | Number of Eggs | Number of Parr ^b | Number of Smolts | Returning Progeny ^c |
|-------------------|------------------|----------|-----------------------------|----------|----------------|-----------------------------|---------------------|--------------------------------|
| | Natural | Hatchery | Natural | Hatchery | | | | |
| 1985 ^d | 316 | --- | 3,883 | --- | 1,227,028 | 90,200 | 35,559 | 392 |
| 1986 | 200 | --- | 3,916 | --- | 783,200 | 102,600 | 51,004 | 467 |
| 1987 | 185 | --- | 4,096 | --- | 757,760 | 79,100 | 52,349 | 228 |
| 1988 | 117 | --- | 3,882 | --- | 454,194 | 69,100 | 35,925 | 502 |
| 1989 | 103 | 3 | 3,883 | 2,606 | 407,767 | 58,600 | 19,107 | 153 |
| 1990 | 128 | 52 | 3,993 | 2,697 | 651,348 | 86,259 | 32,969 | 94 |
| 1991 | 51 | 40 | 3,741 | 2,517 | 291,471 | 54,800 | 30,000 ^e | 7 |
| 1992 | 119 | 81 | 3,854 | 3,295 | 725,521 | 103,292 | 36,749 | 161 |
| 1993 | 112 | 80 | 3,701 | 3,237 | 673,472 | 86,755 | 34,623 | 177 |
| 1994 | 39 | 5 | 4,187 | 3,314 | 179,863 | 12,720 | 4,957 | 12 |
| 1995 | 5 | 0 | 5,224 | 0 | 26,120 | 0 | 75 ^e | 6 |
| 1996 | 53 | 16 | 3,516 | 2,843 | 231,836 | 2,845 | 2,906 | 69 |
| 1997 | 39 | 34 | 3,609 | 3,315 | 253,461 | 32,913 | 25,553 | 791 |
| 1998 | 19 | 7 | 4,023 | 3,035 | 97,682 | 8,453 | 4,849 | 388 |
| 1999 | 1 | 40 | 3,965 | 3,142 | 129,645 | 15,944 | 8,721 | 141 |
| 2000 | 26 | 66 | 3,969 | 3,345 | 323,964 | 44,618 | 29,442 | 448 |
| 2001 | 219 | 78 | 3,612 | 3,252 | 1,044,684 | 63,412 | 42,416 | 257 |
| 2002 | 104 | 195 | 3,981 | 3,368 | 1,070,784 | 72,197 | 64,036 | 212 |
| 2003 | 67 | 51 | 3,789 | 3,812 | 448,275 | 40,900 | 27,724 | 173 |
| 2004 | 117 | 43 | 3,444 | 2,601 | 514,791 | 30,809 | 21,057 | 399 |
| 2005 | 82 | 25 | 3,773 | 2,903 | 381,961 | 21,162 | 17,579 | 739 |
| 2006 | 73 | 36 | 2,887 | 2,654 | 306,295 | --- | 30,228 | 1,720 |
| 2007 | 50 | 31 | 3,847 | 2,869 | 281,289 | --- | 8,529 | 610 |
| 2008 | 95 | 104 | 3,732 | 3,020 | 668,620 | --- | 14,778 | 884 |
| 2009 | 178 | 273 | 3,639 | 3,267 | 1,539,633 | --- | 45,538 | 619 |
| 2010 | 278 | 203 | 3,579 | 3,195 | 1,643,547 | --- | 35,080 | 938 |
| 2011 | 175 | 122 | 4,230 | 3,301 | 1,142,972 | --- | 23,376 | 727 |
| 2012 | 115 | 54 | 3,151 | 2,563 | 500,767 | --- | 12,886 | 213 |
| 2013 | 44 | 20 | 3,798 | 3,185 | 230,812 | --- | 3,831 | 69 |
| 2014 | 105 | 19 | 3,699 | 3,290 | 450,905 | --- | 6,604 | 89 |
| 2015 | 64 | 127 | 3,839 | 3,468 | 686,132 | --- | 14,305 | 45 |
| 2016 | 53 | 101 | 3,704 | 3,179 | 517,391 | --- | 8,058 | 57 |
| 2017 | 12 | 58 | 3,393 | 3,034 | 216,688 | --- | 17,972 | 98 |
| 2018 | 12 | 97 | 2,977 | 2,860 | 313,144 | --- | 16,979 | 8 |
| 2019 | 4 | 7 | 3,420 | 2,841 | 33,567 | --- | 174 | |
| 2020 | 9 | 5 | 3,403 | 2,445 | 42,852 | --- | | |
| 2021 | 12 | 23 | 3,701 | 2,834 | 109,594 | --- | | |

^a 1985 and 1989 mean fecundity of natural females is the average of 1986-88 and 1990-93 brood years.

^b Number of parr estimated from electrofishing (1985-1989), Line transect snorkel surveys (1990-1992), and Total Count snorkel surveys (1993-2005).

^c Numbers do not include down river harvest or other out-of-basin recoveries.

^d The 1985 redd counts were revised on the SASI database to account for all redds during the spawning season (WDFW 2017).

^e Smolt estimates could not be estimated with the GAUSS program for the 1991 and 1995 brood years. Numbers of smolts for those brood years were obtained from estimates in the annual reports.

Table 23. Estimates of Tucannon spring Chinook salmon abundance (*spawned and reared in the hatchery*) by life stage for 1985-2021 broods.

| Brood Year | Females Spawned | | Mean Fecundity ^a | | Number of Eggs | Number of Parr | Number of Smolts | Returning Progeny ^b |
|------------|-----------------|----------|-----------------------------|----------|----------------|----------------|----------------------|--------------------------------|
| | Natural | Hatchery | Natural | Hatchery | | | | |
| 1985 | 4 | --- | 3,883 | --- | 14,843 | 13,401 | 12,922 | 45 |
| 1986 | 57 | --- | 3,916 | --- | 187,958 | 177,277 | 152,725 | 319 |
| 1987 | 48 | --- | 4,096 | --- | 196,573 | 164,630 | 152,165 | 178 |
| 1988 | 49 | --- | 3,882 | --- | 182,438 | 150,677 | 145,146 | 385 |
| 1989 | 28 | 9 | 3,883 | 2,606 | 133,521 | 103,420 | 99,057 | 209 |
| 1990 | 21 | 23 | 3,993 | 2,697 | 126,334 | 89,519 | 85,737 | 28 |
| 1991 | 17 | 11 | 3,741 | 2,517 | 91,275 | 77,232 | 74,064 | 25 |
| 1992 | 28 | 18 | 3,854 | 3,295 | 156,359 | 151,727 | 87,752 ^c | 76 |
| 1993 | 21 | 28 | 3,701 | 3,237 | 168,366 | 145,303 | 138,848 | 138 |
| 1994 | 22 | 21 | 4,187 | 3,314 | 161,707 | 132,870 | 130,069 | 32 |
| 1995 | 6 | 15 | 5,224 | 0 | 85,772 | 63,935 | 62,144 | 177 |
| 1996 | 18 | 19 | 3,516 | 2,843 | 117,287 | 80,325 | 76,219 | 265 |
| 1997 | 17 | 25 | 3,609 | 3,315 | 144,237 | 29,650 | 24,186 | 176 |
| 1998 | 30 | 14 | 4,023 | 3,035 | 161,019 | 136,027 | 127,939 | 793 |
| 1999 | 1 | 36 | 3,965 | 3,142 | 113,544 | 106,880 | 97,600 | 33 |
| 2000 | 3 | 35 | 3,969 | 3,345 | 128,980 | 123,313 | 102,099 | 157 |
| 2001 | 29 | 27 | 3,612 | 3,252 | 184,127 | 174,934 | 146,922 | 127 |
| 2002 | 22 | 25 | 3,981 | 3,368 | 169,364 | 151,531 | 123,586 | 121 |
| 2003 | 17 | 20 | 3,789 | 3,812 | 140,658 | 126,400 | 71,154 | 71 |
| 2004 | 28 | 18 | 3,444 | 2,601 | 140,459 | 128,877 | 67,542 | 120 |
| 2005 | 25 | 24 | 3,773 | 2,903 | 161,345 | 151,466 | 149,466 | 690 |
| 2006 | 18 | 27 | 2,887 | 2,654 | 123,629 | 112,350 | 106,530 | 1,122 |
| 2007 | 27 | 9 | 3,847 | 2,869 | 124,543 | 117,182 | 114,681 | 261 |
| 2008 | 17 | 43 | 3,732 | 3,020 | 193,324 | 183,925 | 172,897 | 643 |
| 2009 | 42 | 54 | 3,639 | 3,267 | 323,341 | 292,291 | 231,437 ^d | 300 |
| 2010 | 39 | 44 | 3,579 | 3,195 | 279,969 | 237,861 | 201,585 | 194 |
| 2011 | 45 | 41 | 4,230 | 3,301 | 325,701 | 305,215 | 259,964 | 711 |
| 2012 | 48 | 47 | 3,151 | 2,563 | 269,514 | 246,033 | 203,510 | 514 |
| 2013 | 48 | 30 | 3,798 | 3,185 | 275,188 | 263,630 | 207,859 | 362 |
| 2014 | 39 | 27 | 3,699 | 3,290 | 231,026 | 226,300 | 221,099 | 458 |
| 2015 | 55 | 20 | 3,839 | 3,468 | 280,519 | 266,134 | 199,686 | 165 |
| 2016 | 31 | 41 | 3,704 | 3,179 | 245,174 | 230,106 | 209,031 | 50 |
| 2017 | 8 | 52 | 3,393 | 3,034 | 181,664 | 166,590 | 144,219 | 17 |
| 2018 | 9 | 67 | 2,977 | 2,860 | 212,973 | 204,364 | 192,521 | 3 |
| 2019 | 7 | 38 | 3,420 | 2,841 | 126,102 | 118,159 | 80,995 | |
| 2020 | 15 | 7 | 3,403 | 2,445 | 68,155 | 66,227 | 62,020 | |
| 2021 | 38 | 9 | 3,701 | 2,834 | 166,237 | 145,707 | | |

^a 1985 and 1989 mean fecundity of natural females is the average of 1986-88 and 1990-93 brood years; 1999 mean fecundity of natural fish is based on the mean of 1986-1998 brood years.

^b Numbers do not include down river harvest or other out-of-basin recoveries.

^c Number of smolts is less than actual release number. 57,316 parr were released in October 1993, with an estimated 7% survival. Total number of hatchery fish released from the 1992 brood year was 140,725. We therefore use the listed number of 87,752 as the number of smolts released.

^d Parr determined to be in excess of program goals were released at Russell Springs and are not included in number of parr and smolts.

Table 24. Percent survival by brood year for juvenile salmon and the multiplicative advantage of hatchery-reared salmon over naturally-reared salmon in the Tucannon River.

| Brood Year | Natural | | | Hatchery | | | Hatchery Advantage | | |
|-------------|-------------|---------------|--------------|-------------|---------------|--------------|--------------------|---------------|--------------|
| | Egg to Parr | Parr to Smolt | Egg to Smolt | Egg to Parr | Parr to Smolt | Egg to Smolt | Egg to Parr | Parr to Smolt | Egg to Smolt |
| 1985 | 7.4 | 39.4 | 2.9 | 90.3 | 96.4 | 87.1 | 12.3 | 2.4 | 30.0 |
| 1986 | 13.1 | 49.7 | 6.5 | 94.3 | 86.2 | 81.3 | 7.2 | 1.7 | 12.5 |
| 1987 | 10.4 | 66.2 | 6.9 | 83.8 | 92.4 | 77.4 | 8.0 | 1.4 | 11.2 |
| 1988 | 15.2 | 52.0 | 7.9 | 82.6 | 96.3 | 79.6 | 5.4 | 1.9 | 10.1 |
| 1989 | 14.4 | 32.6 | 4.7 | 77.5 | 95.8 | 74.2 | 5.4 | 2.9 | 15.8 |
| 1990 | 13.2 | 38.2 | 5.1 | 70.9 | 95.8 | 67.9 | 5.4 | 2.5 | 13.4 |
| 1991 | 18.8 | 54.7 | 10.3 | 84.6 | 95.9 | 81.1 | 4.5 | 1.8 | 7.9 |
| 1992 | 14.2 | 35.6 | 5.1 | 97.0 | 57.8 | 56.1 | 6.8 | 1.6 | 11.1 |
| 1993 | 12.9 | 39.9 | 5.1 | 86.3 | 95.6 | 82.5 | 6.7 | 2.4 | 16.0 |
| 1994 | 7.1 | 39.0 | 2.8 | 82.2 | 97.9 | 80.4 | 11.6 | 2.5 | 29.2 |
| 1995 | 0.0 | 0.0 | 0.3 | 74.5 | 97.2 | 72.5 | --- | --- | --- |
| 1996 | 1.2 | 102.1 | 1.3 | 68.5 | 94.9 | 65.0 | 55.8 | 0.9 | 51.8 |
| 1997 | 13.0 | 77.6 | 10.1 | 20.6 | 81.6 | 16.8 | 1.6 | 1.1 | 1.7 |
| 1998 | 8.7 | 57.4 | 5.0 | 84.5 | 94.1 | 79.5 | 9.8 | 1.6 | 16.0 |
| 1999 | 12.3 | 54.7 | 6.7 | 94.1 | 91.3 | 86.0 | 7.7 | 1.7 | 12.8 |
| 2000 | 13.8 | 66.0 | 9.1 | 95.6 | 82.8 | 79.2 | 6.9 | 1.3 | 8.7 |
| 2001 | 6.1 | 66.9 | 4.1 | 95.0 | 84.0 | 79.8 | 15.7 | 1.3 | 19.7 |
| 2002 | 6.7 | 88.7 | 6.0 | 89.5 | 81.6 | 73.0 | 13.3 | 0.9 | 12.2 |
| 2003 | 9.1 | 67.8 | 6.2 | 89.9 | 56.3 | 50.6 | 9.8 | 0.8 | 8.2 |
| 2004 | 6.0 | 68.3 | 4.1 | 91.8 | 52.4 | 48.1 | 15.3 | 0.8 | 11.8 |
| 2005 | 5.5 | 83.1 | 4.6 | 93.9 | 98.7 | 92.6 | 16.9 | 1.2 | 20.1 |
| 2006 | --- | --- | 9.9 | 90.9 | 94.8 | 86.2 | --- | --- | 8.7 |
| 2007 | --- | --- | 3.0 | 94.1 | 97.9 | 92.1 | --- | --- | 30.4 |
| 2008 | --- | --- | 2.2 | 95.1 | 94.0 | 89.4 | --- | --- | 40.5 |
| 2009 | --- | --- | 3.0 | 90.4 | 79.2 | 71.6 | --- | --- | 24.2 |
| 2010 | --- | --- | 2.1 | 85.0 | 84.7 | 72.0 | --- | --- | 33.7 |
| 2011 | --- | --- | 2.0 | 93.7 | 85.2 | 79.8 | --- | --- | 39.0 |
| 2012 | --- | --- | 2.6 | 91.3 | 82.7 | 75.5 | --- | --- | 29.3 |
| 2013 | --- | --- | 1.7 | 95.8 | 78.8 | 75.5 | --- | --- | 45.5 |
| 2014 | --- | --- | 1.5 | 98.0 | 97.7 | 95.7 | --- | --- | 65.3 |
| 2015 | --- | --- | 2.1 | 94.9 | 75.0 | 71.2 | --- | --- | 34.1 |
| 2016 | --- | --- | 1.6 | 93.9 | 90.8 | 85.3 | --- | --- | 54.7 |
| 2017 | --- | --- | 8.3 | 91.7 | 86.6 | 79.4 | --- | --- | 9.6 |
| 2018 | --- | --- | 5.4 | 96.0 | 94.2 | 90.4 | --- | --- | 16.7 |
| 2019 | --- | --- | 0.5 | 93.7 | 68.5 | 64.2 | --- | --- | 123.9 |
| 2020 | | | | 97.2 | 93.6 | 91.0 | | | |
| 2021 | | | | 87.7 | | | | | |
| Mean | 10.0 | 56.2 | 4.6 | 87.5 | 86.9 | 75.8 | 11.3 | 1.6 | 25.8 |
| SD | 4.8 | 22.7 | 2.8 | 13.5 | 12.2 | 15.0 | 11.2 | 0.6 | 23.2 |

Table 25. Adult returns and SARs of natural salmon to the Tucannon River for brood years 1985-2018. (2017 and 2018 are incomplete brood years included for comparison.)

| Brood Year | Estimated Number of Smolts | Number of Adult Returns, observed (Obs) and expanded (exp) ^a | | | | | | SAR (%) | |
|-----------------------|----------------------------|---|-----|-------|-------|-------|-----|-------------------------|-------------------------|
| | | Age 3 | | Age 4 | | Age 5 | | With Jacks | No Jacks |
| | | Obs | Exp | Obs | Exp | Obs | Exp | | |
| 1985 | 35,559 | 8 | 19 | 110 | 255 | 36 | 118 | 1.10 | 1.05 |
| 1986 ^b | 51,004 | 1 | 2 | 115 | 375 | 28 | 90 | 0.92 | 0.91 |
| 1987 | 52,349 | 0 | 0 | 52 | 167 | 29 | 61 | 0.44 | 0.44 |
| 1988 | 35,925 | 1 | 3 | 136 | 318 | 74 | 181 | 1.40 | 1.39 |
| 1989 | 19,107 | 5 | 12 | 47 | 115 | 23 | 26 | 0.80 | 0.74 |
| 1990 | 32,969 | 3 | 8 | 63 | 72 | 12 | 14 | 0.29 | 0.26 |
| 1991 | 30,000 ^c | 0 | 0 | 4 | 5 | 1 | 2 | 0.02 | 0.02 |
| 1992 | 36,749 | 2 | 2 | 84 | 138 | 16 | 21 | 0.44 | 0.43 |
| 1993 | 34,623 | 1 | 2 | 62 | 100 | 58 | 75 | 0.51 | 0.51 |
| 1994 | 4,957 | 0 | 0 | 8 | 10 | 1 | 2 | 0.24 | 0.24 |
| 1995 | 75 ^c | 0 | 0 | 1 | 1 | 2 | 5 | 8.00 | 8.00 |
| 1996 | 2,906 | 0 | 0 | 27 | 63 | 2 | 6 | 2.37 | 2.37 |
| 1997 | 25,553 | 6 | 14 | 234 | 695 | 29 | 82 | 3.10 | 3.04 |
| 1998 | 4,849 | 3 | 9 | 91 | 259 | 43 | 120 | 8.00 | 7.82 |
| 1999 | 8,721 | 3 | 9 | 44 | 124 | 3 | 8 | 1.62 | 1.51 |
| 2000 | 29,442 | 1 | 3 | 148 | 392 | 16 | 53 | 1.52 | 1.51 |
| 2001 | 42,416 | 0 | 0 | 73 | 246 | 5 | 11 | 0.61 | 0.61 |
| 2002 | 64,036 | 1 | 3 | 68 | 134 | 36 | 75 | 0.33 | 0.33 |
| 2003 | 27,724 | 4 | 7 | 55 | 115 | 21 | 51 | 0.62 | 0.60 |
| 2004 | 21,057 | 4 | 8 | 147 | 352 | 19 | 39 | 1.89 | 1.86 |
| 2005 | 17,579 | 23 | 131 | 260 | 595 | 2 | 13 | 4.20 | 3.46 |
| 2006 | 30,228 | 32 | 116 | 298 | 1,389 | 73 | 215 | 5.69 | 5.31 |
| 2007 | 8,529 | 4 | 41 | 133 | 456 | 22 | 113 | 7.15 | 6.67 |
| 2008 | 14,778 | 10 | 85 | 150 | 693 | 23 | 106 | 5.98 | 5.41 |
| 2009 | 45,538 | 1 | 7 | 94 | 554 | 10 | 58 | 1.36 | 1.34 |
| 2010 | 35,080 | 3 | 91 | 136 | 799 | 17 | 48 | 2.67 | 2.41 |
| 2011 | 23,376 | 3 | 41 | 145 | 619 | 31 | 67 | 3.11 | 2.93 |
| 2012 | 12,886 | 4 | 65 | 64 | 148 | 0 | 0 | 1.65 | 1.15 |
| 2013 | 3,831 | 2 | 8 | 25 | 60 | 1 | 1 | 1.80 | 1.59 |
| 2014 | 6,604 | 9 | 9 | 44 | 79 | 1 | 1 | 1.35 | 1.21 |
| 2015 | 14,305 | 0 | 0 | 36 | 42 | 3 | 3 | 0.31 | 0.31 |
| 2016 | 8,058 | 1 | 2 | 34 | 50 | 1 | 5 | 0.71 | 0.68 |
| 2017 | 17,972 | 3 | 3 | 79 | 95 | --- | --- | 0.55 | 0.53 |
| 2018 | 16,979 | 8 | 8 | --- | --- | --- | --- | 0.05 | 0.00 |
| Mean | | | | | | | | 2.19^d | 2.07^d |
| Geometric Mean | | | | | | | | 1.21^d | 1.15^d |

^a Expanded numbers are calculated from the proportion of each known age salmon recovered in the river and from broodstock collections in relation to the total estimated return to the Tucannon River. Expansions do not include down river harvest or Tucannon River fish straying to other systems.

^b One known (expanded to two) Age 6 salmon was recovered.

^c Numbers of smolts obtained from estimates in the annual reports.

^d The 2017 and 2018 SARs are not included in the mean.

Table 26. Adult returns and SARs of hatchery salmon to the Tucannon River for brood years 1985-2018. (2017 and 2018 are incomplete brood years included for comparison.)

| Brood Year | Estimated Number of Smolts | Number of Adult Returns, observed (obs) and expanded (exp) ^a | | | | | | SAR (%) | |
|-----------------------|----------------------------|---|-----|-------|-----|-------|-----|-------------------------|-------------------------|
| | | Age 3 | | Age 4 | | Age 5 | | With Jacks | No Jacks |
| | | Obs | Exp | Obs | Exp | Obs | Exp | | |
| 1985 | 12,922 | 9 | 19 | 25 | 26 | 0 | 0 | 0.35 | 0.20 |
| 1986 | 152,725 | 79 | 83 | 99 | 220 | 8 | 16 | 0.21 | 0.15 |
| 1987 | 152,165 | 9 | 19 | 70 | 145 | 8 | 14 | 0.12 | 0.10 |
| 1988 | 145,146 | 46 | 99 | 140 | 244 | 26 | 42 | 0.27 | 0.20 |
| 1989 | 99,057 | 7 | 13 | 100 | 179 | 14 | 17 | 0.21 | 0.20 |
| 1990 | 85,737 | 3 | 6 | 16 | 20 | 2 | 2 | 0.03 | 0.03 |
| 1991 | 74,064 | 4 | 5 | 20 | 20 | 0 | 0 | 0.03 | 0.03 |
| 1992 | 87,752 | 11 | 11 | 50 | 63 | 2 | 2 | 0.09 | 0.07 |
| 1993 | 138,848 | 11 | 15 | 93 | 107 | 15 | 16 | 0.10 | 0.09 |
| 1994 | 130,069 | 2 | 4 | 21 | 23 | 4 | 5 | 0.02 | 0.02 |
| 1995 | 62,144 | 13 | 16 | 117 | 157 | 2 | 4 | 0.28 | 0.26 |
| 1996 | 76,219 | 44 | 59 | 100 | 192 | 5 | 14 | 0.35 | 0.27 |
| 1997 | 24,186 | 7 | 13 | 59 | 163 | 0 | 0 | 0.73 | 0.67 |
| 1998 | 127,939 | 36 | 97 | 174 | 546 | 39 | 150 | 0.62 | 0.54 |
| 1999 | 97,600 | 3 | 11 | 5 | 19 | 1 | 3 | 0.03 | 0.02 |
| 2000 | 102,099 | 7 | 26 | 47 | 131 | 0 | 0 | 0.15 | 0.13 |
| 2001 | 146,922 | 7 | 19 | 51 | 107 | 1 | 1 | 0.09 | 0.07 |
| 2002 | 123,586 | 3 | 6 | 60 | 99 | 6 | 16 | 0.10 | 0.09 |
| 2003 | 71,154 | 1 | 2 | 23 | 65 | 2 | 4 | 0.10 | 0.10 |
| 2004 | 67,542 | 7 | 18 | 59 | 98 | 2 | 4 | 0.18 | 0.15 |
| 2005 | 149,466 | 50 | 291 | 180 | 399 | 0 | 0 | 0.46 | 0.27 |
| 2006 | 106,530 | 60 | 402 | 180 | 679 | 19 | 41 | 1.05 | 0.68 |
| 2007 | 114,681 | 7 | 74 | 76 | 171 | 5 | 16 | 0.23 | 0.16 |
| 2008 | 172,897 | 27 | 269 | 104 | 369 | 6 | 5 | 0.37 | 0.22 |
| 2009 | 231,437 | 1 | 8 | 62 | 291 | 1 | 1 | 0.13 | 0.13 |
| 2010 | 201,585 | 2 | 66 | 55 | 113 | 2 | 15 | 0.10 | 0.06 |
| 2011 | 259,964 | 8 | 62 | 113 | 633 | 10 | 16 | 0.27 | 0.25 |
| 2012 | 203,510 | 24 | 184 | 136 | 319 | 3 | 11 | 0.25 | 0.16 |
| 2013 | 207,859 | 100 | 116 | 116 | 246 | 0 | 0 | 0.17 | 0.12 |
| 2014 | 221,099 | 128 | 140 | 166 | 316 | 2 | 2 | 0.21 | 0.14 |
| 2015 | 199,686 | 8 | 39 | 113 | 126 | 0 | 0 | 0.08 | 0.06 |
| 2016 | 209,031 | 9 | 29 | 14 | 21 | 0 | 0 | 0.02 | 0.01 |
| 2017 | 144,219 | 2 | 2 | 15 | 15 | --- | --- | 0.01 | 0.01 |
| 2018 | 192,521 | 3 | 3 | --- | --- | --- | --- | 0.00 | 0.00 |
| Mean | | | | | | | | 0.23^b | 0.18^b |
| Geometric Mean | | | | | | | | 0.15^b | 0.12^b |

^a Expanded numbers are calculated from the proportion of each known age salmon recovered in the river and from broodstock collections in relation to the total estimated return to the Tucannon River. Expansions do not include down river harvest or Tucannon River fish straying to other systems.

^b The 2017 and 2018 SARs are not included in the mean.

As previously stated, overall survival of hatchery salmon to return as adults was higher than for naturally reared fish because of the early-life survival advantage (Table 24). With the exception of eleven brood years (34%), naturally produced fish have been below the replacement level (Figure 12; Table 27). Based on adult returns from the 1985-2017 broods, naturally reared salmon produced only 0.63 adults for every spawner, while hatchery reared fish produced 1.81 adults (based on geometric means).




Figure 12. Return per spawner (with replacement line) for the 1985-2017 brood years (2017 incomplete brood year).

Table 27. Progeny-to-parent survival estimates of Tucannon River spring Chinook salmon from 1985 through 2017 brood years (2017 brood year incomplete).

| Brood Year | Natural Salmon | | | Hatchery Salmon | | | Hatchery to Natural Advantage |
|-----------------------|--------------------|-------------------|-----------------|-----------------|-------------------|-----------------|-------------------------------|
| | Estimated Spawners | Number of Returns | Return/ Spawner | Number Spawned | Number of Returns | Return/ Spawner | |
| 1985 | 695 | 392 | 0.56 | 9 | 45 | 5.00 | 8.9 |
| 1986 | 440 | 467 | 1.06 | 91 | 319 | 3.51 | 3.3 |
| 1987 | 407 | 228 | 0.56 | 83 | 178 | 2.14 | 3.8 |
| 1988 | 257 | 502 | 1.95 | 87 | 385 | 4.43 | 2.3 |
| 1989 | 276 | 153 | 0.55 | 122 | 209 | 1.71 | 3.1 |
| 1990 | 572 | 94 | 0.16 | 78 | 28 | 0.36 | 2.2 |
| 1991 | 291 | 7 | 0.02 | 72 | 25 | 0.35 | 14.4 |
| 1992 | 476 | 161 | 0.34 | 83 | 76 | 0.92 | 2.7 |
| 1993 | 397 | 177 | 0.45 | 91 | 138 | 1.52 | 3.4 |
| 1994 | 97 | 12 | 0.12 | 69 | 32 | 0.46 | 3.7 |
| 1995 | 27 | 6 | 0.22 | 39 | 177 | 4.54 | 20.4 |
| 1996 | 152 | 69 | 0.45 | 74 | 265 | 3.58 | 7.9 |
| 1997 | 105 | 791 | 7.53 | 89 | 176 | 1.98 | 0.3 |
| 1998 | 60 | 388 | 6.47 | 85 | 793 | 9.33 | 1.4 |
| 1999 | 160 | 141 | 0.88 | 122 | 33 | 0.27 | 0.3 |
| 2000 | 201 | 448 | 2.23 | 73 | 157 | 2.15 | 1.0 |
| 2001 | 766 | 257 | 0.34 | 104 | 127 | 1.22 | 3.6 |
| 2002 | 568 | 212 | 0.37 | 93 | 121 | 1.30 | 3.5 |
| 2003 | 329 | 173 | 0.53 | 75 | 71 | 0.95 | 1.8 |
| 2004 | 346 | 399 | 1.15 | 88 | 120 | 1.36 | 1.2 |
| 2005 | 264 | 739 | 2.80 | 95 | 690 | 7.26 | 2.6 |
| 2006 | 202 | 1,720 | 8.51 | 88 | 1,122 | 12.75 | 1.5 |
| 2007 | 211 | 610 | 2.89 | 82 | 261 | 3.18 | 1.1 |
| 2008 | 796 | 884 | 1.11 | 114 | 643 | 5.64 | 5.1 |
| 2009 | 1191 | 619 | 0.52 | 173 | 300 | 1.73 | 3.3 |
| 2010 | 938 | 938 | 1.00 | 161 | 194 | 1.20 | 1.2 |
| 2011 | 849 | 727 | 0.86 | 166 | 711 | 4.28 | 5.0 |
| 2012 | 335 | 213 | 0.64 | 164 | 514 | 3.13 | 4.9 |
| 2013 | 170 | 69 | 0.41 | 149 | 362 | 2.43 | 6.0 |
| 2014 | 294 | 89 | 0.30 | 126 | 458 | 3.63 | 12.0 |
| 2015 | 523 | 45 | 0.09 | 126 | 165 | 1.31 | 15.2 |
| 2016 | 340 | 57 | 0.17 | 118 | 50 | 0.42 | 2.5 |
| 2017 | 249 | 98 | 0.39 | 99 | 17 | 0.17 | 0.4 |
| Mean | | | 1.38 | | | 2.86 | 4.5 |
| Geometric Mean | | | 0.63 | | | 1.81 | 2.9 |

Beginning with the 2006 brood year, the annual smolt goal was increased from 132,000 to 225,000 to help offset for the higher mortality of hatchery-origin fish after they leave the hatchery. This should increase adult salmon returns back to the Tucannon River. However, based on current hatchery SARs the increase in production would still not produce enough adult returns to reach the LSRCP mitigation goal. Hatchery production changes that result in increased survival/return numbers may result in a Proportionate Natural Influence (PNI) of less than 0.5. This level is generally not considered acceptable for supplementation programs. Historically the PNI for the Tucannon Spring Chinook Program has generally been above 0.5 (Appendix I).

The long-term restoration goal for the State of Washington is to provide a total annual return of between 2,400-3,400 hatchery and natural origin spring Chinook salmon back to the Tucannon River (SRSRB 2006) that should include at least 750 natural origin fish over a 10-year geometric mean (population viability threshold) (ICTRT 2008). Natural origin returns had been increasing, but decreased during recent years (2016-2021), likely due primarily to poor ocean conditions (Figure 13).




Figure 13. Tucannon River spring Chinook natural origin returns with the moving ten-year geometric mean (black line) for the 1985-2021 run years.

Fishery Contribution and Out-of-Basin Straying

The original goal of the LSRCP supplementation program was to enhance returns of salmon to the Tucannon River by providing 1,152 adult hatchery origin fish (the number estimated to have been lost to the project area due to the construction and operation of the Lower Snake River hydropower system) to the river from hatchery-reared smolt releases. Such an increase would allow for limited harvest and increased spawning. However, hatchery adult returns have always been below the mitigation goal (Figure 14). Based on CWT recoveries reported to the Regional Mark Information System (RMIS) database (Appendix J), sport and commercial harvest combined accounted for an average of less than 6% of the adult hatchery fish recovered for the 1985-1996 brood years. Increased fishery impacts occurred for the 1997 through 1999 broods when the states implemented mark-selective fisheries in the lower Columbia River (fishery harvest comprised an average of 19% for hatchery fish recoveries). As such, the WDFW subsequently stopped adipose fin clipping spring Chinook hatchery production from the Tucannon River (Gallinat et al. 2001) to lessen non-tribal fishery impacts from the Columbia River, and newly implemented Snake River fisheries. This change in marking has resulted in lower sport fishery impacts. Based on CWT recoveries for the 2000-2017 brood years, harvest (primarily commercial) has accounted for only 5.4% of the hatchery adult CWT recoveries (Appendix J).

Out-of-basin stray rates of Tucannon River spring Chinook have generally been low (Appendix J), with an average of 1.0% of the adult hatchery fish straying to other river systems/hatcheries for brood years 1985-2017 (range 0-20%). Recent (2005-2017 BYs) locations that Tucannon River spring Chinook have strayed are listed in Table 28.




Figure 14. Total escapement for Tucannon River spring Chinook salmon for the 1985-2021 run years.

Table 28. Summary of Tucannon River spring Chinook recovered outside of the Tucannon River and represent possible strays to other areas (2005-2017 brood years).

| Brood Year | CWT Codes | Recovery Location | Recovery Date | Number of CWT Recovered/Expanded |
|---|------------------------|--|----------------------|----------------------------------|
| 2005 | 633477, 633599 | None | N/A | 0/0 |
| 2006 | 634093, 634094, 634194 | Powell Rack, Lochsa River | 08/27/09 | 1/1 |
| 2007 | 634687, 634688 | None | N/A | 0/0 |
| 2008 | 635174, 635175 | None | N/A | 0/0 |
| 2009 | 635565, 635566 | Lower Granite Dam Trap | 10/17/13 | 1/1 |
| 2010 | 636075, 636076 | None | N/A | 0/0 |
| 2011 | 636441, 636442 | Lower Granite Dam Trap | 09/24/14 | 1/1 |
| 2012 | 636585, 636586 | Lewis River Hatchery SF Walla Walla River | 08/31/16 09/13/16 | 1/1 1/1 |
| 2013 | 636742, 636743 | None | N/A | 0/0 |
| 2014 | 636884 | None | N/A | 0/0 |
| 2015 | 637039 | Three Mile Dam, Umatilla River | 09/04/18 | 1/1 |
| 2016 | 637201 | None | N/A | 0/0 |
| 2017 | 637396 | None | N/A | 0/0 |
| Totals | | | | 6/6 |
| Total recovery of Tucannon fish from all locations | | | | 1,968/5,931 |
| Percent stray rate (recovered or expanded) | | | | 0.30%/0.10% |

Adjusted Hatchery SAS

Using CWT recoveries from the RMIS database, we adjusted Tucannon River spring Chinook hatchery smolt-to-adult survival (SAS) to include all known recoveries both from within and outside the Tucannon River. Increased fishing mortality resulted in higher adjusted SAS for the 1997, 1998, and 2006 brood years. With minor exceptions (1997 and 2006 brood years), even after adjustment, hatchery SAS rates were still below the original LSRCP survival assumption of 0.87% (Table 29).

Table 29. Hatchery SAS adjusted for recoveries from outside the Tucannon River subbasin as reported in the RMIS database, 1985-2016 brood years. (Data downloaded from RMIS database on 12/02/21).

| Brood Year | Estimated Number of Smolts | Expanded Return to Tucannon | Expanded Other Returns ^a | Grand Total of CWT Hatchery Origin Recoveries | Original Hatchery SAR (%) | Adjusted Hatchery SAS (%) |
|-----------------------|----------------------------|-----------------------------|-------------------------------------|---|---------------------------|---------------------------|
| 1985 | 12,922 | 45 | 1 | 46 | 0.35 | 0.36 |
| 1986 | 152,725 | 319 | 15 | 334 | 0.21 | 0.22 |
| 1987 | 152,165 | 178 | 2 | 180 | 0.12 | 0.12 |
| 1988 | 145,146 | 385 | 25 | 410 | 0.27 | 0.28 |
| 1989 | 99,057 | 209 | 12 | 221 | 0.21 | 0.22 |
| 1990 | 85,737 | 28 | 0 | 28 | 0.03 | 0.03 |
| 1991 | 74,064 | 25 | 4 | 29 | 0.03 | 0.04 |
| 1992 | 87,752 | 76 | 17 | 93 | 0.09 | 0.11 |
| 1993 | 138,848 | 138 | 11 | 149 | 0.10 | 0.11 |
| 1994 | 130,069 | 32 | 0 | 32 | 0.02 | 0.02 |
| 1995 | 62,144 | 177 | 2 | 179 | 0.28 | 0.29 |
| 1996 | 76,219 | 265 | 4 | 269 | 0.35 | 0.35 |
| 1997 | 24,186 | 176 | 41 | 217 | 0.73 | 0.90 |
| 1998 | 127,939 | 793 | 216 | 1,009 | 0.62 | 0.79 |
| 1999 | 97,600 | 33 | 3 | 36 | 0.03 | 0.04 |
| 2000 | 102,099 | 157 | 1 | 158 | 0.15 | 0.15 |
| 2001 | 146,922 | 127 | 5 | 132 | 0.09 | 0.09 |
| 2002 | 123,586 | 121 | 0 | 121 | 0.10 | 0.10 |
| 2003 | 71,154 | 71 | 0 | 71 | 0.10 | 0.10 |
| 2004 | 67,542 | 120 | 1 | 121 | 0.18 | 0.18 |
| 2005 | 149,466 | 690 | 2 | 692 | 0.46 | 0.46 |
| 2006 | 106,530 | 1,122 | 36 | 1,158 | 1.05 | 1.09 |
| 2007 | 114,681 | 261 | 5 | 266 | 0.23 | 0.23 |
| 2008 | 172,897 | 643 | 4 | 647 | 0.37 | 0.37 |
| 2009 | 231,437 | 300 | 7 | 307 | 0.13 | 0.13 |
| 2010 | 201,585 | 194 | 1 | 195 | 0.10 | 0.10 |
| 2011 | 259,964 | 711 | 24 | 735 | 0.27 | 0.28 |
| 2012 | 203,510 | 514 | 3 | 517 | 0.25 | 0.25 |
| 2013 | 207,859 | 362 | 11 | 373 | 0.17 | 0.18 |
| 2014 | 221,099 | 458 | 2 | 460 | 0.21 | 0.21 |
| 2015 | 199,686 | 165 | 1 | 166 | 0.08 | 0.08 |
| 2016 | 209,031 | 50 | 0 | 50 | 0.02 | 0.02 |
| Mean | | | | | 0.23 | 0.25 |
| Geometric Mean | | | | | 0.15 | 0.16 |

^a Includes expanded RMIS CWT recoveries from sources outside the Tucannon River Subbasin (i.e., sport and commercial fisheries, Tucannon strays in other river systems, etc.).

Alternative Rearing and Release Strategies

Because of the continued low adult returns back to the Tucannon River due to adverse environmental conditions (e.g., poor ocean conditions, drought, floods, habitat, etc.) and resulting hatchery production that is below program goals, managers are currently looking at three alternative hatchery rearing and release strategies in an effort to increase production and improve survival.

Captive Broodstock

If funding can be obtained, a new captive broodstock program will be conducted similar to the previous Tucannon River spring Chinook Captive Broodstock Program (see Gallinat et al. 2009) to provide for a safety-net to reduce the risk of catastrophic loss of the population. The captive broodstock program is expected to operate concurrently while experimental releases are attempted in an effort to improve conventional hatchery supplementation survival.

The goal will be to collect 290,000 eggs/year from captive females when three complete age classes (ages 3 to 5) are spawned concurrently. These eggs are expected to produce 150,000 yearling smolts for release into the Tucannon River at full production. Excess production above this amount could be released into Asotin Creek which is part of the same Ecologically Significant Unit (ESU) as the Tucannon River population and is considered to be functionally extirpated.

Rearing tanks for the initial rearing phase of the captive broodstock program were purchased in 2021, but manufacturing delays prevented us from utilizing them for the BY21 fish. The BY21 fish that were initially collected to start the captive broodstock program were returned to the conventional supplementation program. The tanks will be installed in 2022, and we anticipate using them for BY22 fish as a safety net to start the captive broodstock program. Whether or not they are reared to maturity will depend on available funding.

Releases below Bonneville Dam

If approved, another strategy employed to improve adult performance would be to acclimate and release Tucannon River hatchery spring Chinook smolts at a hatchery below Bonneville Dam (Kalama Falls Fish Hatchery - KFH) and transport the returning adults back to LFH for spawning or adult outplanting. A fish health inspection will be conducted prior to transferring pre-smolts to KFH. Juvenile fish will be transported from LFH in late October for 5 to 6 months of acclimation on Kalama River water to maximize imprinting. Fish taken to KFH will be

tagged with CWT and marked with a left maxillary clip (No AD clip) to visually distinguish them from other stocks in the lower Columbia River. Smolts will be volitionally released in late March/early April. The size of the program would be capped at no greater than 50% of Tucannon River hatchery production, or up to 100,000 smolts, whichever is less, depending on KFH capacity. The initial numbers however will range between 30,000 to 50,000 smolts.

Returning adults from this strategy will be used to:

- 1) Fulfill shortfalls to supplementation broodstock collection goals to produce 225,000 smolts.
- 2) Outplant into the Tucannon River to supplement natural production.

It is expected that ~20% of the adults will not fully recruit to the KFH adult trap but will remain below the trap and spawn in the Kalama River. In addition, it is estimated that 1-2% of the adults might be able to pass above the falls and spawn in the upper Kalama River basin. Therefore, this strategy is contingent on risk assessment to lower Columbia River populations and approval by NOAA Fisheries. A goal of this program is to keep the number of fish that do not recruit to the trap at less than 100 fish. This level is thought to have a minimal impact to the Kalama River population. Discussions with NOAA Fisheries regarding this strategy are on-going.

Experimental Release Strategies

Based on PIT tag analyses conducted by the Fish Passage Center from 2005-2017, the SAR survival of Tucannon Hatchery spring Chinook from Lower Monumental to Bonneville Dam has been lower than the Lower Granite to Bonneville Dam SARs for up-river hatchery stocks, even though the reach the Tucannon hatchery fish migrate through is shorter. It has been hypothesized that the up-river stocks may have a survival advantage due to additional opportunities for barge transportation. While some studies have shown barging has affected homing abilities for both Chinook and steelhead (Quinn 1993; Keefer et al. 2008b; Keefer and Caudill 2014; Bond et al. 2017), evaluating the effect of barge transportation on the SARs of Tucannon River fish has not been possible with the available data. A more recent PIT tag analyses was completed by the Fish Passage Center (July 28, 2020) comparing SAR survival of Tucannon River hatchery spring Chinook with up-river stocks, but this time as a direct comparison of SAR survivals from Lower Monumental to Bonneville Dam for all stocks. Results showed that while Tucannon River hatchery spring Chinook generally track survival of other groups, they consistently had lower juvenile survival.

Historically, the default action for PIT tagged fish that are detected at transportation facilities has been to return them to the river. Beginning with the 2015 migration year, PIT tagged Tucannon

River hatchery spring Chinook have been included in the Comparative Survival Study (CSS) whereby a portion of the tagged fish are returned to the river and a portion are barged. However, the effects of transportation on SARs of Tucannon hatchery fish to date has not been possible based on the low numbers of PIT tagged fish (15,000/year), and poor collection efficiency of PIT tags at Lower Monumental Dam. Power analysis performed by the Fish Passage Center has determined that the number of PIT tagged fish needed to find a significant difference in survival based on historical rates is approximately 15,000 fish/group.

Survival within the Tucannon River itself from the point of hatchery release to detection at Lower Monumental Dam shows potential for improvement. Survival to the Tucannon River instream PIT arrays and Lower Monumental Dam were obtained from the University of Washington (Columbia River Data Access in Real Time [DART]; www.cbr.washington.edu/). Based on DART PIT tag survival of spring Chinook released from Curl Lake AP, average survival to Lower Monumental Dam is less than 60% (Figure 15).




Figure 15. Tucannon River hatchery-origin spring Chinook survival with 95% confidence interval from release at Curl Lake to the Tucannon River instream antenna arrays (TFH = Tucannon Fish Hatchery; UTR = Upper Tucannon River; MTR = Middle Tucannon River; LTR = Lower Tucannon River) and Lower Monumental Dam (LMO) for the 2012 to 2020 migration years.

When sufficient hatchery production is available (i.e., full production of smolts for in-basin release), we will examine three different release strategies (Direct Stream Release at TFH, Direct Stream Release near the Mouth, and Barge Transportation) by PIT tagging a minimum of 15,000 fish per group in an attempt to determine if significant improvements in survival rates can be achieved for Tucannon River spring Chinook hatchery fish. The study will be conducted for a minimum of three BYs with PIT tag detections from returning adults used to determine significant differences among the release groups. Fish used for the study will be transferred from LFH to TFH in October and reared in the same raceway (A-Pond). This is to ensure that all groups will be treated similarly over the fall/winter months prior to PIT tagging and allow for imprinting to the Tucannon River. Imprinting is critical for the successful completion of the spawning migration (Dittman and Quinn 1996), and the parr-to-smolt transformation has been identified as an important period for olfactory imprinting (Dittman et al. 1996). The shift to releases lower in the river for this study could have consequences (survival, adult trapping, and spawning distributions) that are not fully appreciated at this time. Management actions to account for some of these (hauling returning adults upstream, additional trapping locations for broodstock collection/hauling, etc.) may have to be implemented.

Direct Stream Release at TFH Group (Surrogate Control Group)

Fish from this group will be PIT tagged and put back into the A-Pond and released to the outlet channel and Tucannon River. The release will begin in early to mid-April to coincide with the release timing of the other two groups. Historically, fish were released from Curl Lake AP so this group will not represent a true control group.⁴ This group will also contain the remaining hatchery production (CWT only fish) since survival and returns of fish released near the mouth of the Tucannon and fish that are barged are unknown at this time. The PIT tagged fish from this group will be removed from monitor mode at the dams and the CSS (to compare in-river vs. transported survival).

Direct Stream Release near the Mouth Group

Fish from this group will be PIT tagged and placed into a separate rearing vessel for final rearing prior to release (~1 month). These fish will be released below the smolt trap near the mouth of the Tucannon River (eliminates majority of Tucannon in-river mortality) and we will attempt to release them to coincide with the arrival of the surrogate control group in the lower river. There is a concern that this group might stray past the Tucannon River and spawning distribution could

⁴ The use of Curl Lake Acclimation Pond has been an important part of the hatchery program in addressing survival and spawning distribution concerns for hatchery fish over the years. Options to re-use Curl Lake Acclimation Pond (depending upon study results) should be considered in the future.

shift lower in the river. For example, fall Chinook released at the mouth of the Umatilla River failed to imprint and showed poor homing instinct, presumably due to their short time spent in the river (Hayes and Carmichael 2002). Fish from this group will also be removed from monitor mode at the dams and the CSS.

Barge Transportation Group

Fish from this group will be PIT tagged and placed into a separate rearing vessel for final rearing prior to release (~ 1 month). Arrangements will be made with the Corps of Engineers to have a barge available at LFH between 24-30 April (barging at the Snake River Dams begins on 24 April). On the day that the barge is scheduled to arrive at LFH, the fish will be loaded onto a transport truck, hauled to LFH, and put on the barge the same day. This group also has a high chance on straying and shifting spawning distribution lower in the river.

Due to low production numbers for BY20, the decision was made to directly compare the TFH release group (surrogate control group) and the release near the mouth, since these two groups would provide immediate juvenile survival results based on PIT tag detections at the dams. A total of 20,000 PIT tags were used for each group. The BY21 production numbers are higher and should allow all three groups to be compared beginning with 2023 releases. Implementation of this study will be dependent upon the availability of future production and the health status of the hatchery fish.

Conclusions and Recommendations

Washington's LSRCP hatchery spring Chinook salmon program in the Tucannon River has historically failed to return adequate numbers of adults to meet the mitigation goal for Washington (1,152). This has occurred because SARs of hatchery origin fish have been consistently lower than what was originally assumed under the LSRCP program development. However, because of the advantage in survival during early life history stages for fish in the hatchery, the progeny-to-parent ratio for hatchery-produced fish has generally been above replacement and therefore may have sustained the overall Tucannon spring Chinook population during years when the population was at critically low levels. For a while, we had seen a significant rebound of natural origin fish and we came close to reaching the within river hatchery (LSRCP) goal of 1,152 fish in 2009 and 2010. Recent returns have been much lower, which is believed to be the result of poor ocean conditions. System survivals (in-river, migration corridor, and ocean) must increase in the near future for the hatchery program to succeed, the natural run to persist over the short-term, and the natural population to increase to a level where it can be sustainable over the long-term.

Until that time, the evaluation program will continue to document and study life history survivals, straying, carrying capacity, genotypic and phenotypic traits, and examine procedures within the hatchery that can be changed to improve the hatchery program and the natural population. Based on our previous studies and current data we recommend the following:

1. We continue to see annual differences in phenotypic characteristics of returning salmon (i.e., hatchery fish are generally younger and less fecund than natural origin fish), yet other traits such as run and spawn time are little changed over the program's history. Further, genetic analysis to date has detected little change in the natural population that may have resulted from hatchery actions.

Recommendation: Continue to collect as many carcasses as possible for the most accurate age composition data. Collect biological data (length, run timing, spawn timing, fecundity estimates, DNA samples, smolt trapping, and life stage survival) to document the effects (positive or negative) that the hatchery program may have on the natural population.

2. We have documented that hatchery juvenile (egg-parr-smolt) survival rates are considerably higher than naturally reared salmon, and hatchery smolt-to-adult return rates are much lower than their natural origin counterpart. The population is approaching critical minimum abundance thresholds that may require intervention or run the risk of extirpation. We have been considerably below hatchery production goals and ocean conditions have been poor for salmon survival. We need to identify and address the factors that limit hatchery SARs in order to meet the mitigation goals and for natural production to meet recovery goals.

Recommendation: Managers are currently discussing alternative rearing and release strategies in an attempt to increase hatchery fish survival and preserve this stock (e.g., releasing smolts below Bonneville Dam, barge transportation, captive broodstock, etc.). Funding for a captive broodstock program should be sought by WDFW to provide a safety-net for this population due to uncertain environmental conditions and the uncertainty of adult returns from experimental release strategies. As long as adequate numbers of smolts can be produced, WDFW will continue comparing alternative release strategies to determine if survival rates can be improved to provide greater adult returns. Continue to utilize fish carcasses from hatchery operations for stream nutrient enrichment to improve overall productivity and survival of natural-origin Tucannon River spring Chinook.

3. Subbasin and recovery planning for ESA listed species in the Tucannon River have identified factors limiting the spring Chinook population and strategies to recover the population.

Recommendation: Assist population conservation efforts by updating recent carrying capacity/density and straying effects, and productivity estimates of the Tucannon River so that hatchery stocking is appropriate, and hatchery and natural performance is measured against future basin capacity after habitat improvements.

4. Previously, we have documented higher in-river pre-spawn mortality than what was observed historically. The mechanism for this higher loss is thought to be due to a combination of drought years with higher water temperatures and pathogen load. However, the high loss has prompted drastic action within the program, whereby all, or the majority of the returns to the TFH trap between 2015 to 2019 have been collected and held for adult outplanting. Results from the first year (2015) of adult outplants appeared successful, with > 90% of the fish spawning, contrasted to 30% survival of fish left in the river. From 2016-2018, a range of 55-72% of outplanted fish successfully spawned.

Recommendation: Continue to monitor in-river pre-spawn mortality. Continue intensive monitoring of adult outplants, when that strategy is employed, to determine spawning success. Weigh all pertinent information (pre-spawn mortality rates, outplant success, predicted run sizes, risk of holding all fish at one facility, etc.) and inform co-managers and NOAA Fisheries on future adult outplants. An agreed upon population threshold trigger is needed to determine whether to pass fish at the adult trap or hold fish at LFH for outplanting. A trigger has been suggested by M&E staff (allow outplanting below an estimated return of 400 adults) but has yet to be agreed upon by the co-managers.

5. Based on annual redd densities and historical spring Chinook radio tag data, and PIT tag data from the TFH PIT tag array, the Tucannon Fish Hatchery weir/trap has been an impediment to upstream passage of spring Chinook to the better spawning and rearing habitat upstream of the trap. Numerous options to improve attraction into the ladder/trap have been discussed with some recently implemented. A new trap fyke was installed in 2020, which based on PIT tag detections of steelhead, spring Chinook, and bull trout is considerably better at retaining fish in the trap, either for broodstock collections, or for passing fish upstream with minimal delays.

Recommendation: Continue to monitor changes made to the ladder/trap to see if they improve passage conversion and reduce migration delay for all fish species. If improvements are not seen, and passage delays are still unacceptable, seek funding and engineering expertise to modify the design and/or operation of the weir/trap structure.

Literature Cited

- Bond, M. H., P. A. H. Westley, A. H. Dittman, D. Holecek, T. Marsh, and T. P. Quinn. 2017. Combined effects of barge transportation, river environment, and rearing location on straying and migration of adult Snake River fall-run Chinook Salmon. *Transactions of the American Fisheries Society* 146: 60-73.
- Bugert, R., P. LaRiviere, D. Marbach, S. Martin, L. Ross, and D. Geist. 1990. Lower Snake River Compensation Plan Salmon Hatchery Evaluation Program 1989 Annual Report to U.S. Fish and Wildlife Service, AFF 1/LSR-90-08, Cooperative Agreement 14-16-0001-89525. Washington Department of Fisheries, Olympia, Washington.
- Bumgarner, J., L. Ross, and M. Varney. 2000. Lower Snake River Compensation Plan Tucannon River Spring Chinook Salmon Hatchery Evaluation Program 1998 and 1999 Annual Reports to U.S. Fish and Wildlife Service, Cooperative Agreements 1448-14110-98-J057 and CA-14110-9-J070. Washington Department of Fish and Wildlife, Olympia, Washington. Report # FPA00-17.
- Busack, C., and C. M. Knudsen. 2007. Using factorial mating designs to increase the effective number of breeders in fish hatcheries. *Aquaculture* 273: 24-32.
- Cheng, Y. W., and M. P. Gallinat. 2004. Statistical analysis of the relationship among environmental variables, inter-annual variability and smolt trap efficiency of salmonids in the Tucannon River. *Fisheries Research* 70: 229-238.
- Dittman, A. H., and T. P. Quinn. 1996. Homing in Pacific salmon: mechanisms and ecological basis. *Journal of Experimental Biology* 199: 83-91.
- Dittman, A. H., T. P. Quinn, and G. A. Nevitt. 1996. Timing of imprinting to natural and artificial odors by coho salmon (*Oncorhynchus kisutch*). *Canadian Journal of Fisheries and Aquatic Sciences* 53: 434-442.
- Gallinat, M. P. and W-Y Chang. 2013. Phenotypic comparisons among natural-origin, hatchery-origin, and captive-reared female spring Chinook salmon from the Tucannon River, Washington. *North American Journal of Aquaculture* 75 (4): 572-581.
- Gallinat, M. P., J. D. Bumgarner, D. Maxey, S. Roberts, R. Rogers, L. Ross, and M. A. Varney. 2009. Tucannon River spring Chinook Salmon captive broodstock program: Final project completion report (October 1, 1999 – September 30, 2009). Prepared for the U.S. Department of Energy, Bonneville Power Administration. Project Number 200-019-00. 49 pp.
- Gallinat, M. P., J. D. Bumgarner, L. Ross, and M. Varney. 2001. Tucannon River Spring Chinook Salmon Hatchery Evaluation Program 2000 Annual Report to U.S. Fish and Wildlife Service, Cooperative Agreement 1411-09-J070. Washington Department of Fish and Wildlife, Olympia, Washington. Report # FPA01-05.

- Gallinat, M. P., and L. A. Ross. 2014. Tucannon River Spring Chinook Salmon Hatchery Evaluation Program 2013 Annual Report to U.S. Fish and Wildlife Service, Cooperative Agreement F13AC00096. Washington Department of Fish and Wildlife, Olympia, Washington. Report #FPA14-05. 104 p.
- Gallinat, M. P., and L. A. Ross. 2015. Tucannon River Spring Chinook Salmon Hatchery Evaluation Program 2014 Annual Report to U.S. Fish and Wildlife Service, Cooperative Agreement F14AC00010. Washington Department of Fish and Wildlife, Olympia, Washington. Report #FPA15-04. 108 p.
- Gallinat, M. P., and D. E. Kiefel. 2021. Tucannon River Spring Chinook Salmon Hatchery Evaluation Program 2020 Annual Report to U.S. Fish and Wildlife Service, Cooperative Agreement F16AC00033. Washington Department of Fish and Wildlife, Olympia, Washington. Report #FPA21-03. 96 p.
- Groot, C., and L. Margolis. 1991. Pacific salmon life histories. UBC Press. Vancouver, B.C. 564 p.
- Hayes, M. C., and R. W. Carmichael. 2002. Salmon restoration in the Umatilla River: A study of straying and risk containment. *Fisheries* 27 (10): 10-19.
- ICTRT (Interior Columbia Technical Recovery Team). 2008. Current status assessments. U.S. Dept. Commer., NOAA, National Marine Fisheries Service, Northwest Region, Portland, Ore.
- Keefer, M. L., C. C. Caudill, C. A. Peery, and C. T. Boggs. 2008a. Non-direct homing behaviours by adult Chinook salmon in a large, multi-stock river system. *Journal of Fish Biology* 72: 27-44.
- Keefer, M. L., C. C. Caudill, C. A. Peery, and S. R. Lee. 2008b. Transporting juvenile salmonids around dams impairs adult migration. *Ecological Applications* 18 (8): 1888-1900.
- Keefer, M. L., and C. C. Caudill. 2014. Homing and straying by anadromous salmonids: a review of mechanisms and rates. *Reviews in Fish Biology and Fisheries* 24: 333-368.
- Kohler, A. E., T. N. Pearsons, J. S. Zendt, M. G. Mesa, C. L. Johnson, and P. J. Connolly. 2012. Nutrient enrichment with salmon carcass analogs in the Columbia River Basin, USA: a stream food web analysis. *Transactions of the American Fisheries Society* 141: 802-824.
- Meekin, T. K., 1967. Report on the 1966 Wells Dam Chinook tagging study. Report to Douglas County PUD, Contract 001-01-022-4201. Washington Department of Fisheries, Olympia, WA. 41 p. (Available from Douglas County PUD, 1151 Valley Mall Parkway, East Wenatchee, WA 98801.)
- Murdoch, A. R., T. N. Pearsons, and T. W. Maitland. 2009. The number of redds constructed per female spring Chinook salmon in the Wenatchee River basin. *North American Journal of*

- Fisheries Management 29: 441-446.
- Murdoch, A. R., T. N. Pearsons, T. W. Maitland. 2010. Estimating the spawning escapement of hatchery- and natural-origin spring Chinook salmon using redd and carcass data. *North American Journal of Fisheries Management* 30: 361-375.
- Nehlsen, W., J. E. Williams, and J. A. Lichatowich. 1991. Pacific salmon at the crossroads: stocks at risk from California, Oregon, Idaho, and Washington. *Fisheries* 16 (3): 4-21.
- PTAGIS website 2020. Information of the initiations of barge transportation at Lower Monumental Dam.
- Quinn, T. P. 1993. A review of homing and straying of wild and hatchery-produced salmon. *Fisheries Research* 18: 29-44.
- Scheuerell, M. D., P. S. Levin, R. W. Zabel, J. G. Williams, and B. L. Sanderson. 2005. A new perspective of marine-derived nutrients to threatened stocks of Pacific salmon (*Oncorhynchus* spp.). *Canadian Journal of Fisheries and Aquatic Sciences* 62: 961-964.
- Seiler, D., L. Kishimoto, and S. Neuhauser. 1999. 1998 Skagit River wild 0+ Chinook production evaluation. Washington Department of Fish and Wildlife. Olympia, Washington. 73 pp.
- Snake River Salmon Recovery Board (SRSRB). 2006. Technical Document Snake River Salmon Recovery Plan for S.E. Washington. Prepared for the Washington Governor's Salmon Recovery Office. 408 pages, plus appendices.
- Steinhorst, K., Y. Wu, B. Dennis, and P. Kline. 2004. Confidence intervals for fish outmigration estimates using stratified trap efficiency methods. *Journal of Agricultural, Biological, and Environmental Statistics* 9 (3): 284-299.
- Tucannon Subbasin Summary. 2001. L. Gephart and D. Nordheim, editors. Prepared for the Northwest Power Planning Council. Dayton, Washington.
- USACE (U.S. Army Corps of Engineers), 1975. Special Reports: Lower Snake River Fish and Wildlife Compensation Plan. Walla Walla, Washington.
- Washington Department of Fish and Wildlife (WDFW). 2017. Washington Department of Fish and Wildlife Salmonid Stock Inventory Database. WDFW, Olympia. Available: <http://wdfw.wa.gov/conservation/fisheries/sasi/> (May 2017).

**Appendix A: Annual Section 10 Permit #18024 Takes
for 2021, and NEOR/SEWA Terms and Conditions
Biological Opinion Reporting Requirements**

Appendix A. Table 1. Summary of permissible direct take and actual take (in parenthesis) of Snake River spring/summer Chinook salmon for RM&E activities associated with the Tucannon River spring Chinook salmon program not directly related to fish culture for the 2021 calendar year. NMFS must be notified within two days if the number handled, tagged, or killed are exceeded.

| Origin and Life Stage | Take activity | Capture method And location | Total number handled annually (0.5% handling mortalities) | Number of those handled that are marked/tagged annually (1% handling mortalities) | Total number killed or removed annually |
|--------------------------------|---|---|--|--|--|
| Natural-origin juveniles | Capture, handle, tag, tissue sample, and release live animal. | Trapping operations that include a screw trap, beach seines, cast nets, dip nets, and use of backpack electroshock equipment throughout the Tucannon River. | 18,000 (52) | 7,000 (52) | Up to 160 (0) |
| Hatchery-origin juveniles | Capture, handle, tag, tissue sample, and release live animal. | Trapping operations that include a screw trap, beach seines, cast nets, dip nets, and use of backpack electroshock equipment throughout the Tucannon River. | 35,000 (840) | 7,000 (0) | Up to 245 (5) |
| Natural-origin adults & jacks | Capture, handle, tag, tissue sample, and release live animal. | Adult and jack fall back at screw traps. | 5 (0) | 5 (0) (genetic fin-clip or operculum punch – release live.) | Up to 2 ^a (0) |
| Hatchery-origin adults & jacks | Capture, handle, tag, tissue sample, and release live animal. | Adult and jack fall back at screw traps. | 10 (0) | 10 (0) | Up to 2 ^a (0) |

^a In cases where total number killed is not likely to exceed one (1) mortality, NMFS rounds the total mortality up to two (2), so that operations are not halted completely at the first mortality.

Appendix A. Table 2. Summary of permissible direct take and actual take (in parenthesis) of listed Snake River spring/summer Chinook salmon for fish culture purposes for the Tucannon River Spring Chinook salmon program for the 2021 calendar year. NMFS must be notified within two days if the number handled, tagged, or killed are exceeded.

| Origin and Life Stage | Take activity | Capture method and location | Total number handled annually | Number of those handled that are marked/tagged annually (1% trap mortalities) | Total number killed or removed annually |
|---------------------------------|---|--|---|--|--|
| Natural-origin adults | Capture, handle, tag, tissue sample, remove for transport, holding, and outplanting in the Tucannon River, remove for use for broodstock, or release live animal (pass above weir). | Tucannon River adult weir and Lyons Ferry Hatchery ladder ^a | 2,000 (76) | Up to 1,824 ^b (passed live with fin-clip or operculum punch, PIT and/or radio tagged) (0 passed upstream) (0 outplanted upstream) | Up to 232 ^b broodstock and fish used for outplants (76 broodstock) Plus up to 19 adult trap mortalities (0) |
| Natural-origin jacks | Capture, handle, tag, tissue sample, remove for transport, holding, and outplanting in the Tucannon River, remove for use for broodstock, or release live animal (pass above weir). | Tucannon River adult weir and Lyons Ferry Hatchery ladder ^a | 200 (7) | Up to 200 (passed live with fin-clip or operculum punch, PIT and/or radio tagged) (0 passed upstream; 0 outplanted upstream) | Up to 9 broodstock. (7 collected but not used) Plus up to 2 trap mortalities. (0) |
| Hatchery-origin adults | Capture, handle, tag, tissue sample, remove for transport, holding, and outplanting in the Tucannon River, remove for use for broodstock, or release live animal (pass above weir). | Tucannon River adult weir and Lyons Ferry Hatchery ladder ^a | 1,400 ^b (up to 132 removed for broodstock based on sliding scale) (35) | Up to 1,400 ^b (passed live with fin-clip or operculum punch, PIT and/or radio tagged) (0 passed upstream) (0 outplanted upstream) | Up to 232 ^b broodstock and fish held for later outplanting. (22 broodstock) Up to 100% of total handled may be removed, killed, or transported as described in the HGMP (13 strays KO and 0 A.O. DIPs) |
| Hatchery-origin jacks | Capture, handle, tag, tissue sample, remove for transport, holding, and outplanting in the Tucannon River, use for broodstock, remove for adult management, or release live animal (pass above weir). | Tucannon River adult weir and Lyons Ferry Hatchery ladder ^a | 500 (22) | Up to 135 (more may be passed to mimic natural-origin jack proportions, with NMFS concurrence) (passed live with fin-clip or operculum punch) (0 passed upstream) (0 outplanted) | Up to 9 broodstock. (10 collected but not used.) Up to 100% of remainder may be removed, transported, or killed for jack management as described in the HGMP (12 strays KO and 0 A.O. DIP/PSM) |
| Hatchery-origin egg & juveniles | Capture, handle, tag, tissue sample, and release live animal (within hatchery sampling, and research use). | Tucannon Hatchery or Lyons Ferry Hatchery total | 280,125 (166,237 BY21) (Maximum eggs/juveniles on hand annually prior to any juvenile rearing loss) | 280,125 62,762 BY20 CWT 4,730 BY19 PIT tagged | Up to 55,125 total rearing mortality (10,380 BY20) (806 BY21) |
| Hatchery-origin juveniles | Capture, sample, kill (fish health examinations) | Tucannon Hatchery or Lyons Ferry Hatchery total | 170 (15) | 170 (0) | 170 (15) |

^a In years when returns to Tucannon Hatchery are low, adult Chinook arriving at Lyons Ferry Hatchery ladder that are identifiable as Tucannon River hatchery adults may be taken for broodstock.

^b The actual number taken annually will be subject to the sliding scale in the HGMP, in addition to fish that are collected, held, and used for adult outplants in the Tucannon, but may die while holding, or be used as part of the broodstock, and shall not exceed the totals of each origin identified there.

| Appendix A. Table 3. NOAA Terms and Conditions Biological Opinion reporting requirements for Tucannon River spring Chinook. | |
|---|--|
| Metric of Interest | Location within Report |
| Number and composition of broodstock, dates of collection, and number that die. | Appendix B. |
| Numbers, pounds, CV, dates, location, and tag/mark information of hatchery released fish, with precocial maturation rates. | Table 19; Appendix G. |
| Survival rates of Tucannon hatchery-origin fish life stages. | Tables 23 and 24. |
| Disease occurrence at Lyons Ferry Hatchery, Tucannon Hatchery, and Curl Lake AP. | Pages 9 and 31. |
| The number of returning hatchery and natural-origin adults and age structure. | Page 22; Table 11; Figure 5; Appendix C. |
| Distribution of hatchery and natural-origin spawners. | Table 6. |
| pHOS, pNOB, and PNI for the Tucannon River program. | Appendix I. |
| Survival rates of natural-origin fish. | Tables 22 and 24. |
| Smolt-to-adult survival rate (hatchery and natural-origin fish). | Tables 25 and 26. |
| The contribution of spring Chinook from this program into other populations (2005 to 2017 brood years). | Table 28. |
| The contribution of spring Chinook from other programs into the Tucannon River. | Page 27; Table 15; Appendix E. |
| Post release out-of-basin migration timing (median travel time and speed) of juvenile hatchery-origin fish to Lower Monumental Dam. | Table 21. |
| Mean length, coefficient of variation, number, and age of natural-origin juveniles. | Pages 32 to 34. |
| Any problems that may have arisen during hatchery activities. | A large number of strays entered the Tucannon River. We were able to remove all marked strays from the broodstock. |
| Any unforeseen effects on listed fish. | High proportion of strays were spawning in the Tucannon River. |

Appendix B: Spring Chinook Captured, Transported to Lyons Ferry Hatchery, or Returned to the River at the Tucannon Hatchery Trap in 2021

Appendix B. Spring Chinook salmon captured, transported to Lyons Ferry Hatchery, or returned to the river at the Tucannon Hatchery trap in 2021. (Trapping began in February; last day of trapping was September 30).

| Date | Captured in Trap | | Collected for Broodstock | | Passed Upstream | | Killed Outright ^a | |
|--------------------------------|------------------|-----------|--------------------------|-----------|-----------------|----------|------------------------------|-----------|
| | Natural | Hatchery | Natural | Hatchery | Natural | Hatchery | Natural | Hatchery |
| 5/24 | 2 | 1 | 2 | 1 | | | | |
| 5/27 | 4 | 2 | 4 | 1 | | | | 1 |
| 5/28 | | 1 | | 1 | | | | |
| 6/01 | 9 | 1 | 9 | 1 | | | | |
| 6/02 | 6 | 4 | 6 | 3 | | | | 1 |
| 6/03 | 2 | | 2 | | | | | |
| 6/04 | 3 | 3 | 3 | 1 | | | | 2 |
| 6/05 | 12 | 4 | 12 | 3 | | | | 1 |
| 6/07 | 1 | | 1 | | | | | |
| 6/11 | 7 | 5 | 7 | 3 | | | | 2 |
| 6/13 | 5 | 3 | 5 | 3 | | | | |
| 6/14 | 11 | 2 | 11 | 2 | | | | |
| 6/15 | 5 | 2 | 5 | | | | | 2 |
| 6/16 | 3 | 1 | 3 | | | | | 1 |
| 6/17 | 2 | | 2 | | | | | |
| 6/18 | 2 | 1 | 2 | | | | | 1 |
| 6/20 | 3 | 5 | 3 | 1 | | | | 4 |
| 6/21 | 1 | | 1 | | | | | |
| 6/22 | | 2 | | 2 | | | | |
| 6/24 | 1 | | 1 | | | | | |
| 6/25 | 2 | 4 | 2 | 3 | | | | 1 |
| 6/28 | 3 | 2 | 3 | 1 | | | | 1 |
| 6/29 | | 2 | | | | | | 2 |
| 6/30 | | 3 | | 2 | | | | 1 |
| 7/01 | 1 | 1 | 1 | | | | | 1 |
| 7/03 | | 1 | | | | | | 1 |
| 7/04 | | 1 | | | | | | 1 |
| 8/23 | 1 | 2 | 1 | | | | | 2 |
| 8/30 | 1 | | 1 | | | | | |
| Total | 87 | 53 | 87 | 28 | 0 | 0 | 0 | 25 |
| Final Total^b | 83 | 57 | 83 | 32 | 0 | 0 | 0 | 25 |

^a Fin clipped strays that were killed outright at the trap.

^b Corrected numbers after spawning.

**Appendix C: Age Composition by Brood Year for
Tucannon River Spring Chinook Salmon
(1985-2016 BYs)**

Appendix C. Age composition by brood year for natural and hatchery origin Tucannon River spring Chinook salmon (1985-2016 BYs). (Number at age is found in Tables 25 and 26).

| Brood Year | Natural origin | | | Hatchery origin | | |
|-------------------|-----------------------|----------------|----------------|------------------------|----------------|----------------|
| | % Age 3 | % Age 4 | % Age 5 | % Age 3 | % Age 4 | % Age 5 |
| 1985 | 4.85 | 65.05 | 30.10 | 42.22 | 57.78 | 0.00 |
| 1986 | 0.43 | 80.30 | 19.27 | 26.02 | 68.97 | 5.02 |
| 1987 | 0.00 | 73.25 | 26.75 | 10.67 | 81.46 | 7.87 |
| 1988 | 0.60 | 63.35 | 36.06 | 25.71 | 63.38 | 10.91 |
| 1989 | 7.84 | 75.16 | 16.99 | 6.22 | 85.65 | 8.13 |
| 1990 | 8.51 | 76.60 | 14.89 | 21.43 | 71.43 | 7.14 |
| 1991 | 0.00 | 71.43 | 28.57 | 20.00 | 80.00 | 0.00 |
| 1992 | 1.24 | 85.71 | 13.04 | 14.47 | 82.89 | 2.63 |
| 1993 | 1.13 | 56.50 | 42.37 | 10.87 | 77.54 | 11.59 |
| 1994 | 0.00 | 83.33 | 16.67 | 12.50 | 71.88 | 15.63 |
| 1995 | 0.00 | 16.67 | 83.33 | 9.04 | 88.70 | 2.26 |
| 1996 | 0.00 | 91.30 | 8.70 | 22.26 | 72.45 | 5.28 |
| 1997 | 1.77 | 87.86 | 10.37 | 7.39 | 92.61 | 0.00 |
| 1998 | 2.32 | 66.75 | 30.93 | 12.23 | 68.85 | 18.92 |
| 1999 | 6.38 | 87.94 | 5.67 | 33.33 | 57.58 | 9.09 |
| 2000 | 0.67 | 87.50 | 11.83 | 16.56 | 83.44 | 0.00 |
| 2001 | 0.00 | 95.72 | 4.28 | 14.96 | 84.25 | 0.79 |
| 2002 | 1.42 | 63.21 | 35.38 | 4.96 | 81.82 | 13.22 |
| 2003 | 4.05 | 66.47 | 29.48 | 2.82 | 91.55 | 5.63 |
| 2004 | 2.01 | 88.22 | 9.77 | 15.00 | 81.67 | 3.33 |
| 2005 | 17.73 | 80.51 | 1.76 | 42.17 | 57.83 | 0.00 |
| 2006 | 6.74 | 80.76 | 12.50 | 35.83 | 60.52 | 3.65 |
| 2007 | 6.72 | 74.75 | 18.52 | 28.35 | 65.52 | 6.13 |
| 2008 | 9.62 | 78.39 | 11.99 | 41.84 | 57.39 | 0.78 |
| 2009 | 1.13 | 89.50 | 9.37 | 2.67 | 97.00 | 0.33 |
| 2010 | 9.70 | 85.18 | 5.12 | 34.02 | 58.25 | 7.73 |
| 2011 | 5.64 | 85.14 | 9.22 | 8.64 | 88.16 | 3.20 |
| 2012 | 30.52 | 69.48 | 0.00 | 36.36 | 61.46 | 2.17 |
| 2013 | 11.59 | 86.96 | 1.45 | 32.23 | 67.77 | 0.00 |
| 2014 | 10.11 | 88.76 | 1.12 | 30.57 | 69.00 | 0.44 |
| 2015 | 0.00 | 93.33 | 6.67 | 23.64 | 76.36 | 0.00 |
| 2016 | 3.51 | 87.72 | 8.77 | 58.00 | 42.00 | 0.00 |
| Means | 5.91 | 79.92 | 14.17 | 24.84 | 70.42 | 4.74 |

Appendix D: Total Estimated Run-Size of Tucannon River Spring Chinook Salmon (1985-2021)

**Appendix D. Total estimated run-size of spring Chinook salmon to the Tucannon River, 1985-2021.
(Includes breakdown of conventional hatchery supplementation, captive brood progeny, and stray hatchery components).**

| Year | Natural Jacks | Natural Adults | Hatchery Jacks | Hatchery Adults | C.B. Jacks | C.B. Adults | Stray Jacks | Stray Adults | Total Natural | Total Hatchery | Total Run |
|------|------------------|-------------------|-------------------|--------------------|---------------|----------------|----------------|-----------------|------------------|-------------------|--------------|
| 1985 | --- | --- | --- | --- | --- | --- | --- | --- | 844 | 0 | 844 |
| 1986 | --- | --- | --- | --- | --- | --- | --- | --- | 636 | 0 | 636 |
| 1987 | --- | --- | --- | --- | --- | --- | --- | --- | 582 | 0 | 582 |
| 1988 | 19 | 391 | 19 | --- | --- | --- | --- | --- | 410 | 19 | 429 |
| 1989 | 2 | 334 | 83 | 26 | --- | --- | --- | --- | 336 | 109 | 445 |
| 1990 | 0 | 493 | 19 | 220 | --- | --- | 0 | 14 | 493 | 253 | 746 |
| 1991 | 3 | 257 | 99 | 161 | --- | --- | 0 | 0 | 260 | 260 | 520 |
| 1992 | 12 | 379 | 13 | 258 | --- | --- | 0 | 10 | 391 | 281 | 672 |
| 1993 | 8 | 296 | 6 | 221 | --- | --- | 0 | 2 | 304 | 229 | 533 |
| 1994 | 0 | 98 | 5 | 37 | --- | --- | 0 | 0 | 98 | 42 | 140 |
| 1995 | 2 | 19 | 11 | 22 | --- | --- | 0 | 0 | 21 | 33 | 54 |
| 1996 | 2 | 140 | 15 | 63 | --- | --- | 0 | 3 | 142 | 81 | 223 |
| 1997 | 0 | 121 | 4 | 109 | --- | --- | 0 | 9 | 121 | 122 | 243 |
| 1998 | 0 | 85 | 16 | 39 | --- | --- | 0 | 0 | 85 | 55 | 140 |
| 1999 | 0 | 3 | 59 | 162 | --- | --- | 5 | 15 | 3 | 241 | 244 |
| 2000 | 14 | 68 | 13 | 196 | --- | --- | 5 | 41 | 82 | 255 | 337 |
| 2001 | 9 | 701 | 97 | 177 | --- | --- | 13 | 0 | 710 | 287 | 997 |
| 2002 | 9 | 341 | 11 | 546 | --- | --- | 0 | 97 | 350 | 654 | 1,004 |
| 2003 | 3 | 244 | 26 | 169 | --- | --- | 1 | 0 | 247 | 196 | 443 |
| 2004 | 0 | 400 | 19 | 134 | 3 | 0 | 0 | 16 | 400 | 172 | 572 |
| 2005 | 3 | 299 | 6 | 107 | 0 | 14 | 2 | 4 | 302 | 133 | 435 |
| 2006 | 7 | 145 | 2 | 100 | 2 | 2 | 0 | 8 | 152 | 114 | 266 |
| 2007 | 8 | 190 | 18 | 81 | 0 | 19 | 15 | 13 | 198 | 146 | 344 |
| 2008 | 131 | 403 | 291 | 102 | 158 | 82 | 23 | 1 | 534 | 657 | 1,191 |
| 2009 | 116 | 634 | 402 | 403 | 92 | 196 | 13 | 4 | 750 | 1,110 | 1,860 |
| 2010 | 41 | 1,402 | 74 | 679 | 0 | 306 | 4 | 17 | 1,443 | 1,080 | 2,523 |
| 2011 | 85 | 671 | 269 | 212 | 0 | 27 | 12 | 24 | 756 | 544 | 1,300 |
| 2012 | 7 | 806 | 8 | 385 | --- | --- | 0 | 29 | 813 | 422 | 1,235 |
| 2013 | 91 | 660 | 66 | 296 | --- | --- | 2 | 0 | 751 | 364 | 1,115 |
| 2014 | 41 | 857 | 62 | 114 | --- | --- | 0 | 12 | 898 | 188 | 1,086 |
| 2015 | 65 | 667 | 184 | 648 | --- | --- | 6 | 207 | 732 | 1,045 | 1,777 |
| 2016 | 8 | 215 | 120 | 335 | --- | --- | 12 | 62 | 223 | 529 | 752 |
| 2017 | 9 | 60 | 140 | 257 | --- | --- | 19 | 27 | 69 | 443 | 512 |
| 2018 | 0 | 80 | 39 | 316 | --- | --- | 1 | 109 | 80 | 465 | 545 |
| 2019 | 2 | 43 | 29 | 128 | --- | --- | 0 | 1 | 45 | 158 | 203 |
| 2020 | 3 | 53 | 2 | 21 | --- | --- | 0 | 2 | 56 | 25 | 81 |
| 2021 | 8 | 100 | 3 | 15 | --- | --- | 19 | 70 | 108 | 107 | 215 |

Appendix E: Stray Hatchery-Origin Spring Chinook Salmon in the Tucannon River (1990-2021)

Appendix E. Summary of identified stray hatchery origin spring Chinook salmon that escaped into the Tucannon River (1990-2021).

| Year | CWT Code or Fin clip | Agency | Origin (stock) | Release Location / Release River | Number Observed/Expanded ^a | % of Tuc. Run |
|---------------------|----------------------|--------|---------------------|----------------------------------|---------------------------------------|---------------|
| 1990 | 074327 | ODFW | Carson (Wash.) | Meacham Cr./Umatilla River | 2 / 5 | |
| | 074020 | ODFW | Rapid River | Lookingglass Cr./Grande Ronde | 1 / 2 | |
| | 232227 | NMFS | Mixed Col. | Columbia River/McNary Dam | 2 / 5 | |
| | 232228 | NMFS | Mixed Col. | Columbia River/McNary Dam | 1 / 2 | |
| Total Strays | | | | | 14 | 1.9 |
| 1992 | 075107 | ODFW | Lookingglass Cr. | Bonifer Pond/Umatilla River | 2 / 6 | |
| | 075111 | ODFW | Lookingglass Cr. | Meacham Cr./Umatilla River | 1 / 2 | |
| | 075063 | ODFW | Lookingglass Cr. | Meacham Cr./Umatilla River | 1 / 2 | |
| Total Strays | | | | | 10 | 1.3 |
| 1993 | 075110 | ODFW | Lookingglass Cr. | Meacham Cr./Umatilla River | 1 / 2 | |
| | Total Strays | | | | | |
| 1996 | 070251 | ODFW | Carson (Wash.) | Imeqes AP/Umatilla River | 1 / 1 | |
| | LV clip | ODFW | Carson (Wash.) | Imeqes AP/Umatilla River | 1 / 2 | |
| Total Strays | | | | | 3 | 1.3 |
| 1997 | 103042 | IDFG | South Fork Salmon | Knox Bridge/South Fork Salmon | 1 / 2 | |
| | 103518 | IDFG | Powell | Powell Rearing Ponds/Lochsa R. | 1 / 2 | |
| | RV clip | ODFW | Carson (Wash.) | Imeqes AP/Umatilla River | 3 / 5 | |
| Total Strays | | | | | 9 | 2.6 |
| 1999 | 091751 | ODFW | Carson (Wash.) | Imeqes AP/Umatilla River | 2 / 3 | |
| | 092258 | ODFW | Carson (Wash.) | Imeqes AP/Umatilla River | 1 / 1 | |
| | 104626 | UI | Eagle Creek NFH | Eagle Creek NFH/Clackamas R. | 1 / 1 | |
| | LV clip | ODFW | Carson (Wash.) | Imeqes AP/Umatilla River | 2 / 2 | |
| | RV clip | ODFW | Carson (Wash.) | Imeqes AP/Umatilla River | 8 / 13 | |
| Total Strays | | | | | 20 | 8.2 |
| 2000 | 092259 | ODFW | Carson (Wash.) | Imeqes AP/Umatilla River | 4 / 4 | |
| | 092260 | ODFW | Carson (Wash.) | Imeqes AP/Umatilla River | 1 / 1 | |
| | 092262 | ODFW | Carson (Wash.) | Imeqes AP/Umatilla River | 1 / 3 | |
| | 105137 | IDFG | Powell | Walton Creek/Lochsa R. | 1 / 3 | |
| | 636330 | WDFW | Klickitat (Wash.) | Klickitat Hatchery | 1 / 1 | |
| | 636321 | WDFW | Lyons Ferry (Wash.) | Lyons Ferry/Snake River | 1 / 1 | |
| | LV clip | ODFW | Carson (Wash.) | Imeqes AP/Umatilla River | 18 / 31 | |
| | Ad clip | ODFW | Carson (Wash.) | Imeqes AP/Umatilla River | 2 / 2 | |
| Total Strays | | | | | 46 | 13.6 |
| 2001 | 076040 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 1/7 | |
| | 092828 | ODFW | Imnaha R. & Tribs. | Lookingglass/Imnaha River | 1/3 | |
| | 092829 | ODFW | Imnaha R. & Tribs. | Lookingglass/Imnaha River | 1/3 | |
| Total Strays | | | | | 13 | 1.3 |

^a The expansion is based on subsample rates of the proportion of stray carcasses to Tucannon River origin carcasses from the river. Actual counts are not expanded.

Appendix E (continued). Summary of identified stray hatchery origin spring Chinook salmon that escaped into the Tucannon River (1990-2021).

| Year | CWT Code or Fin clip | Agency | Origin (stock) | Release Location / Release River | Number Observed/Expanded ^a | % of Tuc. Run |
|------|----------------------|---------|----------------------|----------------------------------|---------------------------------------|---------------|
| 2002 | 054208 | USFWS | Dworshak | Dworshak NFH/Clearwater R. | 1/29 | |
| | 076039 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 1/8 | |
| | 076040 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 2/16 | |
| | 076041 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 2/16 | |
| | 076049 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 1/8 | |
| | 076051 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 1/8 | |
| | 076138 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 1/8 | |
| | 105412 | IDFG | Powell | Clearwater Hatch./Powell Ponds | 1/4 | |
| | | | Total Strays | 97 | 9.7 | |
| 2003 | 100472 | IDFG | Salmon R. | Sawtooth Hatch./Nature's Rear. | 1/1 | |
| | | | | Total Strays | 1 | 0.2 |
| 2004 | Ad clip | Unknown | Unknown | Unknown | 6/17 | |
| | | | | Total Strays | 17 | 3.0 |
| 2005 | Ad clip | Unknown | Unknown | Unknown | 3/6 | |
| | | | | Total Strays | 6 | 1.4 |
| 2006 | 109771 | IDFG | Sum. Ch. - S Fk Sal. | McCall Hatch./S. Fk. Salmon R. | 1/1 | |
| | 093859 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 1/1 | |
| | Ad clip | Unknown | Unknown | Unknown | 3/6 | |
| | | | | Total Strays | 8 | 3.2 |
| 2007 | 092043 | ODFW | Rogue R. – Cole H. | Cole Rivers Hatchery/Rogue R. | 1/1 | |
| | Ad clip | Unknown | Unknown | Unknown | 9/27 | |
| | | | | Total Strays | 28 | 8.1 |
| 2008 | 092045 | ODFW | Rogue R. – Cole H. | Cole Rivers Hatchery/Rogue R. | 1/1 | |
| | 094358 | ODFW | Grande Ronde R. | Lookingglass/Grande Ronde R. | 1/11 | |
| | 094460 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 1/11 | |
| | Ad clip | Unknown | Unknown | Unknown | 1/1 | |
| | | | | Total Strays | 24 | 2.0 |
| 2009 | 092043 | ODFW | Rogue R. | Cole Rivers Hatch./Rogue R. | 1/3 | |
| | 094532 | ODFW | Imnaha R. | Lookingglass Hatch./Imnaha R. | 1/3 | |
| | 094538 | ODFW | Lostine R. | Lookingglass/Lostine R. | 2/4 | |
| | 100181 | IDFG | Salmon R. Sum. Ck. | Knox Bridge/S. Fork Salmon | 1/1 | |
| | Ad clip | Unknown | Unknown | Unknown | 6/6 | |
| | | | | Total Strays | 17 | 0.9 |
| 2010 | 092737 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 1/6 | |
| | 094351 | ODFW | Lostine R. | Lookingglass/Lostine R. | 1/6 | |
| | Ad clip | Unknown | Unknown | Unknown | 9/9 | |
| | | | | Total Strays | 21 | 0.8 |
| 2011 | 054685 | USFWS | Dworshak | Dworshak Hatchery | 1/1 | |
| | 094591 | ODFW | Catherine Ck. | Lookingglass Hatchery | 2/2 | |
| | 094593 | ODFW | Lookingglass Ck. | Lookingglass Hatchery | 1/1 | |
| | 094665 | ODFW | Lostine R. | Lookingglass Hatchery | 1/6 | |
| | 101381 | IDFG | Clear Ck. | Clearwater Hatchery/Powell | 1/6 | |
| | 102380 | IDFG | S.F. Clearwater | Clearwater Hatchery | 1/6 | |
| | 105081 | IDFG | Selway R. | Clearwater Hatchery/Powell | 1/6 | |
| | Ad clip | Unknown | Unknown | Unknown | 3/8 | |
| | | | | Total Strays | 36 | 2.8 |

^a The expansion is based on subsample rates of the proportion of stray carcasses to Tucannon River origin carcasses from the river. Actual counts are not expanded.

Appendix E (continued). Summary of identified stray hatchery origin spring Chinook salmon that escaped into the Tucannon River (1990-2021).

| Year | CWT Code or Fin clip | Agency | Origin (stock) | Release Location / Release River | Number Observed/ Expanded ^a | % of Tuc. Run |
|------|----------------------|---------|----------------|----------------------------------|--|---------------|
| 2012 | Ad clip | Unknown | Unknown | Unknown | 9/29 | |
| | | | | Total Strays | 29 | 2.3 |
| 2013 | Ad clip | Unknown | Unknown | Unknown | 2/2 | |
| | | | | Total Strays | 2 | 0.2 |
| 2014 | 090471 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 1/1 | |
| | 090485 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 1/1 | |
| | 090282 | ODFW | Lostine R. | Lookingglass/Lostine R. | 1/11 | |
| | | | | Total Strays | 13 | 1.2 |
| 2015 | 090552 | ODFW | Imnaha R. | Lookingglass/Imnaha R. | 1/14 | |
| | 090643 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 6/19 | |
| | 090652 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 15/123 | |
| | 090729 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 3/3 | |
| | Ad clip | Unknown | Unknown | Unknown | 28/54 | |
| | | | | Total Strays | 213 | 12.0 |
| 2016 | 090861 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 1/4 | |
| | 090719 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 12/31 | |
| | 090729 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 2/2 | |
| | 090733 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 1/4 | |
| | 220134 | NPT | Clearwater Mix | NPT Hatchery | 1/4 | |
| | 090652 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 2/2 | |
| | Ad clip | Unknown | Unknown | Unknown | 24/27 | |
| | | | | Total Strays | 74 | 9.8 |
| 2017 | 090910 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 1/1 | |
| | 090918 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 2/6 | |
| | 090861 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 2/6 | |
| | 190418 | Yakama | Yakima R. | Cle Elum Hatch./Yakima River | 1/5 | |
| | Ad clip | Unknown | Unknown | Unknown | 17/28 | |
| | | | | Total Strays | 46 | 9.0 |
| 2018 | 090903 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 2/2 | |
| | 090910 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 3/9 | |
| | 090918 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 5/15 | |
| | Ad clip | Unknown | Unknown | Unknown | 47/84 | |
| | | | | Total Strays | 110 | 20.2 |
| 2019 | Ad clip | Unknown | Unknown | Unknown | 1/1 | |
| | | | | Total Strays | 1 | 0.5 |
| 2020 | Ad clip | Unknown | Unknown | Unknown | 2/2 | |
| | | | | Total Strays | 2 | 2.5 |
| 2021 | 091338 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 1/1 | |
| | 091340 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 2/2 | |
| | 091220 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 2/6 | |
| | 091221 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 1/1 | |
| | 091223 | ODFW | Umatilla R. | Umatilla Hatch./Umatilla River | 5/13 | |
| | 09 | ODFW | Unknown | Unknown | 10/14 | |
| | AD clip | Unknown | Unknown | Unknown | 28/52 | |
| | | | | Total Strays | 89 | 41.4 |

^a The expansion is based on subsample rates of the proportion of stray carcasses to Tucannon River origin carcasses from the river. Actual counts are not expanded.

**Appendix F: Final PIT Tag Detections of Returning
Tucannon River Spring Chinook, 2015 to 2021
Calendar Years**

Appendix F. Final PIT tag detections of returning Tucannon River spring Chinook from fish originally tagged as juveniles from the Tucannon River during the 2015 to 2021 calendar years (Data for the 1995 to 2014 calendar years can be found in Gallinat and Kiefl 2019).

| PIT Tag ID | Release Data | | | Adult Return Final Detection Data ^a | | | |
|-----------------|--------------|-------------|--------------|--|----------|-------------|----------|
| | Origin | Length (mm) | Release Date | OBS | OBS Date | Travel Time | Est. Age |
| 3DD.00775150D8 | W | 118 | 4/28/15 | LTR | 6/14/17 | 778 | 4 |
| 3DD.0077484E81 | H | 133 | 4/06/15 | UTR ^b | 6/04/16 | 425 | 3 |
| 3DD.0077487AD0 | H | 162 | 4/06/15 | UTR | 5/30/16 | 420 | 3 |
| 3DD.007748AE73 | H | 147 | 4/06/15 | UTR | 7/20/16 | 471 | 3 |
| 3DD.007749A8C2 | H | 136 | 4/06/15 | UTR | 9/21/16 | 444 | 3 |
| 3DD.007749DDBD | H | 148 | 4/06/15 | UTR ^b | 6/23/16 | 444 | 3 |
| 3DD.007749EDDD | H | 127 | 4/06/15 | UTR ^b | 7/02/16 | 453 | 3 |
| 3DD.00774A59CE | H | 163 | 4/06/15 | UTR | 6/13/16 | 434 | 3 |
| 3DD.00774A73B1 | H | 138 | 4/06/15 | MTR | 5/31/16 | 421 | 3 |
| 3DD.00774A95A2 | H | 129 | 4/06/15 | UTR ^b | 6/19/16 | 440 | 3 |
| 3DD.00774AC987 | H | 130 | 4/06/15 | UTR ^b | 6/07/16 | 428 | 3 |
| 3DD.007747D619 | H | 176 | 4/06/15 | TDA | 7/19/17 | 835 | 4 |
| 3DD.007747F7ED | H | 137 | 4/06/15 | LMO | 5/29/17 | 784 | 4 |
| 3DD.00774888D6B | H | 129 | 4/06/15 | LTR | 5/27/17 | 782 | 4 |
| 3DD.0077499F22 | H | 141 | 4/06/15 | LTR | 6/10/17 | 796 | 4 |
| 3DD.007749C0F4 | H | --- | 4/06/15 | LMO | 6/10/17 | 794 | 4 |
| 3DD.007749CEEB | H | 134 | 4/06/15 | BON | 5/07/17 | 762 | 4 |
| 3DD.007749D2D4 | H | 149 | 4/06/15 | TFH ^b | 5/30/17 | 785 | 4 |
| 3DD.007749E193 | H | 146 | 4/06/15 | LMO | 6/18/17 | 804 | 4 |
| 3DD.00774A053B | H | 139 | 4/06/15 | TFH | 6/26/17 | 790 | 4 |
| 3DD.00774A2D48 | H | 149 | 4/06/15 | MTR | 7/11/17 | 827 | 4 |
| 3DD.00774A3E6D | H | 128 | 4/06/15 | LTR | 5/05/17 | 760 | 4 |
| 3DD.00774A3F26 | H | 139 | 4/06/15 | TFH | 9/06/17 | 807 | 4 |
| 3DD.00774A5ED9 | H | 158 | 4/06/15 | BON | 5/22/17 | 777 | 4 |
| 3DD.00774A9148 | H | 118 | 4/06/15 | TDA | 6/08/17 | 794 | 4 |
| 3DD.00774A97E7 | H | 139 | 4/06/15 | LMO | 6/09/17 | 795 | 4 |
| 3DD.0077710EA3 | H | 118 | 4/08/16 | LGR | 6/06/17 | 424 | 3 |
| 3DD.007774D735 | H | 133 | 4/08/16 | LGR | 7/03/17 | 420 | 3 |
| 3DD.0077751EB0 | H | 128 | 4/08/16 | TFH ^b | 6/19/17 | 437 | 3 |
| 3DD.0077754705 | H | 124 | 4/08/16 | MCN | 5/30/17 | 417 | 3 |
| 3DD.0077754B3C | H | 123 | 4/08/16 | TFH ^b | 7/06/17 | 452 | 3 |
| 3DD.0077757758 | H | 163 | 4/08/16 | TFH ^b | 7/05/17 | 445 | 3 |
| 3DD.00777577C7 | H | 159 | 4/08/16 | TFH | 6/24/17 | 435 | 3 |
| 3DD.007775AC37 | H | 152 | 4/08/16 | BON | 5/22/17 | 409 | 3 |
| 3DD.007775B4A4 | H | 159 | 4/08/16 | LMO | 6/07/17 | 425 | 3 |

Abbreviations are as follows: BON – Bonneville Dam, TDA – The Dalles Dam, MCN – McNary Dam, ICH – Ice Harbor Dam, LMO – Lower Monumental Dam, LTR – Lower Tucannon River, MTR – Middle Tucannon River, UTR – Upper Tucannon River, TFH – Tucannon Fish Hatchery, LGO – Little Goose Dam, LGR – Lower Granite Dam, AFC – Asotin Creek.

^a PIT tag adult detection systems were in operation beginning in 1988 for LGR, 1998 for BON, 2002 for MCN, 2005 for both ICH and LTR, 2011 for MTR and UTR, and 2012 for TFH.

^b This fish was detected by bypassing the Tucannon River (LGO or LGR detection) before heading back downstream.

Appendix F (continued). Final PIT tag detections of returning Tucannon River spring Chinook from fish originally tagged as juveniles from the Tucannon River during the 2015 to 2021 calendar years (Data for the 1995 to 2014 calendar years can be found in Gallinat and Kiefel 2019).

| PIT Tag ID | Release Data | | | Adult Return Final Detection Data ^a | | | |
|----------------|--------------|-------------|--------------|--|----------|-------------|----------|
| | Origin | Length (mm) | Release Date | OBS | OBS Date | Travel Time | Est. Age |
| 3DD.007775C8C1 | H | 128 | 4/08/16 | TFH | 8/04/17 | 446 | 3 |
| 3DD.007775D09B | H | 126 | 4/08/16 | TFH ^b | 6/13/17 | 431 | 3 |
| 3DD.00777F78DD | H | 161 | 4/08/16 | LMO | 5/31/17 | 418 | 3 |
| 3DD.00777FBA6E | H | 154 | 4/08/16 | LGR | 6/11/17 | 421 | 3 |
| 3DD.0077800113 | H | 135 | 4/08/16 | LGR | 6/04/17 | 421 | 3 |
| 3DD.007780EAC4 | H | 135 | 4/08/16 | LGO | 6/09/17 | 427 | 3 |
| 3DD.007780F56C | H | 150 | 4/08/16 | TFH | 6/21/17 | 439 | 3 |
| 3DD.007781CE48 | H | 140 | 4/08/16 | TFH ^b | 6/07/17 | 420 | 3 |
| 3DD.007781CF34 | H | 137 | 4/08/16 | LMO ^b | 6/08/17 | 424 | 3 |
| 3DD.00778D992C | H | 118 | 4/08/16 | TFH | 6/20/17 | 435 | 3 |
| 3DD.00776F6554 | H | 120 | 4/08/16 | TFH ^b | 6/08/18 | 791 | 4 |
| 3DD.00777169D1 | H | 161 | 4/08/16 | MTR | 5/26/18 | 778 | 4 |
| 3DD.0077719998 | H | 149 | 4/08/16 | TFH | 6/27/18 | 781 | 4 |
| 3DD.007771ADFE | H | 123 | 4/08/16 | TFH | 6/19/18 | 799 | 4 |
| 3DD.007771F0BE | H | 138 | 4/08/16 | BON | 5/24/18 | 776 | 4 |
| 3DD.007771FE88 | H | 113 | 4/08/16 | TFH | 6/20/18 | 794 | 4 |
| 3DD.0077721C1E | H | 126 | 4/08/16 | TFH | 6/07/18 | 787 | 4 |
| 3DD.0077722AB9 | H | 161 | 4/08/16 | BON | 5/27/18 | 779 | 4 |
| 3DD.007772D04C | H | 171 | 4/08/16 | TFH | 6/19/18 | 772 | 4 |
| 3DD.007774B9D1 | H | 165 | 4/08/16 | TFH | 6/14/18 | 796 | 4 |
| 3DD.007774DA7E | H | 148 | 4/08/16 | BON | 5/05/18 | 757 | 4 |
| 3DD.007774DAB6 | H | 154 | 4/08/16 | MTR | 5/22/18 | 774 | 4 |
| 3DD.007775295C | H | 138 | 4/08/16 | TFH ^b | 6/15/18 | 798 | 4 |
| 3DD.007775463E | H | 118 | 4/08/16 | JOD | 5/27/18 | 779 | 4 |
| 3DD.0077756BB3 | H | 118 | 4/08/16 | TFH ^b | 6/20/18 | 802 | 4 |
| 3DD.0077757EDF | H | 106 | 4/08/16 | BON | 4/26/18 | 748 | 4 |
| 3DD.00777583DD | H | 128 | 4/08/16 | TFH | 6/20/18 | 777 | 4 |
| 3DD.0077759EED | H | 137 | 4/08/16 | TDA | 5/03/18 | 755 | 4 |
| 3DD.007775AB57 | H | 166 | 4/08/16 | TFH | 6/12/18 | 792 | 4 |
| 3DD.007775AB97 | H | 102 | 4/08/16 | TFH ^b | 6/13/18 | 791 | 4 |
| 3DD.007775ABD7 | H | 132 | 4/08/16 | BON | 5/19/18 | 771 | 4 |
| 3DD.007775C5A1 | H | 130 | 4/08/16 | LGR | 6/07/18 | 790 | 4 |
| 3DD.007775C7BD | H | 142 | 4/08/16 | JOD | 5/27/18 | 779 | 4 |
| 3DD.007775E060 | H | 117 | 4/08/16 | MTR ^b | 6/02/18 | 785 | 4 |
| 3DD.007775E19A | H | 154 | 4/08/16 | TFH | 6/16/18 | 791 | 4 |

Abbreviations are as follows: BON – Bonneville Dam, TDA – The Dalles Dam, MCN – McNary Dam, ICH – Ice Harbor Dam, LMO – Lower Monumental Dam, LTR – Lower Tucannon River, MTR – Middle Tucannon River, UTR – Upper Tucannon River, TFH – Tucannon Fish Hatchery, LGO – Little Goose Dam, LGR – Lower Granite Dam, AFC – Asotin Creek.

^a PIT tag adult detection systems were in operation beginning in 1988 for LGR, 1998 for BON, 2002 for MCN, 2005 for both ICH and LTR, 2011 for MTR and UTR, and 2012 for TFH.

^b This fish was detected by passing the Tucannon River (LGO or LGR detection) before heading back downstream.

Appendix F (continued). Final PIT tag detections of returning Tucannon River spring Chinook from fish originally tagged as juveniles from the Tucannon River during the 2015 to 2021 calendar years (Data for the 1995 to 2014 calendar years can be found in Gallinat and Kiefel 2019).

| PIT Tag ID | Release Data | | | Adult Return Final Detection Data ^a | | | |
|----------------|--------------|-------------|--------------|--|----------|-------------|----------|
| | Origin | Length (mm) | Release Date | OBS | OBS Date | Travel Time | Est. Age |
| 3DD.007775F701 | H | 134 | 4/08/16 | TFH ^b | 6/03/18 | 780 | 4 |
| 3DD.007780CF9E | H | 118 | 4/08/16 | BON | 4/30/18 | 752 | 4 |
| 3DD.007780FEA9 | H | 129 | 4/08/16 | MTR ^b | 6/03/18 | 786 | 4 |
| 3DD.0077813299 | H | 126 | 4/08/16 | TFH | 6/07/18 | 789 | 4 |
| 3DD.00778C2417 | H | 158 | 4/08/16 | TFH ^b | 6/18/18 | 797 | 4 |
| 3DD.007774F3D6 | H | 110 | 4/08/16 | TFH | 7/18/19 | 1149 | 5 |
| 3DD.0077758E24 | H | 123 | 4/08/16 | UTR | 6/07/19 | 1155 | 5 |
| 3DD.0077510CED | W | 103 | 4/23/16 | TFH ^b | 6/16/18 | 784 | 4 |
| 3DD.0077512587 | W | 100 | 4/25/16 | BON | 5/16/18 | 751 | 4 |
| 3DD.00775159BE | W | 104 | 3/31/16 | BON | 5/10/18 | 770 | 4 |
| 3DD.007751E527 | W | 115 | 4/29/16 | TFH | 6/15/18 | 777 | 4 |
| 384.3B23A8F17E | W | 119 | 3/01/17 | MTR | 6/13/19 | 834 | 4 |
| 3DD.0077B5E4B2 | H | 178 | 4/12/17 | TFH ^b | 6/12/18 | 426 | 3 |
| 3DD.0077B6E3B1 | H | 150 | 4/12/17 | UTR | 6/01/18 | 415 | 3 |
| 3DD.0077B90D27 | H | 154 | 4/12/17 | LMO | 5/30/18 | 413 | 3 |
| 3DD.00778C9423 | H | 116 | 4/12/17 | MTR ^b | 6/21/19 | 800 | 4 |
| 3DD.00778EDD6A | H | 147 | 4/12/17 | MTR | 6/09/19 | 788 | 4 |
| 3DD.00778F01BD | H | 164 | 4/12/17 | UTR ^b | 6/01/19 | 780 | 4 |
| 3DD.0077AE2FFB | H | 115 | 4/12/17 | MTR | 6/11/19 | 790 | 4 |
| 3DD.0077B5EF67 | H | 130 | 4/12/17 | MTR | 6/04/19 | 783 | 4 |
| 3DD.0077B61920 | H | 117 | 4/12/17 | MTR | 6/26/19 | 805 | 4 |
| 3DD.0077B63DEF | H | 177 | 4/12/17 | UTR | 6/01/19 | 780 | 4 |
| 3DD.0077B64FED | H | 121 | 4/12/17 | MTR | 6/09/19 | 788 | 4 |
| 3DD.0077B68776 | H | 119 | 4/12/17 | UTR | 6/03/19 | 782 | 4 |
| 3DD.0077B697B3 | H | 153 | 4/12/17 | BON | 5/11/19 | 759 | 4 |
| 3DD.0077B90306 | H | 118 | 4/12/17 | UTR | 5/28/19 | 776 | 4 |
| 3DD.0077B92203 | H | 117 | 4/12/17 | UTR | 5/30/19 | 778 | 4 |
| 3DD.0077B972B0 | H | 148 | 4/12/17 | TFH | 6/17/19 | 779 | 4 |
| 3DD.0077A5D971 | H | 158 | 4/09/18 | LTR | 7/05/19 | 452 | 3 |
| 3DD.0077A637B7 | H | 117 | 4/09/18 | TFH | 7/12/19 | 441 | 3 |
| 3DD.0077A53DFA | H | 104 | 4/09/18 | LTR ^b | 5/26/20 | 778 | 4 |
| 3DD.0077A7064D | H | 126 | 4/09/18 | LGR | 6/06/20 | 789 | 4 |
| 3DD.00779DCA2C | W | 120 | 4/28/18 | LTR | 5/24/20 | 757 | 4 |
| 3DD.0077BF98E9 | H | --- | 4/18/19 | TFH | 5/27/21 | 770 | 4 |
| 3DD.00778C429A | W | 107 | 4/26/19 | TFH | 7/08/21 | 804 | 4 |

Abbreviations are as follows: BON – Bonneville Dam, TDA – The Dalles Dam, MCN – McNary Dam, ICH – Ice Harbor Dam, LMO – Lower Monumental Dam, LTR – Lower Tucannon River, MTR – Middle Tucannon River, UTR – Upper Tucannon River, TFH – Tucannon Fish Hatchery, LGO – Little Goose Dam, LGR – Lower Granite Dam, AFC – Asotin Creek.

^a PIT tag adult detection systems were in operation beginning in 1988 for LGR, 1998 for BON, 2002 for MCN, 2005 for both ICH and LTR, 2011 for MTR and UTR, and 2012 for TFH.

^b This fish was detected by passing the Tucannon River (LGO or LGR detection) before heading back downstream.

Appendix F (continued). Final PIT tag detections of returning Tucannon River spring Chinook from fish originally tagged as juveniles from the Tucannon River during the 2015 to 2021 calendar years (Data for the 1995 to 2014 calendar years can be found in Gallinat and Kiefel 2019).

| PIT Tag ID | Release Data | | | Adult Return Final Detection Data ^a | | | |
|----------------|--------------|-------------|--------------|--|----------|-------------|----------|
| | Origin | Length (mm) | Release Date | OBS | OBS Date | Travel Time | Est. Age |
| 3DD.00779E27A3 | W | 114 | 4/28/19 | TFH | 6/02/21 | 766 | 4 |
| 3DD.00779F22E5 | W | 111 | 5/02/19 | TFH | 6/12/21 | 772 | 4 |
| 3DD.00779FC6F8 | W | 112 | 5/05/19 | BON | 5/18/21 | 744 | 4 |
| 3DD.00779FD92F | W | 120 | 5/02/19 | TFH | 6/13/21 | 773 | 4 |

Abbreviations are as follows: BON – Bonneville Dam, TDA – The Dalles Dam, MCN – McNary Dam, ICH – Ice Harbor Dam, LMO – Lower Monumental Dam, LTR – Lower Tucannon River, MTR – Middle Tucannon River, UTR – Upper Tucannon River, TFH – Tucannon Fish Hatchery, LGO – Little Goose Dam, LGR – Lower Granite Dam, AFC – Asotin Creek.

^a PIT tag adult detection systems were in operation beginning in 1988 for LGR, 1998 for BON, 2002 for MCN, 2005 for both ICH and LTR, 2011 for MTR and UTR, and 2012 for TFH.

Appendix G: Historical Hatchery Releases (1987-2022 Release Years)

Appendix G. Historical hatchery spring Chinook releases from the Tucannon River, 1987-2022 release years.
(Totals are summation by brood year and release year.)

| Release Year | Brood | Release | | CWT Code ^b | Number CWT | Ad-only marked | Additional Tag/location/cross ^c | Kg | Mean Wt. (g) |
|--------------|-------|-------------------|-----------|-----------------------|----------------|----------------|--|-------|--------------|
| | | Type ^a | Date | | | | | | |
| 1987 | 1985 | H-Acc | 4/6-10 | 34/42 | 12,922 | | | 986 | 76 |
| Total | | | | | 12,922 | | | | |
| 1988 | 1986 | H-Acc | 3/7 | 33/25 | 12,328 | 512 | | 628 | 45 |
| | | " | " | 41/46 | 12,095 | 465 | | 570 | 45 |
| | | " | " | 41/48 | 13,097 | 503 | | 617 | 45 |
| | | " | 4/13 | 33/25 | 37,893 | 1,456 | | 1,696 | 45 |
| | | " | " | 41/46 | 34,389 | 1,321 | | 1,621 | 45 |
| | | " | " | 41/48 | 37,235 | 1,431 | | 1,756 | 45 |
| Total | | | | | 147,037 | 5,688 | | | |
| 1989 | 1987 | H-Acc | 4/11-13 | 49/50 | 151,100 | 1,065 | | 7,676 | 50 |
| Total | | | | | 151,100 | 1,065 | | | |
| 1990 | 1988 | H-Acc | 3/30-4/10 | 55/01 | 68,591 | 3,007 | | 2,955 | 41 |
| | | " | " | 01/42 | 70,459 | 3,089 | | 3,035 | 41 |
| Total | | | | | 139,050 | 6,096 | | | |
| 1991 | 1989 | H-Acc | 4/1-12 | 14/61 | 75,661 | 989 | | 3,867 | 50 |
| | | " | " | 01/31 | 22,118 | 289 | | 1,130 | 50 |
| Total | | | | | 97,779 | 1,278 | | | |
| 1992 | 1990 | H-Acc | 3/30-4/10 | 40/21 | 51,149 | | BWT, RC, WxW | 2,111 | 41 |
| | | " | " | 43/11 | 21,108 | | BWT, LC, HxH | 873 | 41 |
| | | " | " | 37/25 | 13,480 | | Mixed | 556 | 41 |
| Total | | | | | 85,737 | | | | |
| 1993 | 1991 | H-Acc | 4/6-12 | 46/25 | 55,716 | 796 | VI, LR, WxW | 1,686 | 30 |
| | | " | " | 46/47 | 16,745 | 807 | VI, RR, HxH | 507 | 30 |
| Total | | | | | 72,461 | 1,603 | | | |
| 1993 | 1992 | Direct | 10/22-25 | 48/23 | 24,883 | 251 | VI, LR, WxW | 317 | 13 |
| | | " | " | 48/24 | 24,685 | 300 | VI, RR, HxH | 315 | 13 |
| | | " | " | 48/56 | 7,111 | 86 | Mixed | 91 | 13 |
| Total | | | | | 56,679 | 637 | | | |
| 1994 | 1992 | H-Acc | 4/11-18 | 48/10 | 35,405 | 871 | VI, LY, WxW | 1,176 | 32 |
| | | " | " | 49/05 | 35,469 | 2,588 | VI, RY, HxH | 1,234 | 32 |
| | | " | " | 48/55 | 8,277 | 799 | Mixed | 294 | 32 |
| Total | | | | | 79,151 | 4,258 | | | |
| 1995 | 1993 | H-Acc | 3/15-4/15 | 53/43 | 45,007 | 140 | VI, RG, HxH | 1,437 | 32 |
| | | " | " | 53/44 | 42,936 | 2,212 | VI, LG, WxW | 1,437 | 32 |
| | | P-Acc | 3/20-4/3 | 56/15 | 11,661 | 72 | VI, RR, HxH | 355 | 30 |
| | | " | " | 56/17 | 10,704 | 290 | VI, LR, WxW | 333 | 30 |
| | | " | " | 56/18 | 13,705 | 47 | Mixed | 416 | 30 |
| | | Direct | 3/20-4/3 | 56/15 | 3,860 | 24 | VI, RR, HxH | 118 | 30 |
| | | " | " | 56/17 | 3,542 | 96 | VI, LR, WxW | 110 | 30 |
| | | " | " | 56/18 | 4,537 | 15 | Mixed | 138 | 30 |
| Total | | | | | 135,952 | 2,896 | | | |
| 1996 | 1994 | H-Acc | 3/16-4/22 | 56/29 | 89,437 | | VI, RR, Mixed | 2,326 | 26 |
| | | P-Acc | 3/27-4/19 | 57/29 | 35,334 | 35 | VI, RG, Mixed | 1,193 | 30 |
| | | Direct | 3/27 | 43/23 | 5,263 | | VI, LG, Mixed | 168 | 34 |
| Total | | | | | 130,034 | 35 | | | |

Appendix G (continued). Historical hatchery spring Chinook releases from the Tucannon River, 1987-2022 release years. (Totals are summation by brood year and release year.)

| Release Year | Brood | Release | | CWT Code ^b | Number CWT | Ad-only marked | Additional Tag/location/cross ^c | Kg | Mean Wt. (g) |
|--------------|--------|-------------------|-----------|-----------------------|----------------|--------------------------|--|-------|--------------|
| | | Type ^a | Date | | | | | | |
| 1997 | 1995 | H-Acc | 3/07-4/18 | 59/36 | 42,160 | 40 | VI, RR, Mixed | 1,095 | 26 |
| | | P-Acc | 3/24-3/25 | 61/41 | 10,045 | 50 | VI, RB, Mixed | 244 | 24 |
| | | Direct | 3/24 | 61/40 | 9,811 | 38 | VI, LB, Mixed | 269 | 27 |
| Total | | | | | 62,016 | 128 | | | |
| 1998 | 1996 | H-Acc | 3/11-4/17 | 03/60 | 14,308 | 27 | Mixed | 410 | 29 |
| | | C-Acc | 3/11-4/18 | 61/25 | 23,065 | 62 | “ | 680 | 29 |
| | | “ | “ | 61/24 | 24,554 | 50 | “ | 707 | 29 |
| | | Direct | 4/03 | 03/59 | 14,101 | 52 | “ | 392 | 28 |
| Total | | | | 76,028 | 191 | | | | |
| 1999 | 1997 | C-Acc | 3/11-4/20 | 61/32 | 23,664 | 522 | Mixed | 704 | 29 |
| Total | | | | | 23,664 | 522 | | | |
| 2000 | 1998 | C-Acc | 3/20-4/26 | 12/11 | 125,192 | 2,747 | Mixed | 4,647 | 36 |
| Total | | | | | 125,192 | 2,747 | | | |
| 2001 | 1999 | C-Acc | 3/19-4/25 | 02/75 | 96,736 | 864 | Mixed | 4,180 | 43 |
| Total | | | | | 96,736 | 864 | | | |
| 2002 | 2000 | C-Acc | 3/15-4/23 | 08/87 | 99,566 | 2,533 ^e | VI, RR, Mixed | 2,990 | 29 |
| Total | | | | | 99,566 | 2,533^e | | | |
| 2002 | 2000CB | C-Acc | 3/15/4/23 | 63 | 3,031 | 24 ^f | CB, Mixed | 156 | 51 |
| Total | | | | | 3,031 | 24^f | | | |
| 2002 | 2001 | Direct | 5/06 | 14/29 | 19,948 | 1,095 | Mixed | 77 | 4 |
| Total | | | | | 19,948 | 1,095 | | | |
| 2002 | 2001CB | Direct | 5/06 | 14/30 | 20,435 | 157 | CB, Mixed | 57 | 3 |
| Total | | | | | 20,435 | 157 | | | |
| 2003 | 2001 | C-Acc | 4/01-4/21 | 06/81 | 144,013 | 2,909 ^e | VI, RR, Mixed | 5,171 | 35 |
| Total | | | | | 144,013 | 2,909^e | | | |
| 2003 | 2001CB | C-Acc | 4/01-4/21 | 63 | 134,401 | 5,995 ^f | CB, Mixed | 4,585 | 33 |
| Total | | | | | 134,401 | 5,995^f | | | |
| 2004 | 2002 | C-Acc | 4/01-4/20 | 17/91 | 121,774 | 1,812 ^e | VI, RR, Mixed | 4,796 | 39 |
| Total | | | | | 121,774 | 1,812^e | | | |
| 2004 | 2002CB | C-Acc | 4/01-4/20 | 63 | 42,875 | 1,909 ^f | CB, Mixed | 1,540 | 34 |
| Total | | | | | 42,875 | 1,909^f | | | |
| 2005 | 2003 | C-Acc | 3/28-4/15 | 24/82 | 69,831 | 1,323 ^e | VI, RR, Mixed | 2,544 | 36 |
| Total | | | | | 69,831 | 1,323^e | | | |
| 2005 | 2003CB | C-Acc | 3/28-4/15 | 27/78 | 125,304 | 4,760 ^f | CB, Mixed | 4,407 | 34 |
| Total | | | | | 125,304 | 4,760^f | | | |
| 2006 | 2004 | C-Acc | 4/03-4/26 | 28/87 | 67,272 | 270 ^e | VI, RR, Mixed | 2,288 | 34 |
| Total | | | | | 67,272 | 270^e | | | |
| 2006 | 2004CB | C-Acc | 4/03-4/26 | 28/65 | 127,162 | 5,150 ^f | CB, Mixed | 3,926 | 30 |
| Total | | | | | 127,162 | 5,150^f | | | |
| 2007 | 2005 | C-Acc | 4/02-4/23 | 35/99 | 144,833 | 4,633 ^e | VI, RR, Mixed | 8,482 | 57 |
| Total | | | | | 144,833 | 4,633^e | | | |
| 2007 | 2005CB | C-Acc | 4/02-4/23 | 34/77 | 88,885 | 1,171 ^f | CB, Mixed | 5,525 | 61 |
| Total | | | | | 88,885 | 1,171^f | | | |

Appendix G (continued). Historical hatchery spring Chinook releases from the Tucannon River, 1987-2022 release years. (Totals are summation by brood year and release year.)

| Release Year | Brood | Release | | CWT Code ^b | Number CWT | Ad-only marked | Additional Tag/location/cross ^c | Kg | Mean Wt. (g) |
|--------------|--------|-------------------|-----------|-----------------------|----------------|---------------------------|--|--------|--------------|
| | | Type ^a | Date | | | | | | |
| 2008 | 2006 | C-Acc | 4/08-4/22 | 40/93 | 50,309 | 2,426 ^e | VI, LB, Mixed | 2,850 | 54 |
| 2008 | 2006 | C-Acc | 4/08-4/22 | 40/94 | 51,858 | 1,937 ^e | VI, LP, Mixed | 2,106 | 39 |
| Total | | | | | 102,167 | 4,363^e | | | |
| 2008 | 2006CB | C-Acc | 4/08-4/22 | 41/94 | 75,283 | 2,893 ^f | CB, Mixed | 4,493 | 57 |
| Total | | | | | 75,283 | 2,893^f | | | |
| 2009 | 2007 | C-Acc | 4/13-4/22 | 46/88 | 55,266 | 214 ^e | VI, LB, Mixed | 3,188 | 57 |
| 2009 | 2007 | C-Acc | 4/13-4/22 | 46/87 | 58,044 | 1,157 ^e | VI, LP, Mixed | 2,203 | 37 |
| Total | | | | | 113,310 | 1,371^e | | | |
| 2010 | 2008 | C-Acc | 4/2-4/12 | 51/75 | 84,738 | 1,465 ^e | VI, LB, Mixed | 5,672 | 66 |
| 2010 | 2008 | C-Acc | 4/2-4/12 | 51/74 | 84,613 | 2,081 ^e | VI, LP, Mixed | 3,423 | 40 |
| Total | | | | | 169,351 | 3,546^e | | | |
| 2010 | 2009 | Direct | 4/22-4/23 | None | 0 | 52,253 ^f | Oxytet., Mixed | 342 | 7 |
| Total | | | | | 0 | 52,253^f | | | |
| 2011 | 2009 | C-Acc | 4/7-4/25 | 55/66 | 113,049 | 0 ^e | VI, LB, Mixed | 5,767 | 51 |
| 2011 | 2009 | C-Acc | 4/7-4/25 | 55/65 | 117,824 | 564 ^e | VI, LP, Mixed | 4,135 | 35 |
| Total | | | | | 230,873 | 564^e | | | |
| 2012 | 2010 | C-Acc | 4/11-4/23 | 60/76 | 96,984 | 275 ^e | VI, LB, Mixed | 6,400 | 66 |
| 2012 | 2010 | C-Acc | 4/11-4/23 | 60/75 | 102,169 | 2,157 ^e | VI, LP, Mixed | 3,312 | 32 |
| Total | | | | | 199,153 | 2,432^e | | | |
| 2012 | 2011 | Direct | 5/01 | None | 0 | 39,460 ^f | Oxytet., Mixed | 285 | 7 |
| Total | | | | | 0 | 39,460^f | | | |
| 2013 | 2011 | C-Acc | 4/3-4/22 | 64/42 | 27,748 | 1,825 ^f | TFH reared, Mixed | 987 | 33 |
| 2013 | 2011 | C-Acc | 4/3-4/22 | 64/41 | 227,703 | 2,688 ^f | LFH reared, Mixed | 7,691 | 33 |
| Total | | | | | 255,451 | 4,513^f | | | |
| 2014 | 2012 | C-Acc | 4/11-4/23 | 65/86 | 21,101 | 1,916 ^f | TFH reared, Mixed | 746 | 32 |
| 2014 | 2012 | C-Acc | 4/11-4/23 | 65/85 | 179,400 | 1,093 ^f | LFH reared, Mixed | 5,853 | 32 |
| Total | | | | | 200,501 | 3,009^f | | | |
| 2015 | 2013 | C-Acc | 3/27-4/16 | 67/43 | 20,373 | 3,061 ^f | TFH reared, Mixed | 872 | 37 |
| 2015 | 2013 | C-Acc | 3/27-4/16 | 67/42 | 179,494 | 4,931 ^f | LFH reared, Mixed | 6,863 | 37 |
| Total | | | | | 199,867 | 7,992^f | | | |
| 2016 | 2014 | C-Acc | 4/01-4/15 | 68/84 | 216,295 | 4,804 ^f | Mixed | 8,883 | 40 |
| Total | | | | | 216,295 | 4,804^f | | | |
| 2017 | 2015 | C-Acc | 4/04-4/21 | 70/39 | 187,601 | 12,085 ^f | Mixed | 7,883 | 40 |
| Total | | | | | 187,601 | 12,085^f | | | |
| 2018 | 2016 | C-Acc | 4/09-4/27 | 72/01 | 202,952 | 6,079 ^f | Mixed | 11,434 | 55 |
| Total | | | | | 202,952 | 6,079^f | | | |
| 2019 | 2017 | C-Acc | 4/04-5/03 | 73/96 | 140,262 | 3,957 ^f | Mixed | 4,308 | 30 |
| Total | | | | | 140,262 | 3,957^f | | | |
| 2020 | 2018 | Direct | 3/23-3/24 | 74/21 | 185,758 | 6,763 ^f | Mixed | 6,993 | 36 |
| Total | | | | | 185,758 | 6,763^f | | | |
| 2021 | 2019 | Direct | 3/24 | 77/61 | 65,969 | 1,771 ^f | Mixed | 2,629 | 39 |
| 2021 | 2019 | Direct | 3/15 | 77/61 | 12,908 | 347 ^f | High ELISA, Mixed | 537 | 41 |
| Total | | | | | 78,877 | 2,118^f | | | |
| 2022 | 2020 | Direct | 4/11 | 79/43 | 42,046 | 759 ^f | TFH Release | 1,689 | 40 |
| 2022 | 2020 | Direct | 4/22 | 79/43 | 19,974 | 361 ^f | Mouth Release | 743 | 37 |
| Total | | | | | 62,020 | 1,120^f | | | |

^a Release types are: Tucannon Hatchery Acclimation Pond (H-Acc); Portable Acclimation Pond (P-Acc); Curl Lake Acclimation Pond (C-Acc); and Direct Stream Release (Direct).

^b All tag codes start with agency code 63.

^c Codes listed in column are as follows: BWT - Blank Wire Tag; CB - Captive Brood; VI-Visual Implant (elastomer); LR - Left Red, RR - Right Red, LG-Left Green, RG - Right Green, LY - Left Yellow, RY - Right Yellow, LB - Left Blue, RB - Right Blue, LP - Left Purple; Oxytet. - Oxytetracycline Mark; Crosses: WxW - wild x wild progeny, HxH - hatchery x hatchery progeny, Mixed - wild x hatchery progeny.

^d No tag loss data due to presence of both CWT and BWT in fish.

^e VI tag only.

^f No wire.

**Appendix H: Numbers of Fish Species Captured by
Month in the Tucannon River Smolt Trap during the
2021 Outmigration**

Appendix H. Numbers of fish species captured by month in the Tucannon River smolt trap during the 2021 outmigration sampling period (1 October 2020 – 6 July 2021).

| Species | Oct | Nov | Dec | Jan | Feb | Mar | Apr | May | Jun | Jul | Total |
|-------------------------------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|--------------|
| Nat. spring Chinook | | 3 | | | | 2 | 4 | 4 | | | 13 |
| Hatch. spring Chinook | | | | | | 110 | 687 | 43 | | | 840 |
| Fall Chinook | | | | | 28 | 217 | 2483 | 2577 | 541 | | 5846 |
| Coho salmon | | | | | 1 | 18 | 23 | 138 | 97 | | 277 |
| Steelhead < 80 mm | | | | | | | | 5 | 88 | 11 | 104 |
| Steelhead 80-124 mm | 36 | 20 | 1 | | | 1 | | | | | 58 |
| Steelhead ≥ 125 mm | 248 | 64 | 18 | 1 | 20 | 24 | 134 | 1043 | 26 | | 1578 |
| Hat. endemic steelhead | | | | | | | 282 | 1040 | 44 | | 1366 |
| Bull trout | 1 | | 1 | 1 | | | | | | | 3 |
| Pacific lamprey - Ammocoetes | 4 | 14 | 15 | 198 | 242 | 44 | 9 | 22 | 6 | 1 | 555 |
| Pacific lamprey - Macrophthalmia | 1 | 7 | 8 | 77 | 144 | | | 3 | | | 240 |
| Pacific lamprey - Adults | | | | | | | | 5 | 2 | | 7 |
| American shad | | | | 2 | | 1 | | | | | 3 |
| Smallmouth bass | 7 | | | 1 | | 2 | 4 | 12 | 4 | 3 | 33 |
| Pumpkinseed sunfish | 13 | 4 | | 4 | 2 | | 12 | 20 | 7 | 1 | 63 |
| Bluegill | | | | | | | 1 | | 1 | | 2 |
| Chiselmouth | 35 | 15 | 4 | | | 1 | 9 | 7 | 6 | 3 | 80 |
| Longnose dace | 46 | 17 | | | 3 | | 5 | 248 | 34 | 2 | 355 |
| Speckled dace | | | | | | | | 6 | | | 6 |
| Redside shiner | 76 | 19 | 1 | | 1 | | 32 | 261 | 105 | 7 | 502 |
| Bridgelip sucker | 94 | 28 | 11 | 2 | 2 | 14 | 34 | 90 | 14 | 5 | 294 |
| Northern pikeminnow | 4 | 3 | | | 3 | 1 | 1 | 16 | 2 | | 30 |
| Brown bullhead | | | | | | | | | 1 | | 1 |
| Mountain whitefish | | | 1 | | | | | | | | 1 |
| Sculpin sp. | | 1 | | | | | 2 | 1 | | | 4 |

Appendix I: Proportionate Natural Influence (PNI) for the Tucannon Spring Chinook Population (1985-2021)

Appendix I. Proportionate Natural Influence (PNI)^a for the Tucannon River spring Chinook population (1985-2021). Note: Pre-spawn and trap mortalities are excluded from the analysis.

| Year | Spawmed Hatchery Broodstock | | River Spawning Fish | | PNI | PNI < 0.50 |
|------|--------------------------------|---------------------|---------------------|----------------------|------|---------------|
| | Total | % Natural (PNOB) | Total | % Hatchery (PHOS) | | |
| 1985 | 8 | 100.00 | 695 | 0.00 | 1.00 | |
| 1986 | 91 | 100.00 | 440 | 0.00 | 1.00 | |
| 1987 | 83 | 100.00 | 407 | 0.00 | 1.00 | |
| 1988 | 90 | 100.00 | 257 | 0.00 | 1.00 | |
| 1989 | 122 | 45.08 | 276 | 1.09 | 0.98 | |
| 1990 | 62 | 48.39 | 572 | 21.50 | 0.69 | |
| 1991 | 71 | 56.34 | 291 | 32.30 | 0.64 | |
| 1992 | 82 | 45.12 | 476 | 35.92 | 0.56 | |
| 1993 | 87 | 51.72 | 397 | 38.29 | 0.57 | |
| 1994 | 69 | 50.72 | 97 | 0.00 | 1.00 | |
| 1995 | 39 | 23.08 | 27 | 0.00 | 1.00 | |
| 1996 | 75 | 44.00 | 152 | 23.03 | 0.66 | |
| 1997 | 89 | 42.70 | 105 | 35.24 | 0.55 | |
| 1998 | 86 | 52.33 | 60 | 26.67 | 0.66 | |
| 1999 | 122 | 0.82 | 160 | 97.50 | 0.01 | * |
| 2000 | 73 | 10.96 | 201 | 69.15 | 0.14 | * |
| 2001 | 104 | 50.00 | 766 | 19.84 | 0.72 | |
| 2002 | 93 | 45.16 | 568 | 60.56 | 0.43 | * |
| 2003 | 75 | 54.67 | 329 | 25.84 | 0.68 | |
| 2004 | 88 | 54.55 | 346 | 17.34 | 0.76 | |
| 2005 | 95 | 49.47 | 264 | 19.70 | 0.72 | |
| 2006 | 88 | 40.91 | 202 | 24.26 | 0.63 | |
| 2007 | 82 | 62.20 | 211 | 22.27 | 0.74 | |
| 2008 | 114 | 35.09 | 796 | 38.94 | 0.47 | * |
| 2009 | 173 | 50.87 | 1,191 | 49.29 | 0.51 | |
| 2010 | 161 | 50.31 | 938 | 42.22 | 0.54 | |
| 2011 | 166 | 53.61 | 849 | 29.68 | 0.64 | |
| 2012 | 164 | 56.10 | 335 | 30.15 | 0.65 | |
| 2013 | 149 | 62.42 | 170 | 30.59 | 0.67 | |
| 2014 | 126 | 67.46 | 294 | 27.55 | 0.71 | |
| 2015 | 126 | 79.37 | 523 | 66.92 | 0.54 | |
| 2016 | 118 | 44.92 | 340 | 66.47 | 0.40 | * |
| 2017 | 99 | 19.19 | 249 | 80.32 | 0.19 | * |
| 2018 | 138 | 23.91 | 220 | 86.82 | 0.22 | * |
| 2019 | 85 | 28.24 | 22 | 63.64 | 0.31 | * |
| 2020 | 43 | 69.77 | 26 | 34.62 | 0.67 | |
| 2021 | 89 | 84.27 | 75 | 66.67 | 0.56 | |

^a PNI = PNOB/(PNOB + PHOS).

PNOB = Percent natural origin fish in the hatchery broodstock.

PHOS = Percent hatchery origin fish among naturally spawning fish.

**Appendix J: Recoveries of Coded-Wire Tagged
Salmon Released into the Tucannon River for the
1985-2017 Brood Years**

Appendix J. Observed and estimated recoveries of coded-wire tagged salmon released into the Tucannon River with percent return to the Tucannon Basin, out-of-basin returns, and estimated survival and exploitation rates for the 1985-2017 brood years. (Data downloaded from RMIS database on 12/02/21.)

| Brood Year | 1985 | | 1986 | | 1987 | |
|----------------------------------|-----------------|------------------|---------------------|------------------|-----------------|------------------|
| Smolts Released | 12,922 | | 147,037 | | 151,100 | |
| Fish Size (g) | 76 | | 45 | | 50 | |
| CWT Codes^a | 34/42 | | 33/25, 41/46, 41/48 | | 49/50 | |
| Release Year | 1987 | | 1988 | | 1989 | |
| Agency (fishery/location) | Observed Number | Estimated Number | Observed Number | Estimated Number | Observed Number | Estimated Number |
| WDFW | | | | | | |
| Tucannon River | | | 30 | 84 | 28 | 130 |
| Kalama R., Wind R. | | | | | | |
| Treaty Troll | | | 1 | 2 | | |
| Lyons Ferry Hatch. ^b | 32 | 38 | 136 | 280 | 53 | 71 |
| F.W. Sport | | | 1 | 4 | | |
| ODFW | | | | | | |
| Test Net, Zone 4 | 1 | 1 | 1 | 1 | | |
| Treaty Ceremonial | | | 2 | 4 | 1 | 2 |
| Three Mile, Umatilla R. | | | | | | |
| Spawning Ground | | | | | | |
| Fish Trap - F.W. | | | | | | |
| F.W. Sport | | | | | | |
| Hatchery | | | | | | |
| CDFO | | | | | | |
| Non-treaty Ocean Troll | | | 1 | 4 | | |
| Mixed Net & Seine | | | | | | |
| Ocean Sport | | | | | | |
| USFWS | | | | | | |
| Warm Springs Hatchery | | | | | | |
| Dworshak NFH | | | | | | |
| IDFG | | | | | | |
| Hatchery | | | | | | |
| Total Returns | 33 | 39 | 172 | 379 | 82 | 203 |
| Tucannon (%) | 97.4 | | 96.0 | | 99.0 | |
| Out-of-Basin (%) | 0.0 | | 0.0 | | 0.0 | |
| Commercial Harvest (%) | 2.6 | | 1.8 | | 0.0 | |
| Sport Harvest (%) | 0.0 | | 1.1 | | 0.0 | |
| Treaty Ceremonial (%) | 0.0 | | 1.1 | | 1.0 | |
| Other (%) | 0.0 | | 0.0 | | 0.0 | |
| Survival | 0.30 | | 0.26 | | 0.13 | |

^a WDFW agency code prefix is 63.

^b Fish trapped at TFH and held at LFH for spawning.

Appendix J (continued). Observed and estimated recoveries of coded-wire tagged salmon released into the Tucannon River with percent return to the Tucannon Basin, out-of-basin returns, and estimated survival and exploitation rates for the 1985-2017 brood years. (Data downloaded from RMIS database on 12/02/21.)

| Brood Year | 1988 | | 1989 | | 1990 | |
|----------------------------------|-----------------|------------------|-----------------|------------------|---------------------|------------------|
| Smolts Released | 139,050 | | 97,779 | | 85,737 | |
| Fish Size (g) | 41 | | 50 | | 41 | |
| CWT Codes^a | 01/42, 55/01 | | 01/31, 14/61 | | 37/25, 40/21, 43/11 | |
| Release Year | 1990 | | 1991 | | 1992 | |
| Agency (fishery/location) | Observed Number | Estimated Number | Observed Number | Estimated Number | Observed Number | Estimated Number |
| WDFW | | | | | | |
| Tucannon River | 108 | 371 | 61 | 191 | 2 | 6 |
| Kalama R., Wind R. | | | | | | |
| Treaty Troll | | | 2 | 2 | | |
| Lyons Ferry Hatch. ^b | 83 | 86 | 55 | 55 | 19 | 19 |
| F.W. Sport | 1 | 4 | | | | |
| ODFW | | | | | | |
| Test Net, Zone 4 | 3 | 3 | 2 | 2 | | |
| Treaty Ceremonial | 8 | 17 | 4 | 8 | | |
| Three Mile, Umatilla R. | | | | | | |
| Spawning Ground | | | | | | |
| Fish Trap - F.W. | | | | | | |
| F.W. Sport | | | | | | |
| Hatchery | | | | | | |
| CDFO | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| Mixed Net & Seine | | | | | | |
| Ocean Sport | | | | | | |
| USFWS | | | | | | |
| Warm Springs Hatchery | | | | | | |
| Dworshak NFH | 1 | 1 | | | | |
| IDFG | | | | | | |
| Hatchery | | | | | | |
| Total Returns | 204 | 482 | 124 | 258 | 21 | 25 |
| Tucannon (%) | 94.8 | | 95.3 | | 100.0 | |
| Out-of-Basin (%) | 0.2 | | 0.0 | | 0.0 | |
| Commercial Harvest (%) | 0.6 | | 1.6 | | 0.0 | |
| Sport Harvest (%) | 0.8 | | 0.0 | | 0.0 | |
| Treaty Ceremonial (%) | 3.5 | | 3.1 | | 0.0 | |
| Other (%) | 0.0 | | 0.0 | | 0.0 | |
| Survival | 0.35 | | 0.26 | | 0.03 | |

^a WDFW agency code prefix is 63.

^b Fish trapped at TFH and held at LFH for spawning.

Appendix J (continued). Observed and estimated recoveries of coded-wire tagged salmon released into the Tucannon River with percent return to the Tucannon Basin, out-of-basin returns, and estimated survival and exploitation rates for the 1985-2017 brood years. (Data downloaded from RMIS database on 12/02/21.)

| Brood Year | 1991 | | 1992 | | 1992 | |
|----------------------------------|-----------------|------------------|---------------------|------------------|---------------------|------------------|
| Smolts Released | 72,461 | | 56,679 | | 79,151 | |
| Fish Size (g) | 30 | | 13 | | 32 | |
| CWT Codes^a | 46/25, 46/47 | | 48/23, 48/24, 48/56 | | 48/10, 48/55, 49/05 | |
| Release Year | 1993 | | 1993 | | 1994 | |
| Agency (fishery/location) | Observed Number | Estimated Number | Observed Number | Estimated Number | Observed Number | Estimated Number |
| WDFW | | | | | | |
| Tucannon River | | | | | 11 | 34 |
| Kalama R., Wind R. | | | | | | |
| Treaty Troll | | | | | | |
| Lyons Ferry Hatch. ^b | 24 | 24 | 2 | 2 | 45 | 47 |
| F.W. Sport | | | | | | |
| ODFW | | | | | | |
| Test Net, Zone 4 | | | | | | |
| Treaty Ceremonial | 1 | 3 | | | 1 | 1 |
| Three Mile, Umatilla R. | | | | | | |
| Spawning Ground | 1 | 1 | | | 2 | 2 |
| Fish Trap - F.W. | | | 1 | 1 | 5 | 9 |
| F.W. Sport | | | | | 2 | 2 |
| Hatchery | | | | | | |
| CDFO | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| Mixed Net & Seine | | | 1 | 2 | | |
| Ocean Sport | | | | | | |
| USFWS | | | | | | |
| Warm Springs Hatchery | | | | | 3 | 3 |
| Dworshak NFH | | | | | | |
| IDFG | | | | | | |
| Hatchery | | | | | | |
| Total Returns | 26 | 28 | 4 | 5 | 69 | 98 |
| Tucannon (%) | 85.7 | | 40.0 | | 82.7 | |
| Out-of-Basin (%) | 3.6 | | 20.0 | | 14.3 | |
| Commercial Harvest (%) | 0.0 | | 40.0 | | 0.0 | |
| Sport Harvest (%) | 0.0 | | 0.0 | | 2.0 | |
| Treaty Ceremonial (%) | 10.7 | | 0.0 | | 1.0 | |
| Other (%) | 0.0 | | 0.0 | | 0.0 | |
| Survival | 0.04 | | 0.01 | | 0.12 | |

^a WDFW agency code prefix is 63.

^b Fish trapped at TFH and held at LFH for spawning.

Appendix J (continued). Observed and estimated recoveries of coded-wire tagged salmon released into the Tucannon River with percent return to the Tucannon Basin, out-of-basin returns, and estimated survival and exploitation rates for the 1985-2017 brood years. (Data downloaded from RMIS database on 12/02/21.)

| Brood Year | 1993 | | 1994 | | 1995 | |
|----------------------------------|---------------------------|------------------|---------------------|------------------|---------------------|------------------|
| Smolts Released | 135,952 | | 130,034 | | 62,016 | |
| Fish Size (g) | 30-32 | | 25-35 | | 24-27 | |
| CWT Codes^a | 56/15, 56/17-18, 53/43-44 | | 43/23, 56/29, 57/29 | | 59/36, 61/40, 61/41 | |
| Release Year | 1995 | | 1996 | | 1997 | |
| Agency (fishery/location) | Observed Number | Estimated Number | Observed Number | Estimated Number | Observed Number | Estimated Number |
| WDFW | | | | | | |
| Tucannon River | 42 | 138 | 3 | 8 | 36 | 92 |
| Kalama R., Wind R. | | | | | | |
| Treaty Troll | | | | | | |
| Lyons Ferry Hatch. ^b | 66 | 66 | 21 | 21 | 94 | 94 |
| F.W. Sport | | | | | | |
| ODFW | | | | | | |
| Test Net, Zone 4 | | | | | | |
| Treaty Ceremonial | 3 | 3 | | | | |
| Three Mile, Umatilla R. | | | | | | |
| Spawning Ground | 3 | 3 | | | 1 | 1 |
| Fish Trap - F.W. | 1 | 1 | | | | |
| F.W. Sport | | | | | | |
| Hatchery | 1 | 1 | | | 1 | 1 |
| CDFO | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| Mixed Net & Seine | | | | | | |
| Ocean Sport | 1 | 3 | | | | |
| USFWS | | | | | | |
| Warm Springs Hatchery | | | | | | |
| Dworshak NFH | | | | | | |
| IDFG | | | | | | |
| Hatchery | | | | | | |
| Total Returns | 117 | 215 | 24 | 29 | 132 | 188 |
| Tucannon (%) | 94.9 | | 100.0 | | 98.9 | |
| Out-of-Basin (%) | 2.3 | | 0.0 | | 1.1 | |
| Commercial Harvest (%) | 0.0 | | 0.0 | | 0.0 | |
| Sport Harvest (%) | 1.4 | | 0.0 | | 0.0 | |
| Treaty Ceremonial (%) | 1.4 | | 0.0 | | 0.0 | |
| Other (%) | 0.0 | | 0.0 | | 0.0 | |
| Survival | 0.16 | | 0.02 | | 0.30 | |

^a WDFW agency code prefix is 63.

^b Fish trapped at TFH and held at LFH for spawning.

Appendix J (continued). Observed and estimated recoveries of coded-wire tagged salmon released into the Tucannon River with percent return to the Tucannon Basin, out-of-basin returns, and estimated survival and exploitation rates for the 1985-2017 brood years. (Data downloaded from RMIS database on 12/02/21.)

| Brood Year | 1996 | | 1997 | | 1998 | |
|----------------------------------|--------------------|------------------|-----------------|------------------|-----------------|------------------|
| Smolts Released | 76,028 | | 23,509 | | 124,093 | |
| Fish Size (g) | 28 | | 28 | | 35 | |
| CWT Codes^a | 03/59-60, 61/24-25 | | 61/32 | | 12/11 | |
| Release Year | 1998 | | 1999 | | 2000 | |
| Agency (fishery/location) | Observed Number | Estimated Number | Observed Number | Estimated Number | Observed Number | Estimated Number |
| WDFW | | | | | | |
| Tucannon River | 44 | 140 | 17 | 85 | 147 | 680 |
| Kalama R., Wind R. | | | | | | |
| Treaty Troll | | | | | | |
| Lyons Ferry Hatch. ^b | 96 | 99 | 44 | 46 | 83 | 83 |
| F.W. Sport | | | | | 3 | 14 |
| Non-treaty Ocean Troll | | | | | 1 | 2 |
| ODFW | | | | | | |
| Test Net, Zone 4 | | | | | 1 | 1 |
| Treaty Ceremonial | | | | | 5 | 5 |
| Three Mile, Umatilla R. | | | | | | |
| Spawning Ground | | | | | 1 | 1 |
| Fish Trap - F.W. | 1 | 1 | 2 | 2 | 8 | 10 |
| F.W. Sport | | | | | 2 | 4 |
| Hatchery | 2 | 2 | 1 | 1 | | |
| Columbia R. Gillnet | | | 7 | 22 | 32 | 85 |
| Columbia R. Sport | | | 2 | 15 | 17 | 94 |
| CDFO | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| Mixed Net & Seine | | | | | | |
| Ocean Sport | | | | | | |
| USFWS | | | | | | |
| Warm Springs Hatchery | | | | | | |
| Dworshak NFH | | | | | | |
| IDFG | | | | | | |
| Hatchery | 1 | 1 | 1 | 1 | | |
| Total Returns | 144 | 243 | 74 | 172 | 300 | 979 |
| Tucannon (%) | 98.4 | | 76.2 | | 77.9 | |
| Out-of-Basin (%) | 1.6 | | 2.3 | | 1.2 | |
| Commercial Harvest (%) | 0.0 | | 12.8 | | 9.0 | |
| Sport Harvest (%) | 0.0 | | 8.7 | | 11.4 | |
| Treaty Ceremonial (%) | 0.0 | | 0.0 | | 0.5 | |
| Other (%) | 0.0 | | 0.0 | | 0.0 | |
| Survival | 0.32 | | 0.73 | | 0.79 | |

^a WDFW agency code prefix is 63.

^b Fish trapped at TFH and held at LFH for spawning.

Appendix J (continued). Observed and estimated recoveries of coded-wire tagged salmon released into the Tucannon River with percent return to the Tucannon Basin, out-of-basin returns, and estimated survival and exploitation rates for the 1985-2017 brood years. (Data downloaded from RMIS database on 12/02/21.)

| Brood Year | 1999 | | 2000 | | 2001 | |
|----------------------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|
| Smolts Released | 96,736 | | 99,566 | | 144,013 | |
| Fish Size (g) | 43 | | 29 | | 35 | |
| CWT Codes^a | 02/75 | | 08/87 | | 06/81 | |
| Release Year | 2001 | | 2002 | | 2003 | |
| Agency (fishery/location) | Observed Number | Estimated Number | Observed Number | Estimated Number | Observed Number | Estimated Number |
| WDFW | | | | | | |
| Tucannon River | 2 | 12 | 13 | 37 | 6 | 26 |
| Kalama R., Wind R. | | | | | | |
| Treaty Troll | | | | | | |
| Lyons Ferry Hatch. ^b | 6 | 6 | 39 | 39 | 51 | 51 |
| F.W. Sport | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| ODFW | | | | | | |
| Test Net, Zone 4 | | | | | | |
| Treaty Ceremonial | | | | | | |
| Three Mile, Umatilla R. | | | | | | |
| Spawning Ground | | | | | | |
| Fish Trap - F.W. | | | | | | |
| F.W. Sport | | | | | | |
| Hatchery | | | | | | |
| Columbia R. Gillnet | 1 | 3 | 1 | 1 | | |
| Columbia R. Sport | | | | | | |
| CDFO | | | | | | |
| Non-treaty Ocean Troll | | | | | 1 | 5 |
| Mixed Net & Seine | | | | | | |
| Ocean Sport | | | | | | |
| USFWS | | | | | | |
| Warm Springs Hatchery | | | | | | |
| Dworshak NFH | | | | | | |
| IDFG | | | | | | |
| Hatchery | | | | | | |
| Total Returns | 9 | 21 | 53 | 77 | 58 | 82 |
| Tucannon (%) | 86.0 | | 98.7 | | 93.9 | |
| Out-of-Basin (%) | 0.0 | | 0.0 | | 0.0 | |
| Commercial Harvest (%) | 14.0 | | 1.3 | | 6.1 | |
| Sport Harvest (%) | 0.0 | | 0.0 | | 0.0 | |
| Treaty Ceremonial (%) | 0.0 | | 0.0 | | 0.0 | |
| Other (%) | 0.0 | | 0.0 | | 0.0 | |
| Survival | 0.02 | | 0.08 | | 0.06 | |

^a WDFW agency code prefix is 63.

^b Fish trapped at TFH and held at LFH for spawning.

Appendix J (continued). Observed and estimated recoveries of coded-wire tagged salmon released into the Tucannon River with percent return to the Tucannon Basin, out-of-basin returns, and estimated survival and exploitation rates for the 1985-2017 brood years. (Data downloaded from RMIS database on 12/02/21.)

| Brood Year | 2001 | | 2002 | | 2003 | |
|----------------------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|
| Smolts Released | 19,948 | | 121,774 | | 69,831 | |
| Fish Size (g) | 4 | | 39 | | 36 | |
| CWT Codes^a | 14/29 | | 17/91 | | 24/82 | |
| Release Year | 2002 | | 2004 | | 2005 | |
| Agency (fishery/location) | Observed Number | Estimated Number | Observed Number | Estimated Number | Observed Number | Estimated Number |
| WDFW | | | | | | |
| Tucannon River | | | 11 | 47 | 5 | 21 |
| Kalama R., Wind R. | | | | | | |
| Treaty Troll | | | | | | |
| Lyons Ferry Hatch. ^b | | | 58 | 58 | 21 | 21 |
| F.W. Sport | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| ODFW | | | | | | |
| Test Net, Zone 4 | | | | | | |
| Treaty Ceremonial | | | | | | |
| Three Mile, Umatilla R. | | | | | | |
| Spawning Ground | | | | | | |
| Fish Trap - F.W. | | | | | | |
| F.W. Sport | | | | | | |
| Hatchery | | | | | | |
| Columbia R. Gillnet | 1 | 1 | | | | |
| Columbia R. Sport | | | | | | |
| CDFO | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| Mixed Net & Seine | | | | | | |
| Ocean Sport | | | | | | |
| USFWS | | | | | | |
| Warm Springs Hatchery | | | | | | |
| Dworshak NFH | | | | | | |
| IDFG | | | | | | |
| Hatchery | | | | | | |
| Total Returns | 1 | 1 | 69 | 105 | 26 | 42 |
| Tucannon (%) | 0.0 | | 100.0 | | 100.0 | |
| Out-of-Basin (%) | 0.0 | | 0.0 | | 0.0 | |
| Commercial Harvest (%) | 100.0 | | 0.0 | | 0.0 | |
| Sport Harvest (%) | 0.0 | | 0.0 | | 0.0 | |
| Treaty Ceremonial (%) | 0.0 | | 0.0 | | 0.0 | |
| Other (%) | 0.0 | | 0.0 | | 0.0 | |
| Survival | 0.01 | | 0.09 | | 0.06 | |

^a WDFW agency code prefix is 63.

^b Fish trapped at TFH and held at LFH for spawning.

Appendix J (continued). Observed and estimated recoveries of coded-wire tagged salmon released into the Tucannon River with percent return to the Tucannon Basin, out-of-basin returns, and estimated survival and exploitation rates for the 1985-2017 brood years. (Data downloaded from RMIS database on 12/02/21.)

| Brood Year | 2003 | | 2004 | | 2004 | |
|----------------------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|
| Smolts Released | 125,304 | | 67,272 | | 127,162 | |
| Fish Size (g) | 34 | | 34 | | 30 | |
| CWT Codes^a | 27/78 CB | | 28/87 | | 28/65 CB | |
| Release Year | 2005 | | 2006 | | 2006 | |
| Agency (fishery/location) | Observed Number | Estimated Number | Observed Number | Estimated Number | Observed Number | Estimated Number |
| WDFW | | | | | | |
| Tucannon River | 5 | 21 | 24 | 102 | 17 | 73 |
| Kalama R., Wind R. | | | | | | |
| Treaty Troll | | | | | | |
| Lyons Ferry Hatch. ^b | 3 | 3 | 44 | 44 | 36 | 36 |
| F.W. Sport | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| ODFW | | | | | | |
| Test Net, Zone 4 | | | | | | |
| Treaty Ceremonial | | | | | | |
| Three Mile, Umatilla R. | | | | | | |
| Spawning Ground | | | | | | |
| Fish Trap - F.W. | | | | | | |
| F.W. Sport | | | | | | |
| Hatchery | | | | | | |
| Columbia R. Gillnet | | | | | 3 | 14 |
| Columbia R. Sport | | | | | 1 | 4 |
| CDFO | | | | | | |
| Non-treaty Ocean Troll | | | 1 | 1 | | |
| Mixed Net & Seine | | | | | | |
| Ocean Sport | | | | | | |
| USFWS | | | | | | |
| Warm Springs Hatchery | | | | | | |
| Dworshak NFH | | | | | | |
| IDFG | | | | | | |
| Hatchery | | | | | | |
| Total Returns | 8 | 24 | 69 | 147 | 57 | 127 |
| Tucannon (%) | 100.0 | | 99.3 | | 85.8 | |
| Out-of-Basin (%) | 0.0 | | 0.0 | | 0.0 | |
| Commercial Harvest (%) | 0.0 | | 0.7 | | 11.0 | |
| Sport Harvest (%) | 0.0 | | 0.0 | | 3.2 | |
| Treaty Ceremonial (%) | 0.0 | | 0.0 | | 0.0 | |
| Other (%) | 0.0 | | 0.0 | | 0.0 | |
| Survival | 0.02 | | 0.22 | | 0.10 | |

^a WDFW agency code prefix is 63.

^b Fish trapped at TFH and held at LFH for spawning.

Appendix J (continued). Observed and estimated recoveries of coded-wire tagged salmon released into the Tucannon River with percent return to the Tucannon Basin, out-of-basin returns, and estimated survival and exploitation rates for the 1985-2017 brood years. (Data downloaded from RMIS database on 12/02/21.)

| Brood Year | 2005 | | 2005 | | 2006 | |
|----------------------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|
| Smolts Released | 88,885 | | 144,833 | | 75,283 | |
| Fish Size (g) | 61 | | 57 | | 57 | |
| CWT Codes^a | 34/77 CB | | 35/99 | | 41/94 CB | |
| Release Year | 2007 | | 2007 | | 2008 | |
| Agency (fishery/location) | Observed Number | Estimated Number | Observed Number | Estimated Number | Observed Number | Estimated Number |
| WDFW | | | | | | |
| Tucannon River | 78 | 298 | 130 | 494 | 68 | 384 |
| Kalama R., Wind R. | | | | | | |
| Treaty Troll | | | | | | |
| Lyons Ferry Hatch. ^b | 3 | 3 | 96 | 97 | 4 | 5 |
| F.W. Sport | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| ODFW | | | | | | |
| Test Net, Zone 4 | | | 2 | 2 | | |
| Treaty Ceremonial | | | | | | |
| Three Mile, Umatilla R. | | | | | | |
| Spawning Ground | | | | | | |
| Fish Trap - F.W. | | | | | | |
| F.W. Sport | | | | | | |
| Hatchery | | | | | | |
| Columbia R. Gillnet | | | | | 8 | 26 |
| Columbia R. Sport | | | | | | |
| Juv. Marine Seine | 1 | 1 | | | 3 | 3 |
| CDFO | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| Mixed Net & Seine | | | | | | |
| Ocean Sport | | | | | | |
| USFWS | | | | | | |
| Warm Springs Hatchery | | | | | | |
| Dworshak NFH | | | | | | |
| IDFG | | | | | | |
| Hatchery | | | | | | |
| Total Returns | 82 | 302 | 228 | 593 | 83 | 418 |
| Tucannon (%) | 99.7 | | 99.7 | | 93.1 | |
| Out-of-Basin (%) | 0.0 | | 0.0 | | 0.0 | |
| Commercial Harvest (%) | 0.0 | | 0.3 | | 6.2 | |
| Sport Harvest (%) | 0.0 | | 0.0 | | 0.0 | |
| Treaty Ceremonial (%) | 0.0 | | 0.0 | | 0.0 | |
| Other (%) | 0.3 | | 0.0 | | 0.7 | |
| Survival | 0.34 | | 0.41 | | 0.56 | |

^a WDFW agency code prefix is 63.

^b Fish trapped at TFH and held at LFH for spawning.

Appendix J (continued). Observed and estimated recoveries of coded-wire tagged salmon released into the Tucannon River with percent return to the Tucannon Basin, out-of-basin returns, and estimated survival and exploitation rates for the 1985-2017 brood years. (Data downloaded from RMIS database on 12/02/21.)

| Brood Year | 2006 | | 2006 | | 2007 | |
|----------------------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|
| Smolts Released | 50,309 | | 51,858 | | 58,044 | |
| Fish Size (g) | 54 | | 39 | | 37 | |
| CWT Codes^a | 40/93 | | 40/94 | | 46/87 | |
| Release Year | 2008 | | 2008 | | 2009 | |
| Agency (fishery/location) | Observed Number | Estimated Number | Observed Number | Estimated Number | Observed Number | Estimated Number |
| WDFW | | | | | | |
| Tucannon River | 75 | 385 | 85 | 457 | 7 | 42 |
| Kalama R., Wind R. | | | | | | |
| Treaty Troll | | | | | | |
| Lyons Ferry Hatch. ^b | 42 | 75 | 48 | 87 | 31 | 31 |
| F.W. Sport | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| ODFW | | | | | | |
| Test Net, Zone 4 | | | | | | |
| Treaty Ceremonial | | | | | | |
| Three Mile, Umatilla R. | | | | | | |
| Spawning Ground | | | | | | |
| Fish Trap - F.W. | | | | | | |
| F.W. Sport | | | | | | |
| Hatchery | | | | | | |
| Columbia R. Gillnet | 5 | 21 | 2 | 9 | 1 | 5 |
| Columbia R. Sport | | | | | | |
| Juv. Marine Seine | 3 | 3 | 2 | 2 | | |
| CDFO | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| Mixed Net & Seine | | | | | | |
| Ocean Sport | | | | | | |
| USFWS | | | | | | |
| Warm Springs Hatchery | | | | | | |
| Dworshak NFH | | | | | | |
| IDFG | | | | | | |
| Hatchery | | | 1 | 1 | | |
| Total Returns | 125 | 484 | 138 | 556 | 39 | 78 |
| Tucannon (%) | 95.1 | | 97.8 | | 93.6 | |
| Out-of-Basin (%) | 0.0 | | 0.2 | | 0.0 | |
| Commercial Harvest (%) | 4.3 | | 1.6 | | 6.4 | |
| Sport Harvest (%) | 0.0 | | 0.0 | | 0.0 | |
| Treaty Ceremonial (%) | 0.0 | | 0.0 | | 0.0 | |
| Other (%) | 0.6 | | 0.4 | | 0.0 | |
| Survival | 0.96 | | 1.07 | | 0.13 | |

^a WDFW agency code prefix is 63.

^b Fish trapped at TFH and held at LFH for spawning.

Appendix J (continued). Observed and estimated recoveries of coded-wire tagged salmon released into the Tucannon River with percent return to the Tucannon Basin, out-of-basin returns, and estimated survival and exploitation rates for the 1985-2017 brood years. (Data downloaded from RMIS database on 12/02/21.)

| Brood Year | 2007 | | 2008 | | 2008 | |
|----------------------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|
| Smolts Released | 55,266 | | 84,613 | | 84,738 | |
| Fish Size (g) | 57 | | 40 | | 66 | |
| CWT Codes^a | 46/88 | | 51/74 | | 51/75 | |
| Release Year | 2009 | | 2010 | | 2010 | |
| Agency (fishery/location) | Observed Number | Estimated Number | Observed Number | Estimated Number | Observed Number | Estimated Number |
| WDFW | | | | | | |
| Tucannon River | 18 | 113 | 22 | 179 | 35 | 270 |
| Kalama R., Wind R. | | | | | | |
| Treaty Troll | | | | | | |
| Lyons Ferry Hatch. ^b | 32 | 32 | 28 | 28 | 49 | 49 |
| F.W. Sport | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| ODFW | | | | | | |
| Test Net, Zone 4 | | | | | | |
| Treaty Ceremonial | | | | | | |
| Three Mile, Umatilla R. | | | | | | |
| Spawning Ground | | | | | | |
| Fish Trap - F.W. | | | | | | |
| F.W. Sport | | | | | | |
| Hatchery | | | | | | |
| Columbia R. Gillnet | | | 1 | 4 | | |
| Columbia R. Sport | | | | | | |
| Juv. Marine Seine | | | | | | |
| CDFO | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| Mixed Net & Seine | | | | | | |
| Ocean Sport | | | | | | |
| USFWS | | | | | | |
| Warm Springs Hatchery | | | | | | |
| Dworshak NFH | | | | | | |
| IDFG | | | | | | |
| Hatchery | | | | | | |
| Total Returns | 50 | 145 | 51 | 211 | 84 | 319 |
| Tucannon (%) | 100.0 | | 98.1 | | 100.0 | |
| Out-of-Basin (%) | 0.0 | | 0.0 | | 0.0 | |
| Commercial Harvest (%) | 0.0 | | 1.9 | | 0.0 | |
| Sport Harvest (%) | 0.0 | | 0.0 | | 0.0 | |
| Treaty Ceremonial (%) | 0.0 | | 0.0 | | 0.0 | |
| Other (%) | 0.0 | | 0.0 | | 0.0 | |
| Survival | 0.26 | | 0.25 | | 0.38 | |

^a WDFW agency code prefix is 63.

^b Fish trapped at TFH and held at LFH for spawning.

Appendix J (continued). Observed and estimated recoveries of coded-wire tagged salmon released into the Tucannon River with percent return to the Tucannon Basin, out-of-basin returns, and estimated survival and exploitation rates for the 1985-2017 brood years. (Data downloaded from RMIS database on 12/02/21.)

| Brood Year | 2009 | | 2009 | | 2010 | |
|----------------------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|
| Smolts Released | 117,824 | | 113,049 | | 102,169 | |
| Fish Size (g) | 35 | | 51 | | 32 | |
| CWT Codes^a | 55/65 | | 55/66 | | 60/75 | |
| Release Year | 2011 | | 2011 | | 2012 | |
| Agency (fishery/location) | Observed Number | Estimated Number | Observed Number | Estimated Number | Observed Number | Estimated Number |
| WDFW | | | | | | |
| Tucannon River | 4 | 88 | 5 | 125 | 10 | 115 |
| Kalama R., Wind R. | | | | | | |
| Treaty Troll | | | | | | |
| Lyons Ferry Hatch. ^b | 16 | 16 | 40 | 40 | 17 | 17 |
| F.W. Sport | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| Lower Granite Trap | | | 1 | 1 | | |
| ODFW | | | | | | |
| Test Net, Zone 4 | | | | | | |
| Treaty Ceremonial | | | | | | |
| Three Mile, Umatilla R. | | | | | | |
| Spawning Ground | | | | | | |
| Fish Trap - F.W. | | | | | | |
| F.W. Sport | | | | | | |
| Hatchery | | | | | | |
| Columbia R. Gillnet | | | 1 | 2 | | |
| Columbia R. Sport | | | | | | |
| Juv. Marine Seine | | | | | | |
| CDFO | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| Mixed Net & Seine | | | | | | |
| Ocean Sport | 1 | 4 | | | | |
| USFWS | | | | | | |
| Warm Springs Hatchery | | | | | | |
| Dworshak NFH | | | | | | |
| NMFS | | | | | | |
| Juvenile Trawl Sample | | | | | 1 | 1 |
| Total Returns | 21 | 108 | 47 | 168 | 28 | 133 |
| Tucannon (%) | 96.3 | | 98.2 | | 99.2 | |
| Out-of-Basin (%) | 0.0 | | 0.6 | | 0.0 | |
| Commercial Harvest (%) | 0.0 | | 1.2 | | 0.0 | |
| Sport Harvest (%) | 3.7 | | 0.0 | | 0.0 | |
| Treaty Ceremonial (%) | 0.0 | | 0.0 | | 0.0 | |
| Other (%) | 0.0 | | 0.0 | | 0.8 | |
| Survival | 0.09 | | 0.15 | | 0.13 | |

^a WDFW agency code prefix is 63.

^b Fish trapped at TFH and held at LFH for spawning.

Appendix J (continued). Observed and estimated recoveries of coded-wire tagged salmon released into the Tucannon River with percent return to the Tucannon Basin, out-of-basin returns, and estimated survival and exploitation rates for the 1985-2017 brood years. (Data downloaded from RMIS database on 12/02/21.)

| Brood Year | 2010 | | 2011 | | 2011 | |
|----------------------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|
| Smolts Released | 96,984 | | 227,703 | | 27,748 | |
| Fish Size (g) | 66 | | 33 | | 33 | |
| CWT Codes^a | 60/76 | | 64/41 | | 64/42 | |
| Release Year | 2012 | | 2013 | | 2013 | |
| Agency (fishery/location) | Observed Number | Estimated Number | Observed Number | Estimated Number | Observed Number | Estimated Number |
| WDFW | | | | | | |
| Tucannon River | 10 | 122 | 92 | 673 | 5 | 36 |
| Kalama R., Wind R. | | | | | | |
| Treaty Troll | | | | | | |
| Lyons Ferry Hatch. ^b | 22 | 22 | 27 | 27 | 2 | 2 |
| F.W. Sport | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| Lower Granite Trap | | | 1 | 1 | | |
| ODFW | | | | | | |
| Test Net, Zone 4 | | | | | | |
| Treaty Ceremonial | | | | | | |
| Three Mile, Umatilla R. | | | | | | |
| Spawning Ground | | | | | | |
| Fish Trap - F.W. | | | | | | |
| F.W. Sport | | | | | | |
| Hatchery | | | | | | |
| Columbia R. Gillnet | | | 4 | 19 | | |
| Columbia R. Sport | | | | | | |
| Juv. Marine Seine | | | | | | |
| Non-treaty Ocean Troll | | | 1 | 4 | | |
| CDFO | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| Mixed Net & Seine | | | | | | |
| Ocean Sport | | | | | | |
| USFWS | | | | | | |
| Warm Springs Hatchery | | | | | | |
| Dworshak NFH | | | | | | |
| IDFG | | | | | | |
| Hatchery | | | | | | |
| Total Returns | 32 | 144 | 125 | 724 | 7 | 38 |
| Tucannon (%) | 100.0 | | 96.7 | | 100.0 | |
| Out-of-Basin (%) | 0.0 | | 0.1 | | 0.0 | |
| Commercial Harvest (%) | 0.0 | | 3.2 | | 0.0 | |
| Sport Harvest (%) | 0.0 | | 0.0 | | 0.0 | |
| Treaty Ceremonial (%) | 0.0 | | 0.0 | | 0.0 | |
| Other (%) | 0.0 | | 0.0 | | 0.0 | |
| Survival | 0.15 | | 0.32 | | 0.14 | |

^a WDFW agency code prefix is 63.

^b Fish trapped at TFH and held at LFH for spawning.

Appendix J (continued). Observed and estimated recoveries of coded-wire tagged salmon released into the Tucannon River with percent return to the Tucannon Basin, out-of-basin returns, and estimated survival and exploitation rates for the 1985-2017 brood years. (Data downloaded from RMIS database on 12/02/21.)

| Brood Year | 2012 | | 2012 | | 2013 | |
|----------------------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|
| Smolts Released | 179,400 | | 21,101 | | 179,494 | |
| Fish Size (g) | 32 | | 32 | | 37 | |
| CWT Codes^a | 65/85 | | 65/86 | | 67/42 | |
| Release Year | 2014 | | 2014 | | 2015 | |
| Agency (fishery/location) | Observed Number | Estimated Number | Observed Number | Estimated Number | Observed Number | Estimated Number |
| WDFW | | | | | | |
| Tucannon River | 96 | 406 | 7 | 36 | 108 | 233 |
| Kalama R., Wind R. | | | | | | |
| Treaty Troll | | | | | | |
| Lyons Ferry Hatch. ^b | 56 | 58 | 3 | 3 | 85 | 85 |
| F.W. Sport | | | | | | |
| Non-treaty Ocean Troll | 1 | 1 | | | 2 | 4 |
| ODFW | | | | | | |
| Test Net, Zone 4 | | | | | 1 | 1 |
| Treaty Ceremonial | | | | | | |
| Three Mile, Umatilla R. | | | | | | |
| Spawning Ground | | | | | | |
| Fish Trap - F.W. | | | | | | |
| F.W. Sport | | | | | | |
| Hatchery | 1 | 1 | | | | |
| Columbia R. Gillnet | | | | | | |
| Columbia R. Sport | | | | | | |
| Juv. Marine Seine | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| CDFO | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| Mixed Net & Seine | | | | | | |
| Ocean Sport | | | | | | |
| USFWS | | | | | | |
| Warm Springs Hatchery | | | | | | |
| Dworshak NFH | | | | | | |
| NMFS | | | | | | |
| Juvenile Trawl Sample | 1 | 1 | | | 1 | 1 |
| Total Returns | 155 | 467 | 10 | 39 | 197 | 324 |
| Tucannon (%) | 99.4 | | 100.0 | | 98.2 | |
| Out-of-Basin (%) | 0.2 | | 0.0 | | 0.0 | |
| Commercial Harvest (%) | 0.2 | | 0.0 | | 1.2 | |
| Sport Harvest (%) | 0.0 | | 0.0 | | 0.0 | |
| Treaty Ceremonial (%) | 0.0 | | 0.0 | | 0.0 | |
| Other (%) | 0.2 | | 0.0 | | 0.6 | |
| Survival | 0.26 | | 0.18 | | 0.18 | |

^a WDFW agency code prefix is 63.

^b Fish trapped at TFH and held at LFH for spawning.

Appendix J (continued). Observed and estimated recoveries of coded-wire tagged salmon released into the Tucannon River with percent return to the Tucannon Basin, out-of-basin returns, and estimated survival and exploitation rates for the 1985-2017 brood years. (Data downloaded from RMIS database on 12/02/21.)

| Brood Year | 2013 | | 2014 | | 2015 | |
|----------------------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|
| Smolts Released | 20,373 | | 216,295 | | 187,601 | |
| Fish Size (g) | 37 | | 40 | | 40 | |
| CWT Codes^a | 67/43 | | 68/84 | | 70/39 | |
| Release Year | 2015 | | 2016 | | 2017 | |
| Agency (fishery/location) | Observed Number | Estimated Number | Observed Number | Estimated Number | Observed Number | Estimated Number |
| WDFW | | | | | | |
| Tucannon River | 15 | 20 | 155 | 304 | 10 | 65 |
| Kalama R., Wind R. | | | | | | |
| Treaty Troll | | | | | | |
| Lyons Ferry Hatch. ^b | 6 | 6 | 141 | 142 | 111 | 111 |
| F.W. Sport | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| ODFW | | | | | | |
| Test Net, Zone 4 | | | | | | |
| Treaty Ceremonial | | | | | | |
| Three Mile, Umatilla R. | | | | | | |
| Spawning Ground | | | | | | |
| Fish Trap - F.W. | | | | | 1 | 1 |
| F.W. Sport | | | | | | |
| Hatchery | | | | | | |
| Columbia R. Gillnet | 1 | 5 | 1 | 1 | | |
| Columbia R. Sport | | | | | | |
| Juv. Marine Seine | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| CDFO | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| Mixed Net & Seine | | | | | | |
| Ocean Sport | | | | | | |
| USFWS | | | | | | |
| Warm Springs Hatchery | | | | | | |
| Dworshak NFH | | | | | | |
| NMFS | | | | | | |
| Juvenile Trawl Sample | | | 1 | 1 | | |
| Total Returns | 22 | 31 | 298 | 448 | 122 | 177 |
| Tucannon (%) | 83.9 | | 99.6 | | 99.4 | |
| Out-of-Basin (%) | 0.0 | | 0.0 | | 0.6 | |
| Commercial Harvest (%) | 16.1 | | 0.2 | | 0.0 | |
| Sport Harvest (%) | 0.0 | | 0.0 | | 0.0 | |
| Treaty Ceremonial (%) | 0.0 | | 0.0 | | 0.0 | |
| Other (%) | 0.0 | | 0.2 | | 0.0 | |
| Survival | 0.15 | | 0.21 | | 0.09 | |

^a WDFW agency code prefix is 63.

^b Fish trapped at TFH and held at LFH for spawning.

Appendix J (continued). Observed and estimated recoveries of coded-wire tagged salmon released into the Tucannon River with percent return to the Tucannon Basin, out-of-basin returns, and estimated survival and exploitation rates for the 1985-2017 brood years. (Data downloaded from RMIS database on 12/02/21.)

| Brood Year | 2016 | | 2017 | | | |
|----------------------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|
| Smolts Released | 202,952 | | 140,262 | | | |
| Fish Size (g) | 55 | | 30 | | | |
| CWT Codes^a | 72/01 | | 73/96 | | | |
| Release Year | 2018 | | 2019 | | | |
| Agency (fishery/location) | Observed Number | Estimated Number | Observed Number | Estimated Number | Observed Number | Estimated Number |
| WDFW | | | | | | |
| Tucannon River | 9 | 9 | | | | |
| Kalama R., Wind R. | | | | | | |
| Treaty Troll | | | | | | |
| Lyons Ferry Hatch. ^b | 13 | 13 | 2 | 2 | | |
| F.W. Sport | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| ODFW | | | | | | |
| Test Net, Zone 4 | | | | | | |
| Treaty Ceremonial | | | | | | |
| Three Mile, Umatilla R. | | | | | | |
| Spawning Ground | | | | | | |
| Fish Trap - F.W. | | | | | | |
| F.W. Sport | | | | | | |
| Hatchery | | | | | | |
| Columbia R. Gillnet | | | | | | |
| Columbia R. Sport | | | | | | |
| Juv. Marine Seine | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| CDFO | | | | | | |
| Non-treaty Ocean Troll | | | | | | |
| Mixed Net & Seine | | | | | | |
| Ocean Sport | | | | | | |
| USFWS | | | | | | |
| Warm Springs Hatchery | | | | | | |
| Dworshak NFH | | | | | | |
| NMFS | | | | | | |
| Juvenile Trawl Sample | | | | | | |
| Total Returns | 22 | 22 | 2 | 2 | | |
| Tucannon (%) | 100.0 | | 100.0 | | | |
| Out-of-Basin (%) | 0.0 | | 0.0 | | | |
| Commercial Harvest (%) | 0.0 | | 0.0 | | | |
| Sport Harvest (%) | 0.0 | | 0.0 | | | |
| Treaty Ceremonial (%) | 0.0 | | 0.0 | | | |
| Other (%) | 0.0 | | 0.0 | | | |
| Survival | 0.01 | | 0.00 | | | |

^a WDFW agency code prefix is 63.

^b Fish trapped at TFH and held at LFH for spawning.



This program receives Federal financial assistance from the U.S. Fish and Wildlife Service Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972. The U.S. Department of the Interior and its bureaus prohibit discrimination on the bases of race, color, national origin, age, disability and sex (in educational programs). If you believe that you have been discriminated against in any program, activity or facility, please contact the WDFW ADA Program Manager at P.O. Box 43139, Olympia, Washington 98504, or write to

Department of the Interior
Chief, Public Civil Rights Division
1849 C Street NW
Washington D.C. 20240