


STATE OF WASHINGTON

JUNE 2024

# Upper Chehalis River Smolt Production, 2022



by Devin West, Justin Miller-Nelson,  
and Marisa Litz



Washington  
Department of  
**FISH &  
WILDLIFE**

**FPA 24-07**

## Upper Chehalis River Smolt Production, 2022



### **Washington Department of Fish and Wildlife**

Devin West, Justin Miller-Nelson, and Marisa Litz  
Fish Science Division  
1111 Washington Street SE, Olympia WA 98501

## **Acknowledgements**

We would like to thank Charlotte Gruninger for field operations, Eric Walther for providing maps in this report, and the WDFW Ageing and Genetics labs for processing our samples. We would also like to thank Washington Department of Transportation, Washington State Parks, and Lewis County for allowing us to access to our trapping site.

This project was funded by the Washington State Legislature through the Department of Ecology's Office of Chehalis Basin.

**Recommended citation:** West, D., J. Miller-Nelson, and M. Litz. 2024. Upper Chehalis River Smolt Production, 2022, FPA 24-07. Washington Department of Fish and Wildlife, Olympia, Washington.

# Table of Contents

<b>ACKNOWLEDGEMENTS .....</b>	<b>I</b>
<b>TABLE OF CONTENTS .....</b>	<b>II</b>
<b>LIST OF TABLES.....</b>	<b>III</b>
<b>LIST OF FIGURES.....</b>	<b>IV</b>
<b>LIST OF APPENDICES .....</b>	<b>V</b>
<b>EXECUTIVE SUMMARY .....</b>	<b>1</b>
<b>INTRODUCTION .....</b>	<b>3</b>
OBJECTIVES .....	3
<b>METHODS.....</b>	<b>4</b>
STUDY SITE .....	4
TRAP OPERATION .....	6
FISH COLLECTION.....	7
TRAP EFFICIENCY TRIALS.....	8
ASSUMPTION TESTING.....	9
ANALYSIS.....	10
<b>RESULTS.....</b>	<b>11</b>
SUMMARY OF FISH SPECIES ENCOUNTERED .....	11
TRAP OPERATION.....	11
ASSUMPTION TESTING TRIALS .....	11
CHINOOK .....	11
COHO.....	15
STEELHEAD .....	20
<b>DISCUSSION.....</b>	<b>24</b>
BASIN-WIDE CONTEXT .....	24
NEXT STEPS.....	27
<b>REFERENCES .....</b>	<b>29</b>
<b>APPENDICES.....</b>	<b>31</b>

## List of Tables

Table 1. Abundance of Chinook, coho, and steelhead outmigrants that completed their freshwater rearing phase upstream of river kilometer 151.7 (river mile 94.3) of the Upper Chehalis River.....	2
Table 2. Sample rates for biological data collection from wild juvenile salmonids. ....	7
Table 3. Date and length criteria used for field calls of juvenile Chinook salmon. ....	8
Table 4. Date and length criteria used for field calls of juvenile coho salmon. ....	8
Table 5. Date and length criteria used for field calls of juvenile steelhead trout. ....	8
Table 6. Abundance estimate groups defined by species, origin, life stage, and age class. Life stages included in the estimates were transitional (T), and smolt (S). Age classes included in the estimates were subyearling (SY) and yearling (Y). FL = Fork length. ....	9
Table 7. Trap efficiency marks and release locations for each abundance estimate group. Efficiency marks are visible implant elastomer tag (VIE) and passive integrated transponder tag (PIT). ....	9
Table 8. Freshwater ages of wild coho outmigrants (transitionals, smolts) at the Upper Chehalis River screw trap, 2022. Data are scale ages of sampled juveniles by week. ....	20
Table 9. Freshwater ages of wild steelhead outmigrants (transitionals, smolts) at the Upper Chehalis River screw trap, 2022. Data are scale ages of sampled juveniles by week. ....	24
Table 10. Mean monthly stream temperatures °C recorded at Upper Chehalis River smolt trap near river km 151.7, 2022. ....	27

## List of Figures

Figure 1. Upper Chehalis main stem rotary screw trap (46.634747, -123.167972). Anadromous streams represent stream habitat within the predicted coho salmon range of occurrence (299.8 km) using a 0.50 probability decision threshold upstream of the upper Chehalis River rotary screw trap. Non-anadromous streams represent stream habitat outside the predicted coho salmon range of occurrence (919.4 km) upstream of the trap location (Walther 2021). .....	5
Figure 2. Upper Chehalis River trap site and fishing positions. ....	6
Figure 3. Number of outmigrants (top panel) and trap efficiency (bottom panel) by week for wild Chinook subyearlings produced above the Upper Chehalis River smolt trap in 2022. The total estimate is 19,870 with a coefficient of variation (CV) of 7.2%. Error bars and shading around point estimates represent 95% confidence intervals. ....	13
Figure 4. Box plots of fork lengths of wild Chinook subyearling outmigrants (transitionals, smolts) by week at the Upper Chehalis River screw trap, 2022. Boxes represent the median, first and third quartiles, whiskers represent the interquartile ranges, and dots represent outliers. ....	14
Figure 5. Chinook wild transitional and smolt trap efficiency (top), maiden catch (bottom) and flow in cubic feet per second (cfs top & bottom), with trap outages (red) by date at the Upper Chehalis smolt trap in 2022. ....	15
Figure 6. Number of outmigrants (top panel) and trap efficiency (bottom panel) by week for wild coho smolts and transitionals produced above the Upper Chehalis River smolt trap in 2022. The total estimate is 24,434 with a coefficient of variation (CV) of 7.6%. Error bars and shading around point estimates represent 95% confidence intervals. ....	17
Figure 7. Plot of length and age by date for wild coho outmigrants (transitionals, smolts) at the Upper Chehalis River screw trap, 2022. ....	18
Figure 8. Coho wild transitional and smolt trap raw efficiency (top), maiden catch (bottom) and flow in cubic feet per second (cfs, top & bottom) with trap outages (red) by date at the Upper Chehalis smolt trap in 2022. ....	19
Figure 9. Number of outmigrants (top panel) and trap efficiency (bottom panel) by week for wild steelhead smolts and transitionals produced above the Upper Chehalis River smolt trap in 2022. The total estimate is 10,204 with a CV of 19.6%. Error bars and shading around point estimates represent 95% confidence intervals. ....	21
Figure 10. Plot of length and age data by date for wild steelhead outmigrants (transitionals, smolts) at the Upper Chehalis River screw trap, 2022. ....	22
Figure 11. Steelhead wild transitional and smolt trap efficiency (top), maiden catch (bottom) and flow in cubic feet per second (cfs, top & bottom) with trap outages (red) by date at Upper Chehalis smolt trap in 2022. ....	23
Figure 12. Chinook maiden catch and average daily stream temperature (°C) at the Upper Chehalis River smolt trap, 2022. ....	28

## List of Appendices

Appendix A. Decision tree for assigning life stages of juvenile outmigrants developed by the Washington Department of Fish and Wildlife to ensure consistency in data collection protocols across juvenile trapping projects.....	31
Appendix B. Upper Chehalis River missed trapping periods, 2022. ....	32
Appendix C. Mark-recapture data for wild Chinook outmigrants (transitionals, smolts) organized by time period. Dataset includes total marks released (Total Mark), total marks recaptures (Total Recap), total maiden captures (Total Captures), and the proportion of time the trap fished during the time period (Prop Fished). ....	33
Appendix D. Mark-recapture data for wild coho outmigrants (transitionals, smolts) organized by time period. Data are the combined counts of subyearling and yearling coho. Dataset includes total marks released (Total Mark), total marks recaptures (Total Recap), total maiden captures (Total Captures), and the proportion of time the trap fished during the time period (Prop Fished). ....	34
Appendix E. Mark-recapture data for wild Steelhead outmigrants (transitionals, smolts) organized by time period. Dataset includes total marks released (Total Mark), total marks recaptures (Total Recap), total maiden captures (Total Captures), and the proportion of time the trap fished during the time period (Prop Fished). ....	35

## Executive Summary

This report provides the second year of results from a juvenile salmonid monitoring study on the Upper Chehalis River main stem near Pe Ell, Washington in 2022 (Table 1). The primary objective of this study is to describe the freshwater production (e.g., smolt abundance) of Pacific salmon (*Oncorhynchus* spp.) and steelhead trout (*O. mykiss*) in the Upper Chehalis River. Specifically, we describe the abundance, timing, and diversity (body size, age structure) of juvenile outmigrants for wild Chinook (*O. tshawytscha*), coho salmon (*O. kisutch*), and steelhead trout. Based on the location and timing of our study, the results reflect juveniles that completed their freshwater rearing phase in habitats upstream of river kilometer (rkm) 151.7 (river mile 94.3) of the main stem Chehalis River.

To meet the study objectives, a 1.5-meter (5-foot) rotary screw trap was operated near rkm 151.7 (river mile 94.3) of the main stem Chehalis River from March 10 to July 11, 2022.

The Chinook salmon outmigrants quantified were subyearlings, not fry. The majority of Chinook fry ( $\leq 45$  mm fork length) outmigrate when flow conditions are not suitable for smolt trapping in the Chehalis River (e.g., January and February). Therefore, the goal was to estimate the subyearling ( $>45$  mm fork length) component of the Chinook outmigration that generally occurs from March through July. Fork length of Chinook subyearlings increased steadily throughout the trapping period with an average of 50.2 mm ( $\pm 4.5$  mm, standard deviation SD) and 101.8 mm ( $\pm 8.1$  mm SD) in the first and last sampled week of trapping, respectively. During this time roughly 95% of the total catch of wild Chinook outmigrants were ( $>45$  mm). Abundance of wild Chinook subyearling outmigrants was estimated to be 19,870 (95% confidence intervals, CI = 17,383–22,951) with a coefficient of variation (CV) of 7.2%.

The coho outmigrants quantified were yearlings. Scale age data indicated the presence of three age classes, zero-, one-, and two-year-olds. The average fork length of age zero outmigrants was 108.3 mm ( $\pm 4$  mm SD) and age one yearling outmigrants averaged 118 mm ( $\pm 10.5$  mm SD) and age two yearling outmigrants averaged 123.6 mm ( $\pm 12.4$  mm SD). Abundance of wild coho outmigrants was estimated to be 24,434 (95% CI = 21,175–28,403) with a CV of 7.6%.

Steelhead outmigrants were one, two, and three years of age. Fork length averaged 154.5 mm ( $\pm 14.1$  mm SD) for one-year olds, 160.3 mm ( $\pm 14.1$  mm SD) for two-year olds, and 169 mm ( $\pm 14.8$  mm SD) for three-year olds. Abundance of wild steelhead outmigrants was estimated to be 10,204 (95% CI = 7,261–15,252) with a CV of 19.6%.



Table 1. Abundance of Chinook, coho, and steelhead outmigrants that completed their freshwater rearing phase upstream of river kilometer 151.7 (river mile 94.3) of the Upper Chehalis River.

Abundance Group	Origin	Life Stage	Age Class	Abundance (95% Confidence Intervals)	Coefficient of Variation (%)
Chinook	Wild	Transitional, Smolt	Subyearling	19,870 (17,383–22,951)	7.2
Coho	Wild	Transitional, Smolt	Subyearling, Yearling	24,434 (21,175–28,403)	7.6
Steelhead	Wild	Transitional, Smolt	Yearling	10,204 (7,261–15,252)	19.6

## Introduction

The Washington Department of Fish and Wildlife (WDFW) has monitored freshwater production of juvenile Pacific salmon (*Oncorhynchus* spp.) in the Chehalis River since the early 1980s. Over this time, the work has focused on generating wild coho salmon (*O. kisutch*) estimates of smolt abundance at a basin scale. Results from this monitoring program have demonstrated that the Chehalis River has a higher density of wild coho smolts (average 1,011 smolts mi<sup>-2</sup> or 390 smolts km<sup>-2</sup>) than any other western Washington watershed for which data currently exists (Litz 2023). In the 1980s and 1990s, smolt abundance estimates from individual tributaries throughout the Chehalis River were also generated, however, prior to 2019, smolt abundance estimates had not been evaluated for nearly two decades. Furthermore, because the current method for basin scale population estimation utilizes back calculation, estimates are not readily available until returning adults are sampled for coded wire tags (CWT) approximately 18 months following outmigration. Therefore, there is limited information on freshwater production of other salmonid species, including Chinook (*O. tshawytscha*) and chum salmon (*O. keta*) and steelhead trout (*O. mykiss*) in the Chehalis River basin. Recent efforts under the Chehalis Basin Strategy (<http://chehalisbasinstrategy.com/>) to develop a monitoring and adaptive management plan (M&AMT 2021) as part of the larger Aquatic Species Restoration Plan (ASRPSC 2019) have highlighted the need for annual smolt (or juvenile outmigrant) data that will be critical for evaluating variability and trends in freshwater production over time in response to freshwater restoration.

Smolt monitoring activities by WDFW were expanded in 2019 to develop a more comprehensive understanding of freshwater production among multiple species of salmonids across different ecological regions in the Chehalis River basin (e.g., Olympic and Cascade mountains, Willapa Hills). Beginning in 2021, this expanded effort became a long-term component of an integrated status and trends monitoring program used to evaluate salmon and steelhead abundance in the riverine environment in response to habitat restoration, protection actions, and environmental change (M&AMT 2021). Also in 2021, the Upper Chehalis River was selected as an area to monitor smolt production, collect baseline information to better inform restoration projects, and evaluate potential impacts of a proposed Flood Retention Expandable (FRE) facility in the basin. The Upper Chehalis River supports runs of fall and spring run Chinook salmon, coho salmon, and steelhead trout. In fact, the Upper Chehalis River is known to support a relatively large proportion (~15%) of the steelhead population in the entire Chehalis River Basin (Ronne et al. 2020). Additionally, the proposed location for a FRE facility in the main stem Chehalis River at river kilometer (rkm) 174 (river mile 108.2) has highlighted the need for research to fill data gaps about species composition, abundance, distribution, and life history diversity. For these reasons, accurate and unbiased estimates of juvenile salmon and steelhead abundance (e.g., freshwater production) in the Upper Chehalis River are critical for monitoring status and trends of salmon and steelhead populations and their response to habitat alterations.

### Objectives

The primary objective of this study was to estimate the freshwater production of salmon and steelhead in the Upper Chehalis River. Specifically, goals were to describe the abundance, timing, and diversity (body size, age structure) of juvenile outmigrants for wild Chinook salmon, coho salmon, and steelhead. Based on the location and timing of the study, results reflect juveniles that

completed their freshwater rearing phase in habitats upstream of rkm 151.7 (river mile 94.3) of the main stem Upper Chehalis River. This report includes results from the second field season in 2022.

## Methods

### *Study Site*

The Chehalis River is a large coastal watershed in western Washington that drains approximately 6,889 km<sup>2</sup> from the Willapa Hills, Cascade Mountains, and Olympic Mountains into Grays Harbor. The Upper Chehalis River sub-basin has a rain dominant hydrology and arises in the East Fork and West Fork. Primary tributaries to the Upper Chehalis include Thrash, Crim, Rock, and Elk creeks. Land use in the sub-basin is predominately timber production in headwater locations and private residential and agricultural in lower elevation locations. Timber lands are often characterized by steep sloped banks and drainages. River flows in the sub-basin can fluctuate annually from ~18,000 cubic feet per second (cfs or 510 m<sup>3</sup>s<sup>-1</sup>) down to ~20 cfs (0.6 m<sup>3</sup>s<sup>-1</sup>) with sudden and abrupt changes in flows being common. Native anadromous salmonids in the Chehalis River include fall and spring Chinook salmon, coho salmon, winter steelhead, and cutthroat trout (*O. clarkii*). Chum salmon are present in the basin but occur downstream of the smolt trap location in this study. A WDFW acclimation pond is located on Eight Creek, a tributary to Elk Creek. This pond is located upstream of the trapping site and releases approximately 100,000 adipose fin marked coho and 25,000 to 30,000 adipose fin marked steelhead from Skookumchuck hatchery annually (M. Scharpf WDFW, personal communication). Juvenile research is also conducted annually approximately 25 km upstream of the trap site by the Chehalis tribe operating a 5 ft rotary screw trap. Further downstream of this site, rotary screw traps are also operated by WDFW in the Newaukum River at rkm 9.35 and in the main stem Chehalis River at rkm 84.

Like other rivers in western Washington, juvenile Chinook salmon in the Chehalis River migrate downstream over a protracted outmigration period during their first year of life. Yearlings are rarely observed at the Chehalis main stem smolt trap or in the adult returns as determined from otoliths (Campbell et al. 2017; Olson et al. 2023; West et al. 2021). The Chehalis main stem trap is downstream of the Upper Chehalis trap, therefore juvenile Chinook salmon in the Upper Chehalis presumably exhibit a similar life history observed lower in the system. There are two predominant freshwater rearing strategies observed for juvenile Chinook salmon and these are both documented at the Chehalis main stem smolt trap as a bimodal outmigration. The first pulse of outmigrants represent ‘fry’ (defined as juveniles  $\leq 45$  mm fork length, FL), which are individuals that out-migrate almost immediately after emergence. Fry are observed at the smolt trap beginning in mid-March but have been presumably out-migrating since January, based on other fry and smolt traps in the Chehalis River, Puget Sound, and other areas (Anderson and Topping 2018; Gilbertson et al. 2021; Groot and Margolis 1991; Kiyohara and Zimmerman 2012; Zimmerman et al. 2015). The second pulse of Chinook outmigrants represent ‘subyearlings’, which are individuals  $>45$  mm FL that grow in freshwater for weeks to months after emergence and are observed at the smolt trap between the months of April and July.

The trapping location on the Upper Chehalis River (46°38’5.06 N, 123°10’4.47 W) is located at rkm 151.7, approximately 22.4 km downstream of a proposed FRE facility and was selected for multiple reasons (Figure 1). Site selection considerations were typical for selecting a rotary screw trapping site and included fine scale physical characteristics (e.g., access for installation, operation, and removal, water velocities, river depth and width, anchoring locations), broad scale site location

implications (e.g., sites location in the basin and proximity to other trap locations), and landowner permission for access. Site selection was finalized after considering multiple options in the Upper Chehalis River. The location has optimal physical conditions (e.g., flow, depth) within the basin and access was granted for trapping operations.




Figure 1. Upper Chehalis main stem rotary screw trap (46.634747, -123.167972). Anadromous streams represent stream habitat within the predicted coho salmon range of occurrence (299.8 km) using a 0.50 probability decision threshold upstream of the upper Chehalis River rotary screw trap. Non-anadromous streams represent stream habitat outside the predicted coho salmon range of occurrence (919.4 km) upstream of the trap location (Walther 2021).

### Trap Operation

A 1.5 m (5-foot) diameter rotary screw trap (RST) was operated near rkm 151.7 of the Chehalis River. The screw trap used internal flights rotating by water pressure to capture downstream migrants and funnel them into a holding area (livebox) at the back of the trap where fish were held until sampling. In 2022, the trap was scheduled to operate continuously from March 10 through July 11, although unscheduled trap outages did occur due to high flows. Three trapping positions were identified in 2022 to maximize trap efficiency (Figure 2).

Instantaneous water temperature and trap status information (e.g., fishing or not fishing, cone revolutions per minute) were collected at each fish sampling event (“trap check”). Water temperatures in fish holding containers were monitored throughout sampling events. Stream temperature was also monitored with a temperature data logger (HOBO 64K Pendant) deployed adjacent to the trap and cabled to the bank that collected temperature at 30-minute intervals. Data loggers were calibrated according to Winkowski et al. (2018). Stream flow was monitored by the USGS discharge gage near Doty, Washington located in the main stem Chehalis River 13.41 km upstream of the trap (USGS 12020000).



Figure 2. Upper Chehalis River trap site and fishing positions.

## Fish Collection

Fish sampling commenced daily in the morning and was adjusted to earlier times as stream temperatures increased to >18°C throughout the season. Crews monitored river flows and weather several times daily and modified operations in response to environmental conditions, such as earlier or multiple checks to minimize temperature impacts on fish health. Fish were removed from the live box, transferred to 5-gallon buckets, and moved to small dish tubs for sampling. Fish were anaesthetized with tricaine methanesulfonate (MS-222) prior to enumeration and biological sampling. An anaesthetizing solution was created by diluting 10 – 25 ml of a MS-222 solution (5g of MS-222 dissolved in 500 ml of water in a 500 ml container) into 2 – 3 L of water. This solution was replaced as necessary. Samplers continually evaluated fish response to the solution and targeted the lowest dosages needed to complete biological sampling.

During sampling, all fish were identified to species and enumerated. Chinook, coho, and steelhead were further categorized by life stage and age class. Marks associated with trap efficiency trials (see *Trap Efficiency Trials* section) and hatchery origin (clipped adipose fin) were examined on all Chinook, coho, and steelhead. We expected to capture hatchery origin coho and steelhead, released upstream of the trap site at WDFW acclimation ponds located on Eight Creek, a tributary to Elk Creek. Fork length (FL) and scales were collected from a subsample of wild (adipose fin intact) coho and steelhead (Table 2). Only fork lengths were collected from Chinook (no scales).

Table 2. Sample rates for biological data collection from wild juvenile salmonids.

Sample Type	Species	Fry	Parr	Transitional/Smolt
Fork Length	Chinook	1 <sup>st</sup> 10 daily	1 <sup>st</sup> 10 daily	1 <sup>st</sup> 10 daily
	Coho	1 <sup>st</sup> 10 daily	1 <sup>st</sup> 10 daily	All efficiency marked individuals (100 daily)
	Steelhead	1 <sup>st</sup> 10 daily <sup>a</sup>	1 <sup>st</sup> 10 daily	All Efficiency marked individuals (100 daily)
Scales	Chinook <sup>b</sup>	---	---	---
	Coho	---	---	1 <sup>st</sup> 5 daily
	Steelhead	---	---	1 <sup>st</sup> 5 daily

<sup>a</sup>Trout fry included both steelhead/rainbow trout and cutthroat.

<sup>b</sup>No scale samples were collected from Chinook.

Life stage categories followed WDFW protocols developed for the Lower Columbia ESU monitoring program (see Appendix A for life stage decision tree). The five life stage categories include fry, parr, transitional, smolt, and adult. Fry and adults were assigned based on length criteria (fry ≤45 mm FL and adults >300 mm FL [cutthroat], 301 – 499 mm FL [rainbow], or ≥500 mm FL [steelhead]). Parr, transitional, and smolt life stages were assigned based on phenotypic traits. Parr had distinct parr marks or showed no signs of smoltification; transitionals showed initial signs of smoltification (i.e., silvery appearance and faded parr marks); and smolts showed advanced signs of smoltification (i.e., faded parr marks, deciduous scales, silvery appearance, black banding along the trailing edge of the caudal fin, and translucent pectoral and pelvic fins).

Age class represented the number of rearing years in freshwater as measured from scale samples. Over the 35 years of trapping at the main stem Chehalis site, beginning in 1986, yearling Chinook salmon have rarely been observed. Furthermore, the vast majority of juvenile Chinook identified

in the field are assigned to the subyearling age class based on fork length. While extremely rare, individuals >150 mm are encountered that are outside of the fork length range of subyearling outmigrants and get categorized as yearlings in the field. These individuals are often opportunistically sampled for scales to verify age (Table 3). For these reasons we assume Chinook in the Upper Chehalis follow a similar life history as the mainstem Chehalis River. For coho salmon, all fry and parr were classified as subyearlings and all smolts and transitionals were classified as yearlings (Table 4). For steelhead, the field-assigned ‘yearlings’ could be any of 1-, 2-, or 3-year-old individuals that could not be distinguished by length in the field (Table 5). Therefore, the age composition of steelhead was further described using scale data.

Table 3. Date and length criteria used for field calls of juvenile Chinook salmon.

Life Stage	Age Class	Date Range	Length Range (mm FL)
Fry	---	Start – end	≤45
Parr, Transitional, Smolt	Subyearling	Start – end	46 - 150
Transitional, Smolt	Yearling (+)	Start – end	>150

Table 4. Date and length criteria used for field calls of juvenile coho salmon.

Life Stage	Age Class	Date Range	Length Range (mm FL)
Fry	---	Start – end	≤45
Parr	Subyearling	Start – end	>45
Transitional, Smolt	Yearling	Start – end	>45

Table 5. Date and length criteria used for field calls of juvenile steelhead trout.

Life Stage	Age Class	Date Range	Length Range (mm FL)
Fry	---	Start – end	≤45
Parr	NA	Start – end	>45
Transitional, Smolt	Yearling (+)	Start – end	>45
Adult (Resident RBT)	NA	Start – end	300 - 499
Adult (STLH kelt)	NA	Start – end	>500

### *Trap Efficiency Trials*

A single trap, mark-recapture study design stratified by week was used to estimate juvenile salmon and steelhead abundance (Volkhardt et al. 2007). The mark-recapture design consisted of counting maiden caught fish (maiden captures) in the trap and marking a known number of the captured fish for release at an upstream location (marks). Marked fish that were recaptured in the trap after release (recaptures) were enumerated to calculate trap efficiency. Maiden captures, marks, and recaptures were stratified by week to account for heterogeneity in trap efficiency throughout the season. Weekly estimate periods began on Monday and ended on Sunday.

Trap efficiency trials were conducted using selected species, origin, and life stages to estimate outmigrant abundance (Table 6). Species included in the trap efficiency trials were Chinook, coho, and steelhead. All trap efficiency trials were conducted with wild (adipose fin intact) fish. For

Chinook, trap efficiency trials were conducted with transitional and smolt life stages because these were the life stages for which we intended to generate an abundance estimate. Efficiency trials were not conducted on Chinook fry outmigrants as the trap was not operated for the full duration of the early timed outmigration; therefore, no estimate was generated for the Chinook fry life stage. For coho and steelhead, trap efficiency trials were conducted with transitional and smolt life stages. Fry and parr life stages were not included in the trap efficiency trials for coho and steelhead because we assumed that these life stages were not actively outmigrating. Fish in good physical condition were selected for efficiency trials whereas fish in poor physical condition were enumerated and released downstream. The goal was to mark a maximum of 100 fish per species per day and 700 per species per week for efficiency trials. However, this number varied based on fish capture rates throughout the season.

Table 6. Abundance estimate groups defined by species, origin, life stage, and age class. Life stages included in the estimates were transitional (T), and smolt (S). Age classes included in the estimates were subyearling (SY) and yearling (Y). FL = Fork length.

Abundance Group	Origin	Life Stage	Age Class	Note
Chinook	Wild	T, S	SY	FL $\geq$ 45 mm
Coho	Wild	T, S	Y, SY	
Steelhead	Wild	T, S	Y	

Marked fish were released 1.43 kilometers upstream of the trap location at the intersection of River Rd and State Route 6 (Table 7).

Mark types and rotation schedules allowed the data to be stratified by week for the purpose of analysis. This was irrelevant for coho and steelhead, however, because they were marked using individual PIT tags. The different mark types for each species are listed below (Table 7). Releases generally occurred within 1-3 hours of the start of a trap check.

Table 7. Trap efficiency marks and release locations for each abundance estimate group. Efficiency marks are visible implant elastomer tag (VIE) and passive integrated transponder tag (PIT).

Abundance Group	Trap Efficiency Marks			Release location	
	Mark Types	Rotation Schedule	Mark Rotation	Description	Distance upstream of trap (rkm)
Chinook	VIE (n=5)	Weekly	1 week	Intersection	1.43
Coho	PIT	Individual	Individual	Intersection	1.43
Steelhead	PIT	Individual	Individual	Intersection	1.43

### *Assumption Testing*

The six basic assumptions needed to be met for unbiased estimates in mark-recapture studies include: 1) the population is closed, 2) marks are not lost, 3) marking does not affect behavior, 4) initial capture probabilities are homogenous, 5) the second sample is a random representative sample (i.e., marked and unmarked fish are completely mixed), and 6) mark status is reported correctly (Volkhardt et al. 2007). Throughout the season multiple trials were conducted to reduce



the probability of any assumption violations. These included mark/tag retention trials to ensure marks/tags were not lost, mark/tag detection trials to ensure that mark/tags were not missed and that they were reported correctly, and mark-related mortality trials to ensure marking/tagging did not affect behavior or survival.

### *Analysis*

The Bayesian Time-Stratified Population Analysis System (BTSPAS, Bonner and Schwarz 2014) was used to estimate abundance of Chinook, coho, and steelhead (Table 1). This method uses Bayesian P-splines and hierarchical modeling of trap efficiencies to determine abundance with known precision through time, which allows for estimation during missed trapping days and for time strata with minimal efficiency data (Bonner and Schwarz 2011). Data for the analysis were stratified by week and included the total catch of unmarked fish (i.e., maiden captures), marks released, marks recaptured, and proportion of time sampled. The proportion of time sampled each week was included to adjust for missed catch during trap outages.

No trapping occurred from April 3 to April 6, 2022, and April 18 to April 19, 2022, due to high river flows. However, for the missed trapping period, the BTSPAS model produced estimates with known precision using the entire season's dataset by fitting a spline through those dates.

Prior to analysis, marks were removed during periods when the trap did not continuously fish for 48 hours after release because those marks were not available for recapture. For coho and steelhead, two periods were added prior to the trapping season. For Chinook, four periods were added post trapping season. For all species the first and last periods were set to 0 to allow the model to estimate the beginning and tail of each run. For all species estimates, a BTSPAS diagonal model was used. In each case, the model arguments were as follows: number of chains = 4, iterations = 100,000, burn-in = 50,000, sims = 25,000 and a thin rate of 2. Model convergence was assessed by visually inspecting the trace plots and using the potential scale reduction statistic, or Rhat. The Rhat statistic measures the ratio of the average variance draws within each chain to the variance of the pooled draws across chains; if all chains are at equilibrium, these will be the same and Rhat will be 1. If the chains have not converged to a common distribution, the Rhat statistic will be >1. Models were considered to have converged if MCMC chains were fully mixed based on visual inspection, and Rhat was less than 1.002 for all parameters (Gelman et al. 2004). The BTSPAS analysis was executed in R v.3.4.1 (R Core Team, 2017) using R version 2021.1.1 (R Core Team 2021) and the BTSPAS package (Bonner and Schwarz 2014).

## Results

### *Summary of Fish Species Encountered*

A diverse assemblage of fish species was encountered throughout the 2022 trapping season. Native fish included juvenile Chinook and coho salmon, steelhead, rainbow, and cutthroat trout, redbreasted shiner (*Richardsonius balteatus*), smallmouth bass (*Micripterus dolomieu*), dace species (*Rhinichthys spp.*), speckled dace (*R. osculus*), longnose dace (*R. cataractae*), sucker species (*Catostomus spp.*), northern pikeminnow (*Ptychocheilus oregonensis*), Pacific lamprey (*Entosphenus tridentatus*), brook lamprey (*Lampetra planeri*), and sculpin (Cottidae). Non-native fish included rock bass (*Ambloplites rupestris*).

### *Trap operation*

The trap was operated from March 10, 2022, to July 11, 2022. There were eleven occurrences of trap outages (Appendix B). For all eleven events, the outage time was known exactly because the trap stopped fishing either when staff lifted the cone during periods of high flows, debris and maintenance, or there was a trap alarm. All events except for two were under 4 hours in duration. One event began on April 3, 2022, and lasted for three days and the other event began on April 18, 2022, and lasted for one day. Both events were due to high river flows.

### *Assumption Testing Trials*

In 2022, results from the mark retention trials indicated that mark/tag retention was high based on trials that lasted 24 hours. Estimated mark retention was 100% (visible implant elastomer = VIE, 37 tagged) for Chinook and 100% (passive integrated transponder = PIT tag, 31 tagged) for coho and 100% (PIT tag, 6 tagged) for steelhead. A double tag/mark experiment with steelhead and coho also indicated that mark retention was high at 97% (33 total recaps, 32 with PIT tag and scar present at time of recapture) for steelhead and 97.3% (185 total recaps, 180 with PIT tag and scar present at time of recapture) for coho. For all trials, mark/tag related mortality was low. Estimated survival was 100% (VIE, 37 out of 37 tagged) for Chinook and 100% for coho (PIT tag, 31 out of 31 tagged) and 100% for steelhead (PIT tag, 6 out of 6 tagged) over the 24-hour holding period. Differences in initial capture probabilities due to body size were also tested using a Kolmogorov–Smirnov test, which found that the fork length of maiden captures versus recaptures did not differ significantly for Chinook during period 16 ( $D = 0.18$ ,  $p \geq 0.27$ ). PIT tagging coho and steelhead allowed for logistic regression analysis of probability of recapture by fork length. The relationship for coho between probability of recapture and fork length was significant ( $p = 0.02$ ) and the relationship for steelhead was also significant ( $p = 0.05$ ), indicating that tagged coho and steelhead may be more susceptible to recapture than non-tagged fish.

### *Chinook*

The Chinook outmigrant estimate was derived for the ‘subyearling’ life history and included transitionals and smolts. Chinook outmigrants were observed in low numbers the first week of trapping (March 11<sup>th</sup>, trapping period 1), peaked around the fourth week of June, and declined to low numbers again by the last week of trapping (July 5<sup>th</sup>, trapping period 18, Figure 3, Appendix C).

Generally, all Chinook outmigrants were assumed to be Age-0. Chinook ranged from 45 to 117 mm. Fork length of Chinook increased steadily throughout the season with an average of 50.2 mm ( $\pm 4.5$  mm SD) and 101.8 mm ( $\pm 8.1$  mm SD) in the first and last sampled week of trapping, respectively (Figure 4).

A total of 2,117 Chinook subyearling outmigrants were captured of these, 2,004 were marked, and 219 were recaptured (Appendix C; Periods 1 – 18). Modeled weekly trap efficiencies ranged from 10.3 to 11.7%. Trap efficiency and maiden catches may be affected by river flows. In 2022, average daily river flows increased to >100 cubic feet per second (cfs) on March 14, 15 and April 4, 18,19, May 6, 7, 16, 29 and June 5, and 10 (Figure 5). Abundance of wild Chinook subyearling outmigrants was estimated to be 19,870 (95% confidence intervals, CI = 17,383–22,951) with a coefficient of variation (CV) of 7.2%.

In 2021, the total number of adult spring Chinook that spawned in the Chehalis River above our trap site was estimated to be 313 (all NOR) and adult fall Chinook was estimated to be 533 (all NOR), producing an overall smolt-per-adult estimate of 23.2 for the 2021 brood year of naturally spawning Chinook (C. Holt, WDFW, personal communication). Estimating subyearling Chinook productivity through time is a goal of this study going forward.




Figure 3. Number of outmigrants (top panel) and trap efficiency (bottom panel) by week for wild Chinook subyearlings produced above the Upper Chehalis River smolt trap in 2022. The total estimate is 19,870 with a coefficient of variation (CV) of 7.2%. Error bars and shading around point estimates represent 95% confidence intervals.




Figure 4. Box plots of fork lengths of wild Chinook subyearling outmigrants (transitionals, smolts) by week at the Upper Chehalis River screw trap, 2022. Boxes represent the median, first and third quartiles, whiskers represent the interquartile ranges, and dots represent outliers.




Figure 5. Chinook wild transitional and smolt trap efficiency (top), maiden catch (bottom) and flow in cubic feet per second (cfs top & bottom), with trap outages (red) by date at the Upper Chehalis smolt trap in 2022.

### Coho

The coho outmigrant estimate included yearlings in transitional and smolt life stages. Approximately 79% of the outmigrants observed at the trap were categorized as the ‘smolt’ phenotype whereas 21% were categorized as ‘transitional’. Coho outmigrants were observed in low numbers the first week of trapping (March 11<sup>th</sup>, trapping period 1), peaked in late May, and were last observed the week of July 5<sup>th</sup> (trapping period 18, Figure 6, Appendix D).

Scale age data indicated three age classes of the coho juvenile outmigration. A total of 356 scale samples were collected and 95% were successfully aged. Age-1 coho were the dominant age class

(96.1%) and Age-0 and Age-2 were much less prevalent (0.9%) and (3%) (Figure 7, Table 8). Fork lengths averaged 108.3 mm ( $\pm$  4 mm) subyearlings, 118 mm ( $\pm$  10.5 mm) Age-1, and 123.6 mm ( $\pm$  12.4 mm) Age-2.

In 2022, a total of 2,213 coho outmigrants were captured, 1,988 coho were marked, and 183 were recaptured (Appendix D). Modeled weekly trap efficiencies ranged from 8.6% to 9.4%. Trap efficiency and maiden catches may both be affected by river flows. In 2022, average daily river flows increased to >100 cubic feet per second (cfs) on March 14, 15 and April 4, 18, 19, May 6, 7, 16, 29, and June 5 and 10 (Figure 8). The abundance of 2022 wild coho outmigrants was estimated to be 24,434 (95% CI = 21,175–28,403) with a CV of 7.6%.

In 2020, the total number of adult coho spawners in the Chehalis River upstream of the trap site was estimated to be 1,340 (all NOR), producing a smolt-per-spawner estimate of 18.2 for the 2020 brood year of naturally spawning coho (C. Holt, WDFW, personal communication). Estimating coho productivity through time is a goal of this study going forward.




Figure 6. Number of outmigrants (top panel) and trap efficiency (bottom panel) by week for wild coho smolts and transitionals produced above the Upper Chehalis River smolt trap in 2022. The total estimate is 24,434 with a coefficient of variation (CV) of 7.6%. Error bars and shading around point estimates represent 95% confidence intervals.






Figure 7. Plot of length and age by date for wild coho outmigrants (transitionals, smolts) at the Upper Chehalis River screw trap, 2022.




Figure 8. Coho wild transitional and smolt trap raw efficiency (top), maiden catch (bottom) and flow in cubic feet per second (cfs, top & bottom) with trap outages (red) by date at the Upper Chehalis smolt trap in 2022.

Table 8. Freshwater ages of wild coho outmigrants (transitionals, smolts) at the Upper Chehalis River screw trap, 2022. Data are scale ages of sampled juveniles by week.

<b>Period</b>	<b>Start Date</b>	<b>End Date</b>	<b>No. Scales</b>	<b>Age-0</b>	<b>Age-1</b>	<b>Age-2</b>	<b>Not Determined</b>
1	3/07	3/13	8		7		1
2	3/14	3/20	10		10		
3	3/21	3/27	9		9		
4	3/28	4/03	20		20		
5	4/04	4/10	17		16	1	
6	4/11	4/17	21		18		3
7	4/18	4/24	28		26		2
8	4/25	5/01	35		31	1	3
9	5/02	5/08	35		26	4	5
10	5/09	5/15	35		34		1
11	5/16	5/22	35		34		1
12	5/23	5/29	35		34		1
13	5/30	6/05	35		32	2	1
14	6/06	6/12	24		24		
15	6/13	6/19	5	1	2	2	
16	6/20	6/26	1		1		
17	6/27	7/03	2	1	1		
18	7/04	7/10	1	1			

### *Steelhead*

The steelhead outmigrant estimate included both transitional and smolt life stages. Of these life stages, approximately 63% of outmigrants observed were classified as the smolt phenotype compared to 37% transitional. Steelhead outmigrant numbers were low during the first week of trapping March 11 (trapping period 1), peaked mid-May, and were last observed the week of June 28 (trapping period 17) (Figure 9, Appendix E).

Scale age data indicated that the sampled steelhead were one, two, and three years of age (Figure 10, Table 9). Fork length averaged 154 mm ( $\pm$  14.1 mm) for one-year olds, 160 mm ( $\pm$  14.1 mm) for two-year olds, and 169 mm ( $\pm$  14.9 mm) for three-year olds. Age composition of successfully aged steelhead was 34.9% Age-1, 60.1% Age-2, and 5% Age-3.

A total of 575 steelhead outmigrants were captured throughout the season (Appendix E). A total of 523 steelhead were marked and 33 were recaptured. Modeled weekly trap efficiencies ranged from 2.6% to 7.9%. Trap efficiency and maiden catches may both be affected by river flows. In 2022, average daily river flows increased to >100 cubic feet per second (CFS) on March 14, 15 and April 4, 18,19, May 6, 7, 16, 29, and June 5 and 10 (Figure 11). Abundance of wild steelhead outmigrants was estimated to be 10,204 (95% CI = 7,261–15,252) with a CV of 19.6%.

Adult steelhead spawners contributing to the 2022 smolt outmigration came from the 2018 through 2020 brood years. Spawners were estimated to be 963, 1,078, and 917 for these years, respectively

(C. Holt, WDFW, personal communication). More monitoring is required to estimate steelhead productivity above the trap by brood year, but that is a project goal.




Figure 9. Number of outmigrants (top panel) and trap efficiency (bottom panel) by week for wild steelhead smolts and transitionals produced above the Upper Chehalis River smolt trap in 2022. The total estimate is 10,204 with a CV of 19.6%. Error bars and shading around point estimates represent 95% confidence intervals.




Figure 10. Plot of length and age data by date for wild steelhead outmigrants (transitionals, smolts) at the Upper Chehalis River screw trap, 2022.




Figure 11. Steelhead wild transitional and smolt trap efficiency (top), maiden catch (bottom) and flow in cubic feet per second (cfs, top & bottom) with trap outages (red) by date at Upper Chehalis smolt trap in 2022.

Table 9. Freshwater ages of wild steelhead outmigrants (transitionals, smolts) at the Upper Chehalis River screw trap, 2022. Data are scale ages of sampled juveniles by week.

<b>Period</b>	<b>Start Date</b>	<b>End Date</b>	<b>No. Scales</b>	<b>Age-1</b>	<b>Age-2</b>	<b>Age-3</b>	<b>Not Determined</b>
1	3/07	3/13	7	1	3		3
2	3/14	3/20	16	1	9	3	3
3	3/21	3/27	27	6	14	1	6
4	3/28	4/03	35	6	19	4	6
5	4/04	4/10	12		7	2	3
6	4/11	4/17	10	1	8		1
7	4/18	4/24	22	3	6		13
8	4/25	5/01	34	5	27	1	1
9	5/02	5/08	34	5	23	2	4
10	5/09	5/15	32	12	17		3
11	5/16	5/22	33	17	10		6
12	5/23	5/29	32	17	12		3
13	5/30	6/05	15	10	1		4
14	6/06	6/12	6	5	1		
15	6/13	6/19	1	1			
16	6/20	6/26	0				
17	6/27	7/03	1	1			

## Discussion

### *Basin-wide Context*

This report presents results from the 2022 salmon and steelhead smolt outmigration for the Upper Chehalis River and includes the second year of smolt monitoring estimates produced for this portion of the basin. The abundance estimates provided in this report represent juvenile salmonids that completed their freshwater rearing in habitats upstream of the trap location, specifically production from upstream of rkm 151.7. The smolt trap in the Upper Chehalis is one of four juvenile monitoring programs in the Chehalis Basin. By operating multiple smolt traps in the basin, smolt abundance estimates can be partitioned to specific locations, providing finer scale resolution of freshwater production. Habitat upstream of the trap is characterized predominately by timber land, but also includes agricultural land. This area has several tributaries including Elk Creek, Dunn Creek and Crim Creek. In addition to freshwater production from these tributaries, some juveniles emerge from the gravel upstream of the trap location and redistribute to areas downstream during their freshwater rearing period. These fish were not included in the estimates, especially coho salmon which are known to redistribute in a downstream direction during the fall months in search of suitable overwintering habitat (Winkowski et al. 2018).

Due to a proposed Flood Retention Expandable (FRE) facility at rkm 174.1, there has been increased interest in salmon and steelhead abundance and production in the Upper Chehalis. A study completed in 2019 reported that spring Chinook, fall Chinook, coho, and steelhead spawners above the FRE facility represented 1.24%, 3.37%, 2.72% and 15.43% of the total spawners in the Chehalis River Basin, respectively (Ronne et al. 2020). The upper basin is particularly important

for steelhead. In fact, steelhead spawners upstream of the smolt trap contributed anywhere from 12 to 23% of the total basin steelhead abundance from 2013-2019, despite having only 4% of the basin's suitable steelhead spawning habitat (Ronne et al. 2020). This work estimating juvenile steelhead production from the upper basin will be an important element for determining the impacts of the proposed FRE on steelhead production in the Chehalis Basin and provide valuable information on stock status for conservation and management purposes. For example, in 2022, the Secretary of Commerce received a petition to list Olympic Peninsula steelhead Distinct Population Segment (DPS) under the Endangered Species Act (ESA). Steelhead in the Chehalis are part of the Southwest Washington DPS that is not listed, but should that ever change, juvenile production estimates will be important for examining productivity trends.

The estimate of Chinook subyearling outmigrants represents a portion of the total freshwater production of Chinook upstream of the trap location in 2022 since it does not include the earlier timed fry migrants. The subyearling estimate of 19,870 Chinook is the second reportable estimate produced for this portion of the Chehalis basin. Surprisingly, the Chinook production estimate from the Upper Chehalis trap decreased by 83.3% between 2021 (118,834) and 2022 (19,870). Adult spawner abundance of Chinook in the Chehalis River Basin was 42.7% higher in 2020 (2,828 spring Chinook and 15,924 fall Chinook) than 2021 (2,575 spring Chinook and 8,176 fall Chinook), but probably not enough to account for the 83.3% decrease in juvenile production from 2021 to 2022. Therefore, the lower juvenile production in 2022 was probably related to variations in stream conditions between 2021 and 2022. Late fall 2021 and early winter 2022 were characterized by several high flow events above action stage that likely disrupted Chinook redds and affected emerging fry. Emergent fry may have been washed downstream during the high-water events, leaving fewer juveniles to rear in freshwater to the parr/smolt stage. Restoration actions designed to reduce impacts of high flow events (e.g., flooding, scouring) will likely have a positive impact on the survival of juvenile Chinook salmon.

The 2022 Chinook estimate represented 8.0% of the total subyearling production (247,707) above the Lower Chehalis mainstem trap located downstream at rkm 84 (D. Olson, WDFW, personal communication). Generating a 'subyearling' estimate is relevant to habitat restoration planning because the 'subyearling' component of the outmigration represents the numbers of juveniles that are supported by freshwater habitats upstream of the trap site and previous work demonstrated that >95% of adult Chinook returning to the upper Chehalis had a subyearling life history (Campbell et al. 2017). Fry migrants do not spend much time rearing in freshwater habitats but rather make extensive use of downstream, estuary, and nearshore growing environments prior to entering the ocean (Sandell et al. 2014, Beamer et al. 2005). Other studies in western Washington have observed that, within a watershed, the numbers of subyearling Chinook outmigrants are relatively consistent from year to year and concluded that abundance of this life history reflects a freshwater rearing capacity (Anderson and Topping 2018, Zimmerman et al. 2015). Additional Chinook production beyond this capacity appears to migrate downstream as 'fry' in a density-dependent manner (Greene et al. 2005). Extending this density-dependent migration hypothesis to the Chehalis River will require additional years of juvenile monitoring coupled with adult Chinook spawner data above the trap location.

Estimates of annual freshwater production of wild coho smolts in the Chehalis River Basin averaged 2.1 million (0.5 to 3.7 million) since WDFW began monitoring smolt production in the 1980s (Litz 2023). The proportion of coho habitat upstream of our trapping location represents



approximately 6.6% of the rearing habitat relative to the entire basin (Walther 2021). The proportion of juvenile coho salmon from upstream of the trapping location relative to basin-wide production was estimated to be quite low at 0.8% in 2022. Based on this information, it may be that a relatively small proportion of all wild coho in the Chehalis River watershed complete their freshwater rearing in the upper Chehalis, Elk Creek, and other small tributaries upstream of the trap site. Conversely, a larger proportion of wild coho appear to complete their freshwater rearing in the main stem and tributaries downstream of the trap location, which make up approximately 93.4% of coho salmon habitat in the Basin (Walther 2021). Spawning and rearing areas downstream of the trap location include off-channel sloughs and ponds along the main stem river, major tributaries such as the Black, Satsop, Wishkah, and Hoquaim rivers, and smaller tributaries including Porter and Cloquallum Creek.

Our estimate of juvenile coho production in the Upper Chehalis River basin in 2022 was 24,434. This number represents 15.0% of the estimate for coho production above our Lower Chehalis mainstem trap site in 2022. Generating an unbiased and precise estimate for coho at this new location has mainly been possible due to learning how the equipment operates best at this location and adapting field protocols. If rearing habitat is a limiting factor for coho in the Chehalis Basin, as suggested in other streams in western Washington (Reeves et al. 1989), then restoration activities targeting rearing habitat should increase the productivity of coho in the Chehalis Basin, consistent with the goals of the Aquatic Species Restoration Plan (ASRPSC 2019).

This report provides the second reportable estimate of wild steelhead smolt production from the Upper Chehalis River basin upstream of river kilometer 151.7. This estimate represents 28.2% of the estimate for steelhead production above the Lower Chehalis mainstem trap site in 2022. The estimate of 10,204 steelhead outmigrants from the roughly 299.8 anadromous rkm upstream of the trap (Walther 2021) corresponds to 34.0 wild steelhead smolts  $\text{km}^{-1}$ . This smolt density is low compared to other western Washington watersheds where steelhead smolt estimates are available, such as the Coweeman River (average 243 smolts  $\text{km}^{-1}$ ) or the Wind River (average 240 smolts  $\text{km}^{-1}$ ) (T. Buehrens WDFW, personal communication). The reasons for these differences are not yet apparent and may reflect the difference between available versus suitable rearing habitat upstream of the Upper Chehalis River trap. Of note, some studies (Ashcraft et al. 2017, Ronne et al. 2018) identified the Upper Chehalis sub-basin as a particularly productive steelhead spawning area. Over five years of monitoring, surveyors estimated 600-1,000 redds (or 900-1,800 steelhead spawners) in this area of the basin. The Upper Chehalis sub-basin has the high gradient, coarse substrate habitat typically associated with rearing of juvenile steelhead. Another possible explanation is that steelhead parr have the option of rearing downstream of the trap. However, rearing areas downstream of the trap are generally low gradient main stem reaches, off-channel sloughs, and ponds along the main stem river. These habitat types are not considered high quality juvenile steelhead rearing habitat (Burnett et al. 2007). Interestingly, during the first year of trapping, scale results indicated that most juvenile steelhead were Age-1 (71.4% of outmigrants). In the second year of trapping, that proportion was reduced by half to 34.9%, with most steelhead outmigrating at Age-2 (60.1%), consistent with observations throughout western Washington. The majority of smolt and transitional steelhead captured in 2022 were PIT tagged prior to release. Therefore, recoveries of PIT tagged adults in future years could provide insight into marine survival or repeat spawning events. Ideally, installation of a PIT array below the trap could be used to track tagged juvenile and adult returns to further research survival and life history diversity of natural origin steelhead.

### *Next Steps*

The Upper Chehalis River estimates presented here provide critical information for salmon and steelhead smolt production in the basin but trapping in this location presents many challenges. Temperature concerns are prevalent later in the season at this location in the Chehalis River. For example, mean river temperatures from April to July 2022 increased from at 12.5°C to 22.1°C during the period of Chinook outmigration (Figure 12, Table 10). When river temperature exceeds 18°C, sampling protocols were minimized to reduce fish stress. Another challenge faced during the 2022 trapping season was the difference in river flows throughout the trapping season, with decreases of two orders in magnitude between early April and early July. In 2022, the most effective fishing locations for the trap were established, depending on flow. Additionally, several action stage flow events occurred in late 2021 and early 2022, and it is likely that these extreme flow conditions of the river impacted rearing Chinook, whose abundance declined by 83.3% between 2021 and 2022. It is anticipated that restoration treatments such as engineered log jams, that improve rearing habitat by increasing instream complexity and promoting side-channel and floodplain connectivity, would benefit rearing Chinook by providing refuge during high flow events.

Table 10. Mean monthly stream temperatures °C recorded at Upper Chehalis River smolt trap near river km 151.7, 2022.

Month	Mean (°C)
April	12.7
May	14.5
June	18.8
July	22.1




Figure 12. Chinook maiden catch and average daily stream temperature (°C) at the Upper Chehalis River smolt trap, 2022.

In summary, 2022 represents the second year for which wild juvenile Chinook, coho, and steelhead production has been described from this location of the Upper Chehalis River. For all three species, unbiased and precise estimates of smolt abundance were generated and life history characteristics described, including size, age, and timing of outmigration. Life history strategies reflect how the existing habitat contributes to freshwater production of salmon and steelhead. Continued monitoring will provide even greater understanding of how variability in habitat and environmental conditions affects freshwater production over time.

## References

- Anderson, J.H., and P.C. Topping. 2018. Juvenile life history diversity and freshwater productivity of Chinook Salmon in the Green River, Washington. *North America Journal of Fisheries Management* 38:180-193.
- Ashcraft, S., C. Holt, M. Scharpf, M. Zimmerman, and N. VanBuskirk. 2017. Spawner Abundance and Distribution of Salmon and Steelhead in the Upper Chehalis River, 2013-2017, FPT 17-12. Washington Department of Fish and Wildlife, Olympia, Washington.
- ASRPSC (Aquatic Species Restoration Plan Steering Committee). 2019. Chehalis Basin Strategy: Aquatic Species Restoration Plan – Phase I. Office of the Chehalis Basin. Publication 19-06-009. November 2019.
- Beamer, E.M., A. McBride, C.M. Greene, R. Henderson, G.M. Hood, K. Wolf, K. Larsen, C. Rice, and K.L. Fresh. 2005. Skagit River Chinook Recovery Plan. Appendix D. Delta and nearshore restoration for the recovery of wild Skagit River Chinook salmon: Linking estuary restoration to wild Chinook salmon populations, <http://www.skagitcoop.org/index.php/documents/>.
- Burnett, K.M., G.H. Reeves, D.J. Miller, S. Clarke, K. Vance-Borland, and K. Christiansen. 2007. Distribution of salmon-habitat potential relative to landscape characteristics and implications for conservation. *Ecological Applications* 17(1):66-80.
- Bonner, S.J., and C.J. Schwarz. 2011. Smoothing population size estimates for time-stratified mark-recapture experiments using Bayesian P-splines. *Biometrics* 67(4):1498-1507.
- Bonner, S.J., and C.J. Schwarz. 2014. BTSPAS: Bayesian Time-Stratified Population Analysis. R package version.
- Campbell, L.A., A.M. Claiborne, S. Ashcraft, M.S. Zimmerman, and C. Holt. 2017. Investigating Juvenile Life History and Maternal Run Timing of Chehalis River Spring and Fall Chinook Salmon Using Otolith Chemistry, FPT 17-15. Washington Department of Fish and Wildlife, Olympia, Washington.
- Gelman, A., J. Carlin, A. Stern, and D. B. Rubin. 2004. *Bayesian Data Analysis*, 2<sup>nd</sup> Edition. Chapman and Hall/CRC Press. Boca Raton, FL.
- Gilbertson, L., T. Jurasin, R. Coshow, and M. Miller. 2021. Run-Type Composition of Juvenile Chinook Salmon in the Upper Chehalis River Basin in 2020. Technical Report Series 2021-1. Quinault Indian Nation Department of Fisheries.
- Greene, C.M., D.W. Jensen, G.R. Pess, E.A. Steel, and E. Beamer. 2005. Effects of environmental conditions during stream, estuary, and ocean residency on Chinook salmon return rates in the Skagit River, Washington. *Transactions of the American Fisheries Society* 134:1562-1581.
- Groot, C., and L. Margolis, eds. 1991. *Pacific Salmon Life Histories*. UBC Press.
- Kiyohara, K., and M.S. Zimmerman. 2012. Evaluation of juvenile salmon production in 2011 from the Cedar River and Bear Creek, FPA 12-01. Washington Department of Fish and Wildlife, Olympia, Washington.
- Litz, M.N.C. 2023. 2023 wild coho forecasts for Puget Sound, Washington Coast, and Lower Columbia, FPA 23-01. Washington Department of Fish and Wildlife, Olympia, Washington.

M&AMT (Monitoring and Adaptive Management Team). 2021. Aquatic Species Restoration Plan Monitoring and Adaptive Management Plan. Office of the Chehalis Basin.

Olson, D., D. West., J. Winkowski, T. Seamons, and M. Litz. 2023. Chehalis River Smolt Production, 2021. FPA 23-06. Washington Department of Fish and Wildlife, Olympia, Washington.

R Core Team. 2021. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

Reeves, G.H., F.H. Everest, and T.E. Nickelson. 1989. Identification of physical habitats limiting the production of coho salmon in western Oregon and Washington. Gen. Tech. Rep. PNW-GTR-245. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 18 p

Ronne, L., N. Vanbuskirk, C. Holt, and M. Zimmerman. 2018. Spawner Abundance and Distribution of Salmon and Steelhead in the Upper Chehalis River, 2017-2018, FPT 18-09. Washington Department of Fish and Wildlife, Olympia, Washington.

Ronne L., N. VanBuskirk, and M. Litz. 2020. Spawner Abundance and Distribution of Salmon and Steelhead in the Upper Chehalis River, 2019 and Synthesis of 2013-2019, FPT 20-06 Washington Department of Fish and Wildlife, Olympia, Washington.

Sandell, T., J. Fletcher, A. McAninch, and M. Wait. 2014. Grays Harbor Juvenile Fish Use Assessment: 2013 Annual Report. Wild Fish Conservancy, prepared for the Chehalis Basin Habitat Work Group.

Volkhardt, G.C., S.L. Johnson, B.A. Miller, T.E. Nickelson, and D.E. Seiler. 2007. Rotary screw traps and inclined plane screen traps. Pages 235-266 *in* D. H. Johnson, B. M. Shrier, J. S. O'Neal, J. A. Knutzen, X. Augerot, T. A. O-Neil, and T. N. Pearsons, editors. Salmonid field protocols handbook: techniques for assessing status and trends in salmon and trout populations. American Fisheries Society, Bethesda, Maryland.

Walther, E.J. 2021. Salmonid Distribution Models to Support Restoration Planning Across the Fragmented Chehalis River basin, WA. MS Thesis, University of Alaska Fairbanks.


West, D., J. Winkowski, T. Seamons, and M. Litz. 2021. Chehalis River Smolt Production, 2020, FPA 21-06. Washington Department of Fish and Wildlife, Olympia, Washington.

Winkowski, J.J., E.J. Walther, and M.S. Zimmerman. 2018. Summer riverscape patterns of fish, habitat, and temperature across the Chehalis River basin. FPT 18-01. Washington Department of Fish and Wildlife. Olympia, Washington.

Zimmerman, M.S., C. Kinsel, E. Beamer, E.J. Connor, and D.E. Pflug. 2015. Abundance, survival, and life history strategies on juvenile migrant Chinook Salmon in the Skagit River, Washington. Transactions of the American Fisheries Society **144**:627-641.

## Appendices

Appendix A. Decision tree for assigning life stages of juvenile outmigrants developed by the Washington Department of Fish and Wildlife to ensure consistency in data collection protocols across juvenile trapping projects.



Updated 2.8.2016

Appendix B. Upper Chehalis River missed trapping periods, 2022.

<b>Last Time Observed Fishing</b>	<b>Time Stopped Fishing</b>	<b>Method to Determine Trap Not Fishing</b>	<b>Time Start Fishing again</b>	<b>Comments</b>
3/15/2022 0934	1 hr 41 min	Trap Pulled	3/15/2022 1115	Trap position moved
3/21/2022 0900	45 min	Trap Pulled	3/21/2022 0945	Clear log jam
3/22/2022 1230	40 min	Trap Pulled	3/22/2022 1310	Trap position moved
3/28/2022 1000	1 hr 30 min	Trap Pulled	3/28/2022 1130	Trap position moved
4/03/2022 1100	71 hr 50 min	Trap Pulled	4/06/2022 1050	Trap pulled for high flows
4/14/2022 1030	1 hr 10 min	Trap Pulled	4/14/2022 1140	Trap position moved
4/18/2022 1130	2 hr 50 min	Trap Pulled	4/18/2022 1420	Trap position moved
4/18/2022 1846	15 hr 54 min	Trap Pulled	4/19/2022 1040	Trap position moved, pulled for high flows
4/24/2022 1054	3 hr 1 min	Trap Pulled	4/24/2022 1355	Trap position moved
5/07/2022 1745	2 hr 30 min	Trap Stopper	5/07/2022 2015	Trap stopper fixed and debris cleared
5/31/2022 0910	1 hr 30 min	Trap Pulled	5/31/2022 1040	Trap repairs

Appendix C. Mark-recapture data for wild Chinook outmigrants (transitionals, smolts) organized by time period. Dataset includes total marks released (Total Mark), total marks recaptures (Total Recap), total maiden captures (Total Captures), and the proportion of time the trap fished during the time period (Prop Fished).

<b>Period</b>	<b>Start Date*</b>	<b>End Date*</b>	<b>Total Mark</b>	<b>Total Recap</b>	<b>Total Captures</b>	<b>Prop Fished</b>
1	3/08	3/14	0	0	0	1
2	3/15	3/21	0	0	0	1
3	3/22	3/28	0	0	0	1
4	3/29	4/04	4	0	13	1
5	4/05	4/11	10	0	14	1
6	4/12	4/18	11	0	13	1
7	4/19	4/25	18	0	27	1
8	4/26	5/02	15	0	10	1
9	5/03	5/09	10	2	14	1
10	5/10	5/16	8	1	14	1
11	5/17	5/23	58	3	78	1
12	5/24	5/30	150	13	144	1
13	5/31	6/06	221	26	276	1
14	6/07	6/13	363	38	348	1
15	6/14	6/20	177	32	199	1
16	6/21	6/27	479	59	570	1
17	6/28	7/04	354	31	273	1
18	7/05	7/11	122	14	124	1
19	7/12	7/18	1	NA*	NA*	0.001
20	7/19	7/25	1	NA*	NA*	0.001
21	7/26	8/01	1	NA*	0	0.001
22	8/02	8/08	1	NA*	0	0.001

\*Start and End Date reflect the dates of maiden captures to which the release and recapture data are applied for estimation. Release dates start and end one day before the recapture dates.

\*NA's indicate estimated missing data modeled in final analysis.



Appendix D. Mark-recapture data for wild coho outmigrants (transitionals, smolts) organized by time period. Data are the combined counts of subyearling and yearling coho. Dataset includes total marks released (Total Mark), total marks recaptures (Total Recap), total maiden captures (Total Captures), and the proportion of time the trap fished during the time period (Prop Fished).

<b>Period</b>	<b>Start Date*</b>	<b>End Date*</b>	<b>Total Mark</b>	<b>Total Recap</b>	<b>Total Captures</b>	<b>Prop Fished</b>
1	2/22	2/28	1	*NA	0	1
2	3/01	3/07	1	*NA	*NA	1
3	3/08	3/14	7	0	16	1
4	3/15	3/21	10	0	9	1
5	3/22	3/28	9	0	11	1
6	3/29	4/04	16	1	21	1
7	4/05	4/11	34	1	48	1
8	4/12	4/18	25	3	41	1
9	4/19	4/25	52	5	83	1
10	4/26	5/02	164	8	204	1
11	5/03	5/09	442	42	498	1
12	5/10	5/16	230	23	303	1
13	5/17	5/23	465	46	535	1
14	5/24	5/30	380	37	303	1
15	5/31	6/06	107	14	107	1
16	6/07	6/13	36	3	26	1
17	6/14	6/20	5	0	4	1
18	6/21	6/27	1	0	2	1
19	6/28	7/04	2	0	2	1
20	7/05	7/11	1	0	0	1

\*Start and End Date reflect the dates of maiden captures to which the release and recapture data are applied for estimation. Release dates start and end one day before the recapture dates.

\*NA's indicate estimated missing data modeled in final analysis.

Appendix E. Mark-recapture data for wild Steelhead outmigrants (transitionals, smolts) organized by time period. Dataset includes total marks released (Total Mark), total marks recaptures (Total Recap), total maiden captures (Total Captures), and the proportion of time the trap fished during the time period (Prop Fished).

<b>Period</b>	<b>Start Date*</b>	<b>End Date*</b>	<b>Total Mark</b>	<b>Total Recap</b>	<b>Total Captures</b>	<b>Prop Fished</b>
1	2/22	2/28	1	*NA	0	0.001
2	3/01	3/07	1	*NA	*NA	0.001
3	3/08	3/14	5	0	12	1
4	3/15	3/21	16	0	16	1
5	3/22	3/28	41	1	48	1
6	3/29	4/04	48	5	64	1
7	4/05	4/11	23	0	38	1
8	4/12	4/18	10	0	7	1
9	4/19	4/25	38	3	49	1
10	4/26	5/02	59	1	61	1
11	5/03	5/09	57	5	64	1
12	5/10	5/16	62	5	72	1
13	5/17	5/23	79	2	73	1
14	5/24	5/30	60	8	48	1
15	5/31	6/06	15	0	18	1
16	6/07	6/13	6	2	3	1
17	6/14	6/20	1	1	1	1
18	6/21	6/27	0	0	1	1
19	6/28	7/04	1	0	0	1
20	7/05	7/11	0	0	0	1

\*Start and End Date reflect the dates of maiden captures to which the release and recapture data are applied for estimation. Release dates start and end one day before the recapture dates.

\*NA's indicate estimated missing data modeled in final analysis.



Washington Department of  
**FISH & WILDLIFE**

This program receives Federal financial assistance from the U.S. Fish and Wildlife Service Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972. The U.S. Department of the Interior and its bureaus prohibit discrimination on the bases of race, color, national origin, age, disability and sex (in educational programs). If you believe that you have been discriminated against in any program, activity or facility, please contact the Civil Rights Coordinator at 833-885-1012, email: [CivilRightsTeam@dfw.wa.gov](mailto:CivilRightsTeam@dfw.wa.gov) or by mail at P.O. Box 43139, Olympia, WA 98504, or write to:

Department of the Interior  
Chief, Public Civil Rights Division  
1849 C Street NW  
Washington D.C. 20240