

2018

SCOTT FITKIN, District Wildlife Biologist  
JEFF HEINLEN, Assistant District Wildlife  
Biologist



Washington  
Department of  
**FISH and  
WILDLIFE**



# **DISTRICT 6 HUNTING PROSPECTS**

Okanogan County

## TABLE OF CONTENTS

FIRE AND ROAD CONDITIONS UPDATE.....	1
DISTRICT 6 GENERAL OVERVIEW .....	1
ELK .....	3
General Information, Management Goals, and Population Status .....	3
Which GMU Should Elk Hunters Hunt? .....	3
DEER.....	3
General Information, Management Goals, and Population Status .....	3
Which GMU Should Deer Hunters Hunt? .....	4
What to Expect During the 2018 Season .....	6
How to Find and Hunt Mule Deer .....	6
How to Find and Hunt White-tailed Deer .....	7
Deer Areas .....	8
BLACK BEAR.....	8
General Information, Management Goals, and Population Status .....	8
Which GMU Should bear Hunters Hunt? .....	8
What to Expect During the 2017 Season .....	9
COUGAR .....	10
General Information, Management Goals, and Population Status .....	10
Which GMU Should Cougar Hunters Hunt? .....	11
WATERFOWL.....	12
General Information.....	12
FOREST GROUSE .....	13
Species and General Habitat Characteristics.....	13
Harvest Trends and 2017 Prospects .....	13
PHEASANTS .....	14
Species and General Habitat Characteristics.....	14

QUAIL.....	15
Species and General Habitat Characteristics.....	15
TURKEYS.....	15
General Description .....	15
CHUKAR AND GRAY PARTRIDGE .....	16
General Description .....	16
DOVE .....	16
General Description .....	16

## FIRE AND ROAD CONDITIONS UPDATE

As always, check with the Okanogan-Wenatchee National Forest for current information on fire activity, access closures, and campfire restrictions. Significant and possibly complete campfire bans are likely, possibly well into the fall.

In general, U.S. Forest Service (USFS) roads are in somewhat better shape than they were at this time last year. The major washout in Black Canyon has been fully repaired, greatly improving access to the lower portion of GMU 242. A variety of minor secondary state and USFS road closures or washouts remain, particularly in the eastern portion of GMU 224 and the northern portion of GMU 239. In addition, USFS 5130 in the 8-mile drainage remains closed about a mile above the Copper Glance trailhead in the wake of last year's Diamond Creek Fire, and USFS Rd 5160-250 on the upper Chewuch River is closed just past the Andrews Creek Trailhead due to a spring washout. These two closures are currently precluding access to trailheads for the Pasayten Wilderness. Check with appropriate land management agencies for current conditions and updates.

On the plus side, vegetation is generally recovering nicely in recently burned areas and game animals have returned to these portions of the district. We experienced good seasonal growth again this year at all elevations.

For more information, see:

- [Okanogan National Forest, Methow Valley Ranger District](#)
- [DNR Regulated Fire Restrictions](#)
- [InciWeb Current Fire Status](#)
- [Okanogan County Emergency Management](#)

## DISTRICT 6 GENERAL OVERVIEW

District 6 is located along the Canadian border in north-central Washington and encompasses ten game management units: 203 (Pasayten), 204 (Okanogan East), 209 (Wannacut), 215 (Sinlahekin), 218 (Chewuch), 224 (Perrygin), 231 (Gardner), 233 (Pogue), 239 (Chiliwist), and 242 (Alta).


The western two-thirds of the district, stretching from the Okanogan River to the Pacific Crest, lies on the east slope of the Cascade Range and is dominated by mountainous terrain that gets more rugged as you move from east to west. Vegetation in this portion of the district ranges from desert/shrubsteppe at the lowest elevations to various types of conifer forests, culminating in alpine tundra on the higher peaks, which top out at almost 9,000 feet. More than three-quarters of the land base in this portion of the county is in public ownership, offering extensive hunting access. Game is plentiful and dispersed throughout the area for most of the year, concentrating in the lower elevations in winter when deep snows cover much of the landscape.

GMU 204 includes the eastern one-third of the district (from the Okanogan River east to the Okanogan County line) and is moderately rolling terrain, generally rising in elevation as you

move east. The vegetation changes from shrubsteppe near the Okanogan River to a mix of tall grass and conifer forest throughout the remainder of the unit. This portion of the district is roughly a 50-50 patchwork of public and private land, with the public lands generally being higher in elevation. Again, game is plentiful and dispersed throughout.

Weather in the Okanogan District can be quite variable and capable of changing quickly in the fall. Be prepared for everything from warm, sunny days to the possibility of winter temperatures and significant snow at higher elevations by the second week of October.

**Please be respectful of private land and treat landowners and their property the way you would want to be treated if roles were reversed.**



Chewuch River and Pasayten Wilderness – Photos by Scott Fitkin

Agency biologists will run a biological check and information station at the Red Barn in Winthrop both weekends of the modern firearm general deer season. We encourage hunters to stop and provide data to biologists whether they have harvested a deer or not. Data collected assists in assessing herd health and shaping population management.

## ELK

### GENERAL INFORMATION, MANAGEMENT GOALS, AND POPULATION STATUS

Overall, elk numbers are low in District 6. However, conditions vary noticeably between the east and west portions of Okanogan County. The western two-thirds of the district are not currently covered under a Washington Department of Fish and Wildlife (WDFW) elk herd management plan, and the existing harvest strategy (any elk general season) is designed to minimize elk numbers to prevent agricultural damage. As such, elk are quite scarce west of the Okanogan River and very difficult to find without extensive local knowledge.

The eastern portion of the district (GMU 204) is covered by the Selkirk Elk Herd Plan. Its four primary goals are: (1) to preserve, protect, perpetuate, manage, and enhance elk and their habitats to ensure healthy, productive populations and ecosystem integrity, (2) to manage this elk herd for a sustained hunting yield, (3) to manage elk for a variety of recreational, educational, and aesthetic purposes, including hunting, scientific study, cultural and ceremonial uses by Native Americans, biodiversity, wildlife viewing, and photography, and (4) to manage elk and elk habitat to minimize human conflicts and agricultural damage. More specifically, GMU 204 supports part of the Pend Oreille subherd population, where the current management objective is to gradually increase elk numbers while addressing the above four goals. As a result, this unit is now managed with an any bull harvest during general seasons. Elk are not currently abundant enough to warrant a survey effort in District 6, but observations suggest numbers continue to increase in GMU 204 and improve harvest opportunity accordingly.

2017 District 6 elk harvest statistics: [District 6 General Season Elk Harvest](#)

### WHICH GMU SHOULD ELK HUNTERS HUNT?

As noted above, GMU 204 is the only GMU in District 6 with a significant number of elk. Within this unit, elk tend to be most numerous in the area from Havillah through the Chesaw Wildlife Area, the Waconda Summit / Mount Annie area, and USFS lands bordering the Colville Reservation. In the rest of the district, finding animals is extremely difficult unless you have up-to-date knowledge on one of the few small bands of elk that wax and wane in the western portion of the county.

## DEER

### GENERAL INFORMATION, MANAGEMENT GOALS, AND POPULATION STATUS

District 6 supports perhaps the largest migratory mule deer herd in the state and Okanogan County has long been prized by hunters for its mule deer hunting. The district also supports significant numbers of white-tailed deer, particularly in GMUs 204 and 215. The District 6 deer management objective is for a stable to modestly increasing population within the social tolerance limits for nuisance and damage issues.



Okanogan District mule deer and white-tailed deer – Photos by Scott Fitkin


Surveys from the last couple of years indicate a modest decrease in deer populations in the wake of the recent extreme fires, severe droughts, and two modestly tough winters. However, as burned winter range continues to recover, landscape carrying capacity and deer numbers are expected to increase.

### **WHICH GMU SHOULD DEER HUNTERS HUNT?**


All units in District 6 support significant numbers of deer, include large blocks of accessible public land (except GMU 209), and offer good to excellent deer hunting opportunity. Mule deer are abundant throughout the county, with the highest densities occurring in the western two-thirds of the district.

Overall, white-tailed deer are less numerous than mule deer in Okanogan County, and in contrast to mule deer, white-tailed deer abundance generally increases as you move east in the district. The largest population is in GMU 204, where white-tailed deer comprise about half of the overall deer population. Although white-tailed deer numbers are less abundant in the western portion of the district, they are still found in most all drainages up to mid-elevations, particularly those with significant riparian vegetation. The highest concentrations in this area are in the Sinlahekin Valley and surrounding drainages. In many areas west of GMU 204 and outside of the Sinlahekin Wildlife Area, white-tailed deer frequent private lands. Prospective hunters wishing to target white-tailed deer may want to seek permission in advance of the season to access individual ownerships.

Hunters harvested 1,966 (1,707 bucks, 259 antlerless) deer in District 6 during the 2017 general seasons. This total is about a 25 percent dip below the 5-year average, but above the 10-year low of 1,811 animals taken in 2008. Correspondingly, the general season success rates for all weapon types fell below their respective five year averages and broke out as follows: Modern – 13 percent, Muzzleloader – 24 percent, Archery – 27 percent, and Multiple – 25 percent.



**Figure 1.** District 6 2017 general season hunter success by weapon type and GMU.



**Figure 2.** District 6 5-year average general season hunter success by weapon type and GMU.

GMU 204 (the district’s largest unit) yielded the greatest overall general season harvest of 615 animals. In the western portion of District 6, GMU 215 produced the most harvest with 274 deer taken.

For more information, see the 2017 District 6 deer harvest statistics: [District 6 General Season Deer Harvest](#)

## **WHAT TO EXPECT DURING THE 2018 SEASON**

Lower than average estimated fawn recruitment following the 2015-16 and 2016-17 winters likely means a dip in 2.5 and 3.5-year-old buck availability. However, the 2017 raw post season buck:doe ratio remained strong at 19 bucks per 100 does, and with almost half of those being greater than or equal to 3 point, older age class buck availability looks good. Overall, total general season harvest and success rates are anticipated to be around the 2017 numbers, somewhere above the 10-year low and below the 5-year average. Opportunities for older age class bucks during the permit seasons look good.

Several lightning-caused wildfires began in August in District 6, but the eventual impact of those is yet unknown. The district is currently in the midst of an extended hot and dry period. If this weather pattern continues, expect the high country to be drier than usual. If so, deer may be more concentrated on cooler and comparatively moister northerly aspects. Otherwise, expect deer to be widely dispersed during the general seasons. Regardless, expect significant campfire restrictions and check with local agencies for updates.

## **HOW TO FIND AND HUNT MULE DEER**

During the early general seasons, deer will generally be widely distributed on the landscape and not yet concentrated in migration areas or on winter range. Mature bucks in particular are often at high elevations in remote locations as long as succulent vegetation is available. In general, older, higher elevation burns, including the Tripod, Thirty-mile, Farewell, and Needles Fires, are producing high quality summer forage and are a good bet for significant deer activity. Although mule deer will use a variety of habitat types, they will often forage well into fairly open environments, particularly at dawn and dusk. As a result, they can often be glassed and stalked from considerable distance.

During the high hunt, deer will definitely still be spread across the landscape and are found in good numbers throughout the Pasayten Wilderness. Easier access to higher, more open country for hunters on foot is located at the Harts Pass and Iron Gate trailheads at the western and eastern ends of the wilderness respectively. For those with horses, the Andrews Creek offers access to good deer terrain further in. As mentioned above, the Billy Goat and Thirty-Mile trailheads are currently inaccessible due to road closures.

For Youth, Senior, and Disabled Hunters holding antlerless tags, does are spread across the landscape even more so than bucks during the general season, so permit holders should be able to find antlerless animals anywhere they have legal access.

During the late permit seasons, the majority of deer will have moved to winter range areas at lower elevations, often on more southerly slopes, to participate in the breeding season. In District 6, WDFW wildlife areas and immediately adjacent federal lands are good bets for high deer numbers in late fall, although in low snow years, some mature bucks may linger at higher elevations. In exceptionally mild years, hunters may have to go a bit higher than usual to find deer concentrations. Some GMU-specific recommendations for late mule deer permit holders are as follows:

GMU 215: Look for deer on the south facing slopes in the Toats Coulee drainage, open portions of the Sinlahekin Wildlife Area (SWA), and south facing slopes of the major drainages to the west of the Sinlahekin, including Cecil, Sarsapkin, and Sinalhekin creeks and their tributaries.

GMU 218: The Rendezvous Unit of the Methow Wildlife Area (MWA), and the Cub Creek, Buck Lake, and Lower Boulder Creek area of the Okanogan National Forest (ONF) are good bets.

GMU 224: Favorite spots are portions of the MWA and adjacent ONF lands in the southern portion of the unit. This includes more open habitat in drainages such as Pearrygin, Ramsay, Bear, Blue Buck, Beaver, and Frazier creeks.

GMU 231: Check out the Big Buck portion of the MWA, as well as the Virginia Ridge, Thompson Ridge, and Little Bridge Creek areas of the ONF.

GMU 233: The main unit and Pogue Mountain Unit of the Scotch Creek Wildlife Area (SCWA), the Carter Mountain Unit of the SWA, and public land in the Salmon Creek Drainage are good places to start.

GMU 239: The Texas Creek Unit of the MWA and the Chiliwist Unit of the SWA along with adjacent DNR land offer good opportunities. Upper portions of Finley, Benson, and Texas creeks on the ONF are also worth a look.

GMU 242: Look for deer on the Golden Doe Unit of the MWA and on south facing slopes on public land in the Libby Creek and Gold Creek drainages.

## **HOW TO FIND AND HUNT WHITE-TAILED DEER**

White-tailed deer are typically far less migratory than mule deer and generally favor brushier country with denser cover, primarily at lower and middle elevations. Look for white-tailed deer along stream drainages and in other areas with riparian vegetation or thick cover. Like mule deer, white-tailed deer are most active at dawn and dusk, but often don't venture as far into larger openings unless under the cover of darkness. Look for white-tailed deer in edge habitats where denser cover abruptly transitions into more open meadows. Many white-tailed deer hunters will wait patiently at a stationary position along an obvious game trail or the forest edge, often employing the use of a blind or tree stand.

During the late permit season, some white-tailed deer summering at modestly high elevations will move a little ways downslope, but most will be in the same areas they inhabited during summer. GMU-specific recommendations for late permit holders and late archery season in the western portion of the district are as follows:

GMU 215: White-tailed deer are abundant on the SWA and Chopaka Unit of the SCWA.

GMU 218: Look for deer in the Eight-mile drainage, along the Chewuch River, and in the lower half of the Rendezvous Unit of the MWA (despite the open habitat).

GMU 224: Brushier areas along Bear Creek, Upper Beaver Creek and its tributaries, and basin drained by the West Fork Salmon Creek west of Conconully are good bets.

GMU 231: Good possibilities include the huntable portion of the Big Valley Unit of the MWA, and the portion of the unit in the Twisp River Valley (north of the Twisp River Road). White-tailed deer can sometimes be encountered on the south slopes of the Big Buck Unit of the MWA as they move uphill off private land.

GMU 233: Despite the open terrain, the Happy Hill area of the SCWA is productive, along with the Buzzard Lake Unit of the SWA and adjacent DNR lands.

GMU 239: White-tailed deer can be found on ONF land in the South Summit area between Loup Loup Pass and Leecher Mountain and in wetter areas in the western portion of the Chiliwist Unit of the SWA and adjacent DNR lands.

GMU 242: Productive areas include the brushy areas along the river and in the northern half of the Golden Doe Unit of the MWA, as well as the valley bottom of the Twisp River drainage. Public land along Libby and Gold Creeks is also a possibility.

Antlerless white-tailed deer permit holders should look for animals in the same areas mentioned above with the added expectation of a few more deer in the higher reaches of areas like the Twisp River and Eight-mile drainages than might be expected during the late season.

## DEER AREAS

For those hunters with second deer permits in Deer Areas 2012 -2016, remember that those permits are good **only on private land**. Permit holders are responsible for making contact with private landowners to secure hunting access.

# BLACK BEAR

## GENERAL INFORMATION, MANAGEMENT GOALS, AND POPULATION STATUS

Black bears are abundant and well distributed throughout District 6 and are managed for sustainable harvest and diverse age structure. Monitored demographic parameters suggest the local population and associated harvest appear to be relatively stable, so hunting prospects in the district should be good.

For hunters pursuing black bear in the northern Cascades, it is critical to positively identify the bear species, as endangered grizzly bears potentially also inhabit these areas. WDFW's website features some interactive training materials on how to tell the difference between black and grizzly bears. [Click here](#) to view the Interactive Bear Identification Program and take the Bear Identification Test.

## WHICH GMU SHOULD BEAR HUNTERS HUNT?

All GMUs in the Okanogan District provide good black bear hunting opportunity. In 2017, hunters posted a success rate of 7.4 percent and harvested 103 black bears from the western portion of the district in the Okanogan Bear Management Unit (BMU 5), both of which were right at the 5-year averages. GMU 204 in the Northeastern BMU yielded 47 animals, just below the 5-year average.

For more information, see the 2017 District 6 black bear harvest statistics:

- [Okanogan BMU Black Bear Harvest](#)
- [Northeastern BMU Black Bear Harvest](#)




Figure 3. Okanogan District 5-year average black bear harvest success rates by GMU.

### WHAT TO EXPECT DURING THE 2018 SEASON

In general, at the beginning of bear season, animals are likely to be found at middle elevations in wetter areas where berries are peaking. As the season progresses, expect bears to follow the ripening berries to higher elevations. As we move into fall, animals will range over a wider gradient to take advantage of a variety of late season food sources. This year, after a late onset for service berries, the hot weather is accelerating the development of the later crops and timing should be about average as bear season arrives.



Okanogan District black bear – Photo by Scott Fitkin

## COUGAR

### GENERAL INFORMATION, MANAGEMENT GOALS, AND POPULATION STATUS

The District 6 cougar population is healthy and dispersed throughout the landscape. In the Okanogan District, cougars are now managed by a harvest guideline at the individual GMU level to better promote stable population structure and high quality sustainable harvest, while also minimizing human-cougar conflicts.

Cougars follow the deer herds, which means they will be spread throughout the district through late October and concentrate more at lower elevations as deer move to winter range. Much cougar foraging activity takes place at night, so the best opportunities to spot the cats on the move are at dawn and dusk.



Cougar – Photo by Scott Fitkin

**WHICH GMU SHOULD COUGAR HUNTERS HUNT?**

All Okanogan District GMUs support cougars and are open to hunting. After January 1, individual PMUs (one or more GMUs) close on short notice once the harvest guideline has been reached, and hunters are responsible for knowing if a unit is open or closed. This information is available on the WDFW hotline (1-866-364-4868) or [online](#).

<b>PMU Hunt Area (GMUs)</b>	<b>Harvest Guideline</b>	<b>2013-2014 Harvest</b>	<b>2014-2015 Harvest</b>	<b>2015-2016 Harvest</b>	<b>2016-2017 Harvest</b>	<b>2017-2018 Harvest</b>
203	4-6	0	0	0	0	0
204	6-8	5	1	7	2	6
209, 215	3-5	2	4	5	3	5
218, 231	4-6	3	2	2	3	0
224	2-3	2	1	1	3	2
233,239	3-4	0	1	3	5	1
242,243	6-7	4	3	4	3	2

**Table 1.** District 6 cougar harvest guidelines and 5-year harvest by GMU.

As is typical, last season’s harvest fell short of the guidelines in several GMUs, and control-related mortality remained modest. As a result, cougar numbers should be robust and hunting opportunities in District 6 should be good in 2018-19. A summary table of the harvest guideline by PMU is presented below.

For more information on District 6, see the 2017-2018 cougar harvest statistics for PMUs 13-18: [State-wide cougar harvest by PMU](#).


## WATERFOWL

### GENERAL INFORMATION

The Okanogan District offers modest waterfowl hunting opportunities as compared to many other areas of the state. The largest concentrations of birds occur at the southern edge of District 6, at the mouth of the Okanogan River and on the Columbia River. The main stem of the Okanogan and Upper Similkameen rivers and the larger lakes and potholes in the Okanogan Watershed are good secondary sites. Good public river access can be found at the Washburn Island Unit of the Wells Wildlife Area, the Driscoll-Eyhott Island Unit of the Sinlahekin Wildlife Area, and the Similkameen-Chopaka Unit of the Scotch Creek Wildlife Area.



Barrows golden eye pair – Photo by Scott Fitkin



Canada geese in a Methow Valley grain field – Photo by Scott Fitkin

Water levels in local potholes remained high again in 2018, but are coming down a bit with the prolonged hot and dry weather. River levels are currently running around average or a little below normal and are likely to be similar going into the hunting season. Aside from water levels, waterfowl hunting opportunities are mostly dependent on the number of migrants coming from Canada and Alaska and how long water remains ice-free throughout the district.

For more information, see the 2017 District 6 waterfowl harvest statistics: [Okanogan County Small Game Harvest](#)

## FOREST GROUSE

### SPECIES AND GENERAL HABITAT CHARACTERISTICS


The Okanogan supports strong populations of ruffed, dusky (blue), and spruce grouse, which are found throughout the forested areas of the district. Ruffed grouse are generally associated with deciduous tree cover at lower to middle elevations, particularly in riparian habitats. During hunting season, dusky (blue) grouse are generally found in the mid to upper elevation conifer forests, often moving to ridges as snow begins to accumulate. Spruce grouse are located in higher elevation conifer forests throughout the district on a year-round basis.

Forest grouse populations are likely below historical norms within the boundaries of recent wildfires, including the massive Carlton Complex, Okanogan Complex, Tunk Block, and Tripod fires. These fires burned in some of the best and most densely occupied forest grouse habitat in the district. However, grouse habitat within the burns is improving annually (particularly in the Tripod Fire area), and bird numbers outside of burned areas appear to be relatively stable.

### HARVEST TRENDS AND 2018 PROSPECTS

The sprawling landscape of Okanogan yielded a mixed harvest of 6,788 dusky, ruffed, and spruce grouse, down 8 percent over last year. However, grouse harvest per unit effort increased 31 percent in 2017, suggesting the cool and wet spring weather was not a factor last year after all. Spring conditions appeared favorable this year and anecdotal observations suggest dusky grouse productivity may have been good, creating expectations for harvest and success rates equal to or better than last season.

For more information, see the 2017 District 6 forest grouse harvest statistics: [Okanogan County Small Game Harvest](#) .



Male dusky grouse and female spruce grouse – Photos by Scott Fitkin

## PHEASANTS

### SPECIES AND GENERAL HABITAT CHARACTERISTICS

Pheasants occur at low densities and in a patchy distribution throughout the Okanogan Watershed portion of District 6, with the majority of harvested birds coming from pheasant release sites. This year, pheasants will again be released at the Bureau of Reclamation’s Hegdal and Kline sites, and at the Chilliwist Unit of the Sinlahekin Creek Wildlife Area. What little wild production exists within the county comes mostly from private land. Hunters should seek permission in advance of the season to access private property.

The release sites are mapped on the [GoHunt](#) website, as well as in the Eastern Washington Pheasant Release booklet found here: [Eastern Washington Pheasant Release Program](#). Hunters are reminded that nontoxic shot is required for ALL upland bird hunting on ALL pheasant release sites statewide.

Hunters bagged 706 pheasants last year in Okanogan County. Both the total harvest and harvest per unit effort were down from the previous year, but right at the 5-year average.

For more information, see the 2017 District 6 pheasant harvest statistics: [Okanogan County Small Game Harvest](#)



Pheasant release – Photo by WDFW

## QUAIL

### SPECIES AND GENERAL HABITAT CHARACTERISTICS

Quail are locally abundant and widespread throughout the district’s lower elevation shrubsteppe and open pine forest habitats. District 6 wildlife areas offer good access to quail habitat. Anecdotal observations this spring and summer suggest quail production has been good again this year, with some birds having multiple clutches, suggesting 2018 harvest will likely be similar to last year. In 2017, hunters took 8,583 quail in Okanogan County. Harvest and harvest per unit effort were up 17 percent and 22 percent respectively as compared to the 5-year average.

For more information, see the 2017 District 6 quail harvest statistics: [Okanogan County Small Game Harvest](#)

## TURKEYS

### GENERAL DESCRIPTION

Turkeys are found in scattered groups throughout the district and often concentrate on private land near agriculture areas. Prospective hunters should seek permission in advance of the season to access private land. The fall turkey permit season occurs within GMUs 218-231 and 242, with the majority of the birds being located in the latter two units. In recent years, winter conditions and declines in supplemental feeding by private individuals have reduced turkey numbers

substantially in the Methow Valley, although most lower-elevation drainages in GMU 242 still harbor birds.

## CHUKAR AND GRAY PARTRIDGE

### GENERAL DESCRIPTION

In general, gray partridge populations are widely distributed and patchy throughout the district's shrubsteppe habitats, but appear to be increasing in numbers and distribution over time. Birds are seen frequently on the Indian Dan, Chiliwist, and Methow wildlife areas. Scattered groups of chukars are found in the rocky areas in lower elevations of District 6. The steep hills along the Similkameen River in the north part of the Okanogan Valley hold good numbers of birds.

Combined harvest of chukar and gray partridge in 2017, as well as catch per unit effort, rose substantially from the previous season suggesting populations rebounded nicely as winter conditions eased. This year's average winter combined with good forage production this spring suggest another good year is in the offing. For more information, see the 2017 District 6 partridge and chukar harvest statistics: [Okanogan County Small Game Harvest](#).



## DOVE

### GENERAL DESCRIPTION

Similar to chukar and partridge, dove harvest and harvest per unit effort were up significantly in 2017. The past moderate winter and good forage growth this year make the outlook good for doves in 2018. Look for doves in planted food crops in the Sinlahekin and at lower elevations on other public land. Hunting success will depend on warm weather keeping the birds in the area through the season.

For more information, see the 2017 District 6 mourning dove harvest statistics: [Okanogan County Small Game Harvest](#).