



Pacific Salmon and Wildlife

Ecological Contexts, Relationships, and Implications for Management

Special Edition Technical Report



PREFACE

This Updated Special Edition Technical Report is a product of the project: *Wildlife –Habitat Relationships in Oregon and Washington* (Johnson and O’Neil 2001). The integration of the information on the ecology and management of species in terrestrial, freshwater, and marine environs has been a focus of the project since its inception. For the first time, this Special Edition Technical Report synthesizes fundamental and crucial information linking salmon with wildlife species and the broader aquatic and terrestrial realms in which the co-exist. Readers will find this scientifically-robust report to greatly strengthen our collective understanding of the role that salmon play in the population so Pacific Northwest wildlife species, the ecology of freshwater ecosystems, and how management activities such as hatcheries and harvest can impact these aspects.

We thank the authors who contributed important time and effort to the making of this report. Efforts on this Special Edition Technical Report have been supported by a large number of entities, including the 34 *Project Partners* and *Contributing Sponsors of Wildlife –Habitat Relationships in Oregon and Washington*:

Birds of Oregon Project; Birds of Washington Project; Confederated Tribes of the Warm Springs; Donavin Leckenby; Environmental Protection Agency – Corvallis Lab.; Federal Highways Administration; Fish and Wildlife Information Exchange; National Fish and Wildlife Foundation; Northwest Power Planning Council; Oregon Cooperative Wildlife Resources Institute; Oregon/Washington Partners in Flight; Pacific States Marine Fish Commission; Quileute Indian Tribe; Rocky Mountain Elk Foundation; Society for Northwestern Vertebrate Biology; USDA Forest Service; USDI Bureau of Land Management;’ USDI Fish and Washington Community, Trade, and Economic Development’; Washington Department of Natural Resources; Washington Department of Transportation; Washington Forest Protection Association; Weyerhaeuser Company; Wildlife Management Institute. Contribution Sponsors are Paul F. And Teresa J. Roline; Oregon Chapter of The Wildlife Society; and the WA Chapter of the Wildlife Society. The Washington Department of Fish and Wildlife and the Northwest Habitat Institute are the lead organizations on this project.

In particular, we thank the Multi-Species Framework Project/Northwest Power Planning Council, Washington Department of Natural Resources, Washington Department of Fish and Wildlife, and the Northwest Habitat Institute for their important contributions to the development and printing of this Special Edition Technical Report.

Additional copies of this report can be acquired from the web page: <http://www.wa.gov/wdfw/hab/salmonwild/>, or contacting David H. Johnson, WDFW, Habitat Program, 600 Capitol Way N., Olympia, WA 98501-1091; email: johnsdhj@dfw.wa.gov

Special changes have been made to Table 8 to correct numerical errors in previous version.

Cover photo: Glaucous-winged gull feeding on chum salmon carcass: Photo by C. Jeff Cederholm. Cover border: Northwest Indian artwork by Bill Martin of the Makah Tribe, Neah Bay, Washington. Cover layout and design: Charley Barrett of the Northwest Habitat Institute, Corvallis, Oregon.

PACIFIC SALMON AND WILDLIFE - ECOLOGICAL CONTEXTS, RELATIONSHIPS, AND IMPLICATIONS FOR MANAGEMENT

C. Jeff Cederholm, Washington Department of Natural Resources, Olympia, WA. ¹

David H. Johnson, Washington Department of Fish and Wildlife, Olympia, WA.

Robert E. Bilby, National Marine Fisheries Service, Seattle, WA.

Lawrence G. Dominguez, Washington Department of Natural Resources, Olympia, WA.

Ann M. Garrett, Private Consultant, Kirkland, WA.

William H. Graeber, Washington Department of Natural Resources, Olympia, WA.

Eva L. Greda, Washington Department of Fish and Wildlife, Olympia, WA.

Matt D. Kunze, Private Consultant, Spokane, WA.

Bruce G. Marcot, United States Forest Service, PNW Lab, Portland, OR.

John F. Palmisano, Private Consultant, Portland, OR.

Rob W. Plotnikoff, Washington Department of Ecology, Olympia, WA.

William G. Percy, Oregon State University, Corvallis, OR.

Charles A. Simenstad, University of Washington, Seattle, WA.

Patrick C. Trotter, Private Consultant, Seattle, WA.

Recommended Citation:

Cederholm, C. J., D. H. Johnson, R. E. Bilby, L.G. Dominguez, A. M. Garrett, W. H. Graeber, E. L. Greda, M. D. Kunze, B.G. Marcot, J. F. Palmisano, R. W. Plotnikoff, W. G. Percy, C. A. Simenstad, and P. C. Trotter. 2000. Pacific Salmon and Wildlife - Ecological Contexts, Relationships, and Implications for Management. Special Edition Technical Report, Prepared for D. H. Johnson and T. A. O'Neil (Managing directors), Wildlife-Habitat Relationships in Oregon and Washington. Washington Department of Fish and Wildlife, Olympia, Washington.

ACKNOWLEDGMENTS

We thank the Washington Departments of Fish and Wildlife (WDFW) and Natural Resources (WDNR) for supporting this project. We appreciate greatly the thoughtful review given by Peter Bisson of the United States Forest Service; Jim Lichatowich, Private Consultant; Hal Michael, Jr., of the WDFW; Tom Mumford of WDNR; Pat Slaney of the British Columbia Watershed Restoration Program; and Mary F. Willson, of the Great Lakes Program of The Nature Conservancy. We also thank Luis Prado of WDNR and Darrell Pruett of the WDFW for graphic artwork on some of the figures in this report and Billie Lynn Wyckoff for help and considerable patience with putting this publication together.

¹C. Jeff Cederholm. Washington Department of Natural Resources, Business Systems Support Division. 1111 Washington St. SE. PO Box 47014, Olympia, WA 98504-7014. Phone: (360) 902-1609, FAX: (360) 902-1789, E-mail: jeff.cederholm@wadnr.gov.

TABLE OF CONTENTS

ACKNOWLEDGMENTS – i

TABLE OF CONTENTS – ii

ABSTRACT - iii

INTRODUCTION - 1

GENERAL SALMON LIFE HISTORY - 7

FRESH WATER AND TERRESTRIAL HABITAT RELATIONSHIPS OF SALMON 1 - 18

 Freshwater Habitat - 18

 Riparian Habitat - 20

 Habitat Forming Processes - 23

 Beaver - 24

ESTUARY HABITAT - 26

OCEAN HABITAT - 29

ECOLOGICAL RELATIONSHIPS OF SALMON - 30

Macroinvertebrates - 30

 Freshwater Macroinvertebrates And Salmon – 30

 Structure And Function In Macroinvertebrate Communitie - 30

 Physical and Chemical Influences On Macroinvertebrate Distribution - 30

 Significance Of Macroinvertebrate Life Cycles - 31

 Aquatic Macroinvertebrates As A Food Source for Salmon - 31

 Salmon as a Food Source For Aquatic Macroinvertebrates - 32

Vertebrate Wildlife - 34

 Vertebrate Wildlife And Salmon - 34

 Wildlife Species With A Relationship To Salmon - 35

 Wildlife Response To Salmon Congregations - 37

 Review Of Wildlife Relationships By Salmon Life Stages - 39

 Key Ecological Functions (KEFs) Provided To The Ecosystems Through Salmon-Wildlife Interactions - 43

SALMON FISHERIES - 49

 Spawning Escapement Goals - 56

UNDERSTANDING SALMON RELATIONSHIPS - 59

 Salmon As A Key Linkage In Biodiversity And Productivity - 59

 Biomass of Salmon Runs As An Energy Source - 60

 Pacific Salmon Provide Key Ecological Functions - 65

 Salmon As Vectors In Broader Nutrient Cycling - 66

 Maintaining the Salmon Link in Nutrient Cycling - 68

WITH AN ECOSYSTEM PERSPECTIVE IN MIND - WHERE DO WE GO FROM HERE? - 72

 Desired Future Condition - 74

IMPLICATIONS FOR FISHERIES MANAGEMENT - 75

LITERATURE CITED - 80

APPENDIX TABLES I-VII - 115

ABSTRACT

There are seven indigenous salmon and trout of the genus *Oncorhynchus* in Washington and Oregon (chinook, coho, chum, sockeye, and pink salmon, and steelhead and cutthroat trout), for this paper we will collectively call them salmon. Their habitat extends from the smallest inland streams to the vast North Pacific Ocean, an area of freshwater, estuarine, and ocean habitats in excess of 4 million km². Due to past commercial fisheries, habitat loss, hatchery problems, and more recently a changing ocean environment, salmon populations have shown substantial decline over the past several decades. Many salmon stocks in Washington and Oregon are now listed as either threatened or endangered, under the Federal Endangered Species Act.

Early in the 1900's and up until relatively recently, commercial fishing permanently diverted massive quantities of nutrients away from Washington and Oregon rivers, and their respective fish and wildlife inhabitants. Recent calculations by Gresh et al. ^{174a} indicate that only 3 percent of the marine-derived biomass once delivered by anadromous salmon to the rivers of Puget Sound, the Washington Coast, Columbia River, and the Oregon Coast is currently reaching those streams. There have also been many other losses of salmon habitat during this period caused by: river channel clearing and channelization, log driving and splash damming, extensive land clearing, major water diversions, livestock grazing, mining runoff pollution, logging road associated erosion and removal of the old growth forest, filling and diking of wetlands and estuaries, hydroelectric dam development, urban runoff, water and sediment contamination with toxicant, and recently recognized human induced oligotrophication of waterways. Over fishing and habitat degradation, together with a background of a changing ocean environment, have cumulatively reduced stock resilience. A century of hatchery programs have failed to rebuild the wild runs, and in many cases, likely contributed to their further declines. Modern salmon management techniques have become highly sophisticated, however, they have not been able to keep pace with the salmon population declines.

The life history of anadromous salmon covers time spent in freshwater, estuaries, and the ocean. Freshwater habitats are mainly used for spawning, incubation and juvenile rearing; estuaries are where juveniles put on critical rapid growth and make important osmoregulatory adjustments as they transition between fresh and saline waters; and the ocean is where significant feeding results in most of the body mass of the returning adults. Throughout their life salmon feed on a wide variety of prey organisms, including many kinds of freshwater and marine invertebrates and fishes; and at the same time, are fed upon by a wide variety of invertebrate and vertebrate predators and scavengers.

Juvenile salmon are known to feed directly on salmon carcass flesh, salmon eggs, and aquatic macroinvertebrates that may have previously fed on salmon carcasses. Research has uncovered significant contributions of nutrient from spawning salmon to the collector-gatherer macroinvertebrate community. Caddisflies, stoneflies, and midges are involved in processing the microbially conditioned salmon carcass flesh. Increase in aquatic macroinvertebrate density from the introduction of salmon carcasses stimulates feeding by early life stages of select salmon species. Other stages of the salmon life cycle also contribute to the macroinvertebrate food base, such as some stonefly nymphs, when they scavenge dead pink and chum

Of this group of wildlife species, 9 species had a *Strong-Consistent* relationship, 58 a *Recurrent* relationship, 25 an *Indirect* relationship, and 65 had a *Rare* relationship (the tally is more than 138 because 19 species have more than one type of relationship with salmon). These species were further examined as to the life cycle stage of salmon to which they were linked. The five salmon life stages, and the number of wildlife species associated with each (in parenthesis) were: *Incubation* (23); *Freshwater Rearing* (49); *Saltwater* (63); *Spawning* (16); and *Carcasses* (83) (this tally of wildlife species totals more than 137 because 66 species of wildlife are associated with salmon at several life stages).

Salmon act as an ecological process vector, important in the transport of energy and nutrients between the ocean, estuaries, and freshwater environments. The flow of nutrients back upstream via spawning salmon and the ability of watersheds to retain them plays a vital role in determining the overall productivity of salmon runs. As a seasonal resource, salmon directly affect the ecology of many aquatic and terrestrial consumers, and indirectly affect the entire food web. The challenge for salmon, wildlife, and land managers is to recognize and account for the importance of salmon not only as a commodity resource to be harvested for human consumption, but also for their crucial role in supporting overall ecosystem health. It is also important that naive view of wildlife as only consumers of salmon be abandoned. Many species of wildlife for which hard earned environmental laws and significant conservation efforts have been established (e.g., grizzly bears, bald eagles, river otters, killer whales, beaver), play key roles in providing for the health and sustainability of the ecosystems upon which salmon depend. As the health of salmon populations improves, increases in the populations of many of the associated wildlife species would be expected. Salmon and wildlife are important co-dependent components of regional biodiversity, and deserve far greater joint consideration in land-management planning, fishery management strategies, and ecological studies than they have received in the past.

Most measurements in this text are in metric system, however, conversion to English are given in Table 1.

Table 1. Metric to English Conversions.

Metric	English	To Convert From Metric to English Multiply Metric by:
millimeter (mm)	inch (in)	0.039
meter (m)	feet (ft)	3.281
square kilometer (km²)	square mile (mi²)	0.386
kilometer (km)	mile (mi)	0.621
gram (g)	ounce (oz)	0.035
kilogram (kg)	pounds (lbs)	2.205
cubic meter (m³)	cubic yard (yd³)	1.308
metric ton (mt)	ton (t)	1.102

INTRODUCTION

The landscapes of Washington and Oregon at first glance appear to have some disconnect between the terrestrial and ocean environments. Yet, abundant rivers and streams flow from the interior to the coastal zones, actively connecting the freshwater, estuarine, and ocean systems. Within these environments there are countless abiotic and biotic processes which form a highly integrated ecosystem.



*Plate #1. The White River
flowing from Mt. Rainier, WA.
(Photo by: Larry Dominquez)*

Key inhabitants include wild anadromous Pacific salmon (*Oncorhynchus* spp.) (anadromous fishes are those that spend much of their lives feeding in the ocean and migrate to freshwater to breed), 605 identified common vertebrate wildlife species, and numerous species of macroinvertebrates and other fishes. Complex relationships have evolved within and between anadromous salmon and these inhabitants that may be important for maintaining this ecosystem. Past highly exploitive fisheries, poor land use practices, an over-reliance on salmon hatcheries, and a changing ocean environment, have all contributed to many salmon stock declines in these states³⁶⁷. It has been suggested that future salmon conservation will need to take an ecosystem approach if wild stocks are to survive⁴⁸⁵. The purpose of this paper is to identify known relationships between wild salmon and wildlife, discuss the ecological context of these relationships, and to suggest new ways of managing the salmon resource with an ecosystem perspective in mind.

We define wild salmon as indigenous species that are the progeny of streambed spawners. This definition is used to distinguish wild salmon from hatchery (artificially) propagated salmon. The genus *Oncorhynchus* includes both salmon and trout, however for our purpose we collectively refer to them simply as salmon. Wildlife are divided into two main categories, indigenous macroinvertebrates (aquatic and terrestrial) and vertebrates (amphibians, reptiles, fishes, birds, and mammals) found in Washington and Oregon.

It is important to recognize that the ecosystem of Washington and Oregon salmon can be hemispheric in scale; reaching from local inland watersheds, where spawning occurs, all the way to ocean feeding grounds north of the Aleutian Islands of Alaska, west to the Asian side of the Pacific Ocean, and back again. This ecosystem spans an area of freshwater and ocean habitat in excess of 4 million km². The essence of the salmon is that it links together what humans generally consider distant, diverse, and separate ecosystems, and relatively long time spans.



Plate #2. Indian fishing for salmon at Celilo Falls on the lower Columbia River. (Photo by: Archive of The Spokesman Review, Spokane, WA).

Scientific knowledge of salmon in Washington and Oregon was preceded by a rich legacy of aboriginal culture, which wove them into everyday life^{98, 431, 379}. Salmon were an important food staple and a basis of many legends of the native people of these states, particularly those that lived along rivers and marine areas. Salmon were consumed by natives in large quantities, for example, Craig and Hacker^{109 cited in 379} calculate that pre-contact catches of salmon in the Columbia basin alone ranged between 4.5 and 5.6 million fish annually. Most of the salmon caught at that time were consumed within their respective river drainage and some were traded with distant tribes. Columbia River tribal records indicate that salmon were transported long distances inland, including trade routes over the Continental Divide.

“The Wishram and Wasco (tribes along the lower Columbia River near Celilo Falls) seem to have been the focal point in the most extensive trade network in the plateau -- one that reached to the mouth of the Columbia and out onto the plains east of the Rockies. They traded dried fish (salmon) for bison hides and other commodities that originated on the plains.”^{177 cited in 379}. Some of the earliest Euro-Americans to view Pacific salmon traveled to the Northwest with the Lewis and Clark expedition in 1805. Near the confluence of the Columbia and Snake rivers, they observed salmon in unimaginable abundance: *“The number of dead Salmon on the Shores & floating in the river is incredible (sic) to say...”*^{William Clark in 1805, p 252 cited in 119}.

European settlement and commercial development of Washington and Oregon brought significant habitat problems for the salmon; resulting in many physical, chemical, hydrological, and biological modifications to the environment. Varied effects on salmon habitat are often interrelated in complex

ways and the effects of various activities and ecosystem modifications can be cumulative⁵¹³. Some of the more harmful habitat losses caused by man have been: river channel clearing and channelization, log driving and splash damming, extensive land clearing, major water diversions, livestock grazing, mining runoff pollution, logging road associated erosion and removal of the old growth forest, filling and diking of wetlands and estuaries, hydroelectric dam development, urban runoff, water and sediment contamination with toxicants, and recently recognized human induced oligotrophication of waterways^{208, 296, 101, 245, 453, 485, 365, 174, 59, 16, 543}.




Plate #3. Clearcut logging along Matheny Creek, Washington. (Photo by: Jeff Cederholm).



Fishery exploitation of Columbia River salmon by Euro-Americans became a major factor after the middle to late 1800s. To ensure primary access to the salmon, commercial fisheries were strategically located downstream of popular Indian fishing grounds. The principle means used to catch the salmon were gillnets, traps, seines, and fish wheels^{121, 458, 274}. It was reported that "...on a single spring day in 1913 the Seufert brothers' wheel no.5 turned a record catch

Plate #4. Urbanization and habitat loss in the Puyallup River estuary and floodplain, Tacoma, Washington. (Photo by: Washington Department of Natural Resources Photo and Mapping section).




Plate #5. Grand Coulee Dam, an impassable hydroelectric dam on the Columbia River, Washington. (Photo by: Larry Dominguez)

of 70,000 pounds”¹⁰³. After the 1870s and up to the early 1900s, the Columbia River salmon fishery grew from 1 to 40 canneries^{478, 368 cited in 365}.

Fish wheels were prohibited on the Columbia River after 1935³⁶⁵. Commercial landings of Columbia River salmon and steelhead peaked between 1880 and 1930, and then went into a long term decline up to present times³⁶⁵. Depletion of the prime spring and summer chinook probably started earlier, however, as the fishery shifted to the less desirable coho and fall chinook²⁷⁵. One estimate of annual pre-Euro-American salmon and steelhead run size for the Columbia River ranges between 0.2 and 16.3 million fish.

Early attempts to increase salmon catches using salmon hatcheries began as early as the 1870s, when concerns about over fishing led the Oregon and Washington Fish Propagating Company to construct a salmon-breeding station on the Clackamas River³⁶⁵. By the 1960s, with the advent of the

Plate #6. Fish wheel scow on the Columbia River at the Cascades, loaded with blueback (sockeye) salmon, c. 1895. (Photo by: Oregon Historical Society: OrHi-214, reproduction number 340)






Plate #7. Salmon Catch at the Seattle Wharf. (Photo by: Washington State Historical Society, Tacoma, WA. Negative No. 1994.123.121)

Oregon Moist Pellet medicated food, hatchery salmon production increased dramatically. Total annual Columbia-Snake River system hatchery production (Washington, Idaho, and Oregon) reached 216 million smolts in 1989³⁹⁰. The catches of salmon in the Columbia River fisheries in these times were high relative to recent times; however, they were largely hatchery fish. In the early 1990s about 95% of the coho, 70% of the spring chinook, more than 50% of the fall chinook, more than 80% of the summer chinook, and 70% of the steelhead production in the Columbia-Snake River system were of hatchery origin³⁹⁰. By the middle




Plate #8. Canned salmon at the Apex fish company (1913). Photo by: Curtis. Washington State Historical Society. Negative No. 27683

1990s there were well over 100 state, federal, tribal and private salmon hatcheries in Washington and Oregon. With the increasing reliance on artificial propagation, concerns became greatly heightened that contemporary hatchery programs were having negative effects on the genetic diversity and persistence of wild populations, and that increasing releases of hatchery fish could not override other factors contributing to the overall decline of salmon ^{307, 365}. The history of artificial propagation reveals a recurring cycle of technological optimism followed by pessimism.

While many attempts have been made at remedying the threats of habitat loss, over fishing, and hatchery impacts, they have not been enough to prevent the widespread decline of wild salmon stocks in these states. Recent publications have chronicled the low abundance of wild salmon stocks along the Pacific Coast in the lower 48 states ^{367, 53, 542, 387, 181}. In 1991, the American Fisheries Society ³⁶⁷ published a list of 214 naturally spawning stocks of salmon, steelhead, and cutthroat from California, Oregon, Idaho, and Washington, including: 101 stocks at high risk of extinction, 58 at moderate risk of extinction, 54 stocks of special concern, and one classified as threatened under the Federal Endangered Species Act (ESA) of 1973. Fifty-eight of these stocks occur along the Oregon coast, 41 along the Washington coast and in Puget Sound, and 76 within the Columbia River basin. In spite of past salmon habitat degradation and over fishing, however, some stocks remain healthy ²²¹. Since 1990, the National Marine Fisheries Service (NMFS) has received a number of petitions to list Pacific salmon stocks as threatened or endangered under the ESA, and the first salmon stock of this area to be listed as endangered was the Snake River sockeye, in November 1991 ^{366a}.

GENERAL SALMON LIFE HISTORY

There are seven species of Pacific salmon and trout of the genus *Oncorhynchus* in Washington and Oregon, and they include: chum (*O. keta*), pink (*O. gorbuscha*), sockeye (*O. nerka*), chinook (*O. tshawytscha*), and coho salmon (*O. kisutch*); and rainbow (called steelhead when anadromous) (*O. mykiss*) and coastal cutthroat trout (*O. clarki clarki*). Some of these species, including the sockeye salmon (kokanee) and rainbow and cutthroat trouts, have both anadromous and nonanadromous forms. Salmon life history patterns follow a basic theme (Figure 1).




Figure 1. Generalized anadromous and nonanadromous (resident) Pacific salmon life histories, showing freshwater, estuary, and ocean components (the original diagram was from Nicolas and Hankin³⁷² and later modified by Spence et al.⁴⁸⁵).

Salmon evolved in habitats that are typically characterized by accessible cool, clean water with abundant woody debris or other forms of cover, relatively clean spawning gravels, food, and a balanced population of predators. In the temperate ecosystem of the Pacific Northwest, the freshwater environment is less productive than the ocean environment, particularly estuaries and coastal upwelling zones¹⁷⁹, therefore Pacific salmon evolved an ocean feeding phase in their life history¹⁷⁹. A typical anadromous salmon life history has five main stages: (1) spawning and egg incubation, (2) freshwater rearing, (3) seaward migration, (4) ocean rearing, and (5) return migration to freshwater to spawn and the deposition of marine derived nutrients into the freshwater ecosystem (Figure 1). Each species has slightly different temporal varieties of the anadromous life history (Figure 2). Simply put, life history means: "...what the salmon do, where they do it, when they do it, and how they do it"²⁷⁵.




Figure 2. Temporal phases of the anadromous Pacific salmon life history; while in freshwater, estuary, and ocean environments.

The various stages of development may have many different timings, depending on species and location, as typified for Puget Sound (Skagit-Samish Basins) in Figures 3-A, coastal Washington (Queets-Quinault Basins) in Figure 3-B, and coastal Oregon in Figure 3-C. The nonanadromous (resident) life history is typified by spawning and rearing in freshwater without going to sea.

Timing of salmon freshwater life phases in the Skagit-Samish Basins WRIA 03-04 (DeShazo 1980; Williams et al. 1975; WDF et al. 1993; WDFW 1996).




Figure 3A. Timing of salmon freshwater life phases in the Skagit-Samish Basins WRIA 03-04 ^{118, 556, 542, 541}.



Figure 3B. Timing of salmon freshwater life phases in the Queets-Quinault Basin WRIA 21 ^{162, 402, 542, 541}.

Timing of salmon freshwater life phases in drainages of the Oregon Coast (Busby et al. 1996; Johnson et al. 1999; Myers et al. 1998; Nicholas and Hankin 1988; T. Nickelson, personal communication; Sumner 1962; Weitkamp et al. 1995)



Figure 3C. Timing of salmon freshwater life phases in drainages of the Oregon Coast ^{79, 234, 356, 372, 505, 549.}

The chum, pink, sockeye, chinook, and coho salmon all die after spawning just once, a life history strategy known as *semelparity*³²². This life strategy has evolved because of the need to have a greater portion of the energy obtained from ocean feeding devoted to gamete production and juvenile survival. Consequently, survival after spawning no longer offered an advantage to these species³²². Conversely, the death of the spawning run offered a substantial survival advantage to the overall population. By enriching the environment of the juveniles their growth and survival rates could be increased. The *iteroparous*, repeat spawning strategy, typical of the rainbow and cutthroat, probably occurred in the headwater reaches of larger rivers, where nonanadromous populations could be maintained year round. These fish generally were smaller in size, less fecund, and had sparser distribution and lower abundance than the anadromous forms. However, by retaining iteroparity, calamitous losses of young due to floods or drought, could be compensated for in subsequent breeding seasons³²².

Chum, pink, sockeye, chinook, and coho salmon all spawn sometime between August and February, at a time when stream flows are increasing and water temperatures are declining. Rainbow and cutthroat spawn between January and June, when stream flows are decreasing and water temperatures are increasing. For



Plate #9. Chum salmon spawning in Kennedy Creek, Washington. (Photo by: Jeff Cederholm)

successful development of eggs to occur, the gravel should be relatively stable and clean of fine sediments. Pacific salmon are able to clean gravels by purging them of fine sand and silt particles during redd (spawning nest) excavation; but subsequent sediment transport processes and bedload flux can return this environment to the pre-spawning condition^{414, 401a}. After approximately 2-4 months of incubation, salmon fry swim up through the gravel and emerge into the stream. Actual emergence time will depend on species and race of fish, for example chum and chinook emerge in late winter-early spring, while coho emerge in middle-late spring, and rainbow and cutthroat emerge in late spring to mid summer. Emerging fry can vary widely in size at emergence, ranging from 20+ mm nonanadromous cutthroat to 35-40 mm chinook. Upon emergence, fry actively feed on a variety of aquatic insects, and for those that freshwater rear for extended periods of time (particularly coho), the proportion of terrestrial food items in the diet may increase to over 30%³³⁸. Larger-sized juvenile salmon such as older aged rainbow and cutthroat prey on a mixed diet of aquatic and terrestrial macroinvertebrates, and may supplement their diet with occasional salmon eggs or fry^{498, 521}.

Sockeye fry are known to feed on cladocerans, copepods, and gammarid amphipods in lakes³³⁷.

After a summer of rearing in fresh water, juvenile coho average approximately 50 to 90 mm in length, and may weigh 2 to 5 g each^{128, 87}. Summer low-flow is a crucial time in the life of juvenile salmon having extended freshwater rearing. During this period the volume of aquatic habitat shrinks to a minimum, which can intensify inter- and intra-specific competition⁹. Declining streamflow conditions also may cause some fish (i.e., chinook, coho) to emigrate to estuaries^{196, 522}, where they continue to rear. Where species overlap in fresh water, a number of temporal and behavioral differences facilitate coexistence. For example, age-0 coho (juveniles that have not gone through a winter in fresh water) prefer slow deep habitats called pools, while steelhead age-0 prefer fast-water habitats called riffles, and therefore, are able to coexist in fresh water by partitioning the available food and space resources⁹⁶. On the infertile coast, chinook fry fill a temporal niche in spring prior to steelhead emergence, and thus coexist in a similar habitat until they often migrate to sea after 90 days of rearing in freshwater²⁸². Thus, the ratio of fast-water to slow-water habitats and their temporal utilization in a particular stream reach, during spring and summer, can influence the relative proportions of species and age classes of a salmon community.

Upon the first rains and high waters of fall, coastal species (juvenile coho, steelhead, and cutthroat) make a directed migration to seasonally alternate rearing habitats. Juvenile coho and cutthroat exhibit major immigrations into side-channel swamps⁸⁰ and riverine ponds^{398, 400, 94, 162}, located along river flood plains. Juvenile coho, steelhead, and cutthroat are known to immigrate into small “runoff” tributaries (valley-wall tributaries) of rivers^{94, 271}. Presumably these immigrations are to avoid high flows and turbidity of main rivers, as well as to take advantage of good feeding conditions during winter^{398, 399}. In contrast, interior (Idaho) juvenile chinook are known to move out of tributaries and into main rivers to over-winter, likely to avoid winter ice conditions in the tributaries⁵⁷.

After completing their freshwater stage, juvenile salmon of all anadromous forms undergo a physiological change called smoltification that includes osmoregulatory adjustments which prepare them to enter saltwater. For example, chum and pink salmon are nearly smolts upon emergence from the gravel, going directly to estuaries and the ocean^{259, 440}; while chinook¹⁹⁷ and coho⁴⁴² may either go directly to sea the first spring or summer of their life, or remain in freshwater for a whole year before smolting. Sockeye may rear in freshwater for one or two years before smolting⁷⁷, and steelhead²⁸³ and cutthroat^{160, 162} may not smolt for two or three years or more.

Once in the estuary or ocean, most salmon prefer to feed on such prey as amphipods, copepods, euphausiids, squid, herring, sandlance, rockfish, and anchovy^{83, 83a, 197}. While in the ocean, most salmon species migrate long distances to feeding grounds along the North Pacific coast^{178, 135, 136, 196, 194}. In contrast, some chinook stocks remain as “blackmouth feeders” in Puget Sound and the Strait of Georgia⁵⁷³, never going to the open ocean; and anadromous cutthroat may only range several kilometers from their natal stream without overwintering in the ocean³⁹⁵. Depending on the species, salmon use the ocean as a feeding ground in vastly different ways, some stay close to the North American Continent (i.e., chinook, coho, and cutthroat) and others (i.e., sockeye, chum, pink, and steelhead) forage far out into the north and western Pacific Ocean; but all survivors eventually return marine derived nutrients back to their rivers of origin at adulthood (Figure 4).

During their anadromous life history salmon make important ecological contributions (as prey) to various




Figure 4. Generalized ocean migration routes and biomass accrual of six species of Pacific salmon originating in Washington, Oregon, and British Columbia.

predators in the Pacific Northwest ecosystem, regardless of whether a particular individual salmon completes all life history stages or not (Figure 5). It is not uncommon for overall salmon survival rates to average 0.1% from egg to spawning adult.

Chum Salmon (*Oncorhynchus keta*)

Chum, or “dog” salmon spawn in the late summer into winter, depending on location. They deposit their eggs in a redd, and the incubation period can last from 3-5 months, depending on water temperature. There are mainly two races of chum salmon in Washington, the summer chum and the more abundant fall chum²⁵⁹. Some chum spawn intertidally, but most spawn in the lower reaches of rivers and streams. Chum salmon prefer medium sized gravel that is free of excessive amounts of sand²⁵⁹. The freshwater phase of juvenile chum salmon is virtually over upon fry emergence from the gravel,

Some Food Web Beneficiaries




Figure 5. Some food web beneficiaries of Pacific salmon nutrient in freshwater, estuary, and ocean environments.

at which time the fry migrate to protected marine waters and estuaries. Here they rear for several weeks before migrating to the ocean⁴⁴⁰. The length of time spent in the ocean can vary, but generally chum salmon grow to 3 or 4 years of age. Chum in Hood Canal can reach weights of 3.1 to 6.2 kilograms, depending on sex, stock, and year²⁵⁹ (Table 2).

Pink Salmon (*Oncorhynchus gorbuscha*)

Pink, or “humpy” salmon, like the chum salmon, use freshwater almost exclusively as an incubation environment, preferring to feed in estuaries and saltwater. Like the chum salmon, there are both summer and fall pink salmon, with falls predominating in run size. Pink salmon often spawn in the lower reaches of rivers and streams, and many are known to spawn in the intertidal areas¹⁹⁸. Spawning is usually complete by fall or early winter. Upon emergence from the gravel, pink salmon fry go directly to the ocean for rearing. Pink salmon are unique in that they have a strict 2 year life span, and in Washington the odd year cycle dominates⁵⁷³. Adult pink salmon average about 1.8 kilograms in weight⁵⁷³ (Table 2).

Sockeye Salmon (*Oncorhynchus nerka*)

Sockeye salmon, also called the “red” salmon, are unique because they generally use rivers that have an accessible lake environment in their drainage¹⁴⁶. Most sockeye adults enter freshwater in early to mid-summer, hold in a lake, and spawn in fall. Typically sockeye spawn in inlet or outlet tributaries of lakes, while some sockeye spawn in upwelling water along lake shorelines. Sockeye embryos have adapted to the reduced oxygen environments typical of the upwelling areas in lakeshore, stream, and spring areas⁷⁷. Reduced egg size, egg features that enhance oxygen transfer capabilities to the embryo, and a generally longer incubation period than other Pacific salmon, are some of those beneficial adaptations. The length of time spent in freshwater as juveniles varies from 1 to 2 years; while an additional 1 to 3 years is spent in the ocean rearing to adulthood. Most adult sockeye average between 1.5 to 3.6 kilograms in weight⁵⁷³. There is a nonanadromous form of sockeye called “kokanee”. The kokanee is much smaller in size at adulthood because it spends its entire life feeding in freshwater. Adult kokanee weigh about 0.5 to 1 kilogram. (Table 2).

Chinook Salmon (*Oncorhynchus tshawytscha*)

The chinook, or “king” salmon, is classified into three distinct varieties (spring, summer, fall) based on the season they enter freshwater as adults. They tend to spawn in large rivers and their tributaries, preferring deeper water and larger gravel substrate than the other species. Spawning usually occurs between August and November, depending on the particular variety. After fry emergence in winter and early spring, some “ocean-type” chinook swim downstream to the ocean within several weeks, while others “stream-type” spend up to a year feeding in freshwater before they migrate to sea¹⁹⁷. Migrant chinook exhibit major use of estuaries, particularly large marsh habitat (i.e., Skagit River, Fraser River) where they feed on amphipods. During their winter in freshwater, juvenile chinook have been found buried within gravel spaces, presumably to escape high stream flow conditions^{56,57}. The length of time spent at sea varies, but most chinook salmon return to spawn at age 3 to 5 years. Chinook have been known to reach weights of over 45 kilograms, however, most range between 5 and 10 kilograms⁵⁷³ (Table 2).

Coho Salmon (*Oncorhynchus kisutch*)

The ubiquitous coho salmon, also called the “silver” salmon, occur in almost every accessible coastal stream. Coho spawn in October through December, depending on the particular stock⁴⁴². Fry emerge from the gravel in April and May⁵¹⁰, and spend a summer feeding in pools or slow moving river side channels⁸⁰. In the fall, when stream flows increase, coho juveniles move downstream from their summer rearing habitats and immigrate into small flood plain tributaries and riverine ponds^{80,94,401}. Most smolting coho migrate to sea between the months of April and June, at age-1, but in cooler waters may remain a second year in freshwater. Coho spend about a year feeding in the ocean, then return to their natal stream to spawn at age-3⁴³⁹. Coho average between 3.6 to 5.4 kilograms as adults⁵⁷³ (Table 2).

Rainbow Trout (*Oncorhynchus mykiss*)

Rainbow trout are found in most coastal and interior rivers, favoring cold-fast flowing environments⁴⁹⁸. There are two main subgroups of rainbow, the anadromous and nonanadromous forms. The anadromous rainbow are called steelhead. There are two main varieties of steelhead, the so called “winter-runs” and “summer-runs”. Winter-run steelhead tend to predominate in coastal and Puget Sound rivers; while summer-run dominate in the interior Columbia and Snake River drainage. Nonanadromous forms of rainbow tend to exist in the headwaters of many rivers where there are steelhead.

During freshwater rearing, juvenile steelhead are often found in riffle environments, where they are able to minimize competition with juvenile coho⁹, but parr are also common in pools (Pat Slaney, personal communication). In the fall and early winter, juvenile steelhead redistribute and take up overwinter residence in small runoff tributaries^{94,560}, avoiding riverine ponds where juvenile coho reside⁹⁴. Juveniles are also known to bury themselves in the substrate during winter, presumably for protection against high flow conditions⁵⁶. Generally, juvenile steelhead spend two years in freshwater before smolting, however, some spend 1 year or 3 years. They spend an additional 1 to 4 years in the ocean, and return to spawn at age 2 to 5. Steelhead may reach weights of 13 kilograms, but most weight in the 3 to 6 kilogram class (Table 2).

Coastal Cutthroat Trout (*Oncorhynchus clarki clarki*)

Coastal cutthroat trout, also called “harvest trout”, are found in most Washington and Oregon coastal streams⁵²¹. Two main sub-groups of coastal cutthroat have been found living in tributaries of coastal rivers, the anadromous and nonanadromous forms. Anadromous (sea-run) cutthroat enter freshwater from the ocean in late summer and fall to either feed or spawn²³⁸. The nonanadromous form is characterized by both the non-migratory (fluvial) and within river migratory (potamodromous) types¹⁶². Spawning occurs in late winter through spring. After incubation fry emerge during late spring and early summer and take up residence in shallow riffles and pools in small headwater tributaries, usually upstream of all other species of salmon. This spatial segregation allows younger individuals to avoid direct competition with other salmon juveniles¹⁶⁶. In preparation for winter, juvenile cutthroat are known to *move downstream* and immigrate into flood plain tributaries to over-winter, similar to the movements of juvenile coho and steelhead^{94,162}. Anadromous cutthroat spend 2 to 5 years in fresh water before going to sea¹⁶⁰; however, they seldom over-winter in saltwater³⁹⁵. A large sea-run cutthroat trout would be 24 inches long (Table 2).

Table 2. Key sources of life history information for the 7 salmon species of Washington and Oregon.

Species	Reference
chum	Wydoski and Whitney ⁵⁷³ , Salo ⁴⁴⁰ , Koski ²⁵⁹ , Bjornn and Reiser ⁵⁶ , Everest et al. ¹⁴⁰ .
pink	Wydoski and Whitney ⁵⁷³ , Heard ¹⁹⁸ , Bjornn and Reiser ⁵⁶ , Everest et al. ¹⁴⁰ .
sockeye	Wydoski and Whitney ⁵⁷³ , Burgner ⁷⁷ , Foerster ¹⁴⁶ , Bjornn and Reiser ⁵⁶ , Everest et al. ¹⁴⁰ .
chinook	Wydoski and Whitney ⁵⁷³ , Healey ¹⁹⁷ , Bjornn and Reiser ⁵⁶ , Bjornn ⁵⁷ , Everest et al. ¹⁴⁰ .
coho	Wydoski and Whitney ⁵⁷³ , Sandercock ⁴⁴² , Tagart ⁵¹⁰ , Bustard and Narver ⁸⁰ , Salo and Bayliff ⁴³⁹ , Cederholm and Scarlett ⁹⁴ , Peterson and Reid ⁴⁰¹ , Bjornn and Reiser ⁵⁶ , Everest et al. ¹⁴⁰ .
steelhead	Wydoski and Whitney ⁵⁷³ , Stolz and Schnell ⁴⁹⁸ , Bjornn and Reiser ⁵⁶ , Allee ⁹ , Winter ⁵⁶⁰ , Cederholm and Scarlett ⁹⁴ , Everest et al. ¹⁴⁰ .
cutthroat	Wydoski and Whitney ⁵⁷³ , Glova ¹⁶⁶ , Trotter ⁵²¹ , Johnston ²³⁸ , Garrett ¹⁶² , Fuss ¹⁶⁰ , Pearcy ³⁹⁵ , Cederholm and Scarlett ⁹⁴ , Everest et al. ¹⁴⁰ , Hall et al. ¹⁸¹ .

FRESHWATER AND TERRESTRIAL HABITAT RELATIONSHIPS OF SALMON

Freshwater Habitat

Freshwater habitat of salmon includes all the physical, chemical and biological elements within the aquatic environment. Geology, climate, topography, disturbance history, nutrients from returning salmon, and characteristics of the riparian vegetation typically govern the characteristics and the distribution of habitat types in a watershed. Components of freshwater habitat include:

Physical Characteristics - channel width and depth, substrate composition, pool and riffle frequency, pool types, channel roughness.

Water Quality and Quantity - temperature, dissolved oxygen, dissolved nutrients, dissolved and particulate organic matter, hydrography.

Cover factors - interstitial spaces (space between gravels), undercut banks, woody debris, water surface disturbance.

Biological Factors - food availability, salmon carcass nutrient inputs, competition, predation, disease, parasites, and functioning riparian conditions.

Climate and regional geology determine habitat conditions at large spatial scales. The type of bedrock, the glacial history, and precipitation patterns contribute to landscape and channel morphology³⁸. In Washington, the South Fork of the Stillaguamish River (like many Puget Sound rivers) was heavily glaciated, creating a landscape of steep, highly dissected hill slopes in the headwaters, and terraces formed by glacial outwash in the valley bottoms³⁸. The land form dictates the channel gradient, from gentle hill slope morphology on terraces, to steep and precipitous conditions in the headwaters, and plays a large role in shaping the salmon habitat characteristics in the channels³³. Streams draining outwash terraces, which have little topographic relief, typically exhibit gradients less than 4% and contain abundant pool habitat. Stream channels on these terraces support many of the anadromous fish populations in the watershed³³.



Plate #10. Low gradient stream showing large woody debris formed habitat in Monroe Creek, Washington. (Photo by: Jeff Cederholm).

This change of channel conditions from the headwaters downstream is typical of many watersheds in the Pacific Northwest and correlates well with shifts in the fish communities . Montgomery and Buffington ³³² described various channel forms and have developed a classification system based on channel size, gradient, and the presence of roughness elements. The geomorphic classification of stream channels allows one to better understand the distribution of the various salmon species within a watershed. For example, the anadromous forms are usually located in the downstream low gradient reaches, where they have unimpeded upstream and downstream passage; while the resident nonanadromous forms are located in the middle to

upper steep headwater areas. The distribution of various species of salmon in a drainage has been discussed by Reeves et al. ⁴¹⁷.

The abundance of fish in a stream is greatly affected by the stream's capacity to produce food. Many of



Plate #11. Hoh River on the western Olympic Peninsula. (Photo by: Jeff Cederholm).

the factors influencing stream productivity change predictably with changes in stream size, a pattern termed the *river continuum* ⁵²⁹. Productivity is influenced by nutrient availability, input of organic matter from external sources, and the capacity for the channel to store and process organic matter, and light.

Differences in these factors can be very large and lead to a high degree of variability in production of fish, including salmon and trout populations

²²³. Some of the highest freshwater production values have been reported for trout in New Zealand spring streams, $54.7 \text{ g/m}^2/\text{y}$ ¹⁰; however, production values from the Pacific Northwest are generally low when compared with other regions of the world, often below $1.0 \text{ g/m}^2/\text{y}$ and very rarely over $5.0 \text{ g/m}^2/\text{y}$ ⁵⁵.

Riparian Habitat

Many of the functional and structural attributes of stream habitat are created and maintained through interaction with riparian vegetation. Riparian areas constitute the interface between aquatic and terrestrial ecosystems^{506, 171}, performing a number of vital functions that affect the quality of salmon habitats as well as providing habitat for a large variety of terrestrial plants and animals. Riparian areas influence streams and consequently salmon habitat in a variety of ways⁴⁸⁵, including:

Shade - which dampens seasonal and diel fluctuations in stream temperature and controls primary and secondary production.

Streambank stabilization - provides erosion resistant roots that bind soil particles together, thus facilitating bank building during high flow events by slowing the stream velocities.

Sediment control - regulates sediment flow from upland areas by acting as a filter, or storing sediments in the primary flood plain.

Litter Input - contributes a significant amount of organic matter to streams, which acts as an important food resource for aquatic communities.

Large woody debris (LWD) - provides important structure to the stream channel for energy dissipation, fish habitat, and salmon carcass retention.

Nutrient input - riparian zones mediate the flow of nutrients to the stream and are, therefore, important regulators of stream production. Some riparian species such as red alder (*Alnus rubra*) also fix atmospheric nitrogen therefore augmenting N availability to the ecosystem.

Microclimate - streamside soils and vegetation can have a significant effect on moderating the climate within riparian zones.

Streamside vegetation moderates water temperature, and this relationship is influenced by elevation, air temperature, stream width, water depth, and aspect⁴¹. Removal of riparian vegetation has been associated with increased maximum water temperatures, and diurnal fluctuations in water temperature during summer; and decreased winter water temperatures⁴¹. Small, low-elevation streams are the most susceptible to summer water temperature increases caused by canopy removal⁵⁰⁴. The biological consequences of elevated water temperature on aquatic communities are complex. There is little information indicating direct mortality of fishes as a result of temperature changes related to riparian canopy removal⁴¹; however, reductions in growth rate^{51, 572}, changes in life history²¹⁵, changes in competitive interactions between species⁴¹⁸, reductions in fecundity of adults⁴⁰, and an increased susceptibility to disease^{386, 363}, have all been documented. Some species of amphibians and aquatic macroinvertebrates also are thermally intolerant and elevated water temperatures may have detrimental impacts on their populations^{327, 137, 78}.

Riparian vegetation increases streambank stability and resistance to erosion. Roots from woody and herbaceous vegetation bind soil particles together, helping to maintain bank integrity during erosive high-streamflow events^{506, 485}. Riparian vegetation also facilitates bank-building during high flow events by slowing stream velocities, which in turn helps to filter sediments and debris from suspension. This combining action helps to stabilize and rebuild streambanks, allowing the existing channel to narrow and deepen, and increases the effectiveness of riparian vegetation in providing bank stability and shade¹³². During over-bank flows, water is slowed and fine silts are deposited in the flood plain, increasing future productivity of the riparian zone⁴⁸⁵.

Forested riparian areas generate much of the organic matter that provides the energy source for the trophic systems of small streams. In one study of forested headwater channels in Oregon, Sedell et al.⁴⁵² determined that over 90% of the in-channel organic matter was provided from the surrounding terrestrial

environment. A 10-m wide stream in western Washington received over 75% of its annual organic matter supply from terrestrial sources⁴⁶. Even though this source of organic matter decreases relative to autochthonous organic matter in larger channels, it remains vital to stream productivity.

Large woody debris has been shown to be a critical structural component in Pacific Northwest streams, forming pools, waterfalls, and overhead cover; and it also regulates the transport of sediment, gravel and organic matter, for fish and other aquatic biota^{50,49}. In forested watersheds LWD provides the most common obstruction, often forming pools in various types of fluvial channels³³². Without this material, pool abundance and size is decreased⁴⁵, reducing habitat complexity and potentially reducing the diversity of the fish community. In addition to numerous habitat and morphological functions⁵⁰, wood (organic debris) helps retain salmon carcasses in streams for biological activity⁹⁰. The capacity of many small streams to retain carcasses has probably been reduced by human activities, and this could have serious impacts on the food chain of fishes, and on the available food supply of many carnivorous wildlife species⁹¹.

Riparian areas play a key role in determining the concentration of nutrients in stream water⁴². The presence of even a narrow riparian buffer can profoundly influence stream water chemistry. Uptake and storage of various elements carried by groundwater can be considerable, even where input rates have been substantially altered as a result of upslope land uses²⁸⁴. Riparian vegetation composition can influence nitrogen input to streams. Early successional vegetation in riparian areas in the Pacific Northwest is often dominated by red alder, a nitrogen (N) fixing species. As a result, the N content of litter beneath these riparian stands is 1.5 to 3-fold higher than sites where conifer species are the dominant component of the overstory^{128a}. The result is higher N levels in the riparian soils⁵⁸ and increased delivery of N to the stream channel. Higher N levels in stream water may elevate primary production and decomposition (heterotrophy) in the channel and increase food availability for the invertebrate community. Increased invertebrate production may elevate food availability for stream-dwelling fishes, amphibians, and other insect feeders such as bats and flycatchers.

The riparian area may act as either a source or sink of organic matter and sediment during flood flows. The manner in which the stream and riparian area interact at these times depends upon the morphology and vegetation of the riparian zone and the intensity of the discharge event³⁴. The structure and abundance of riparian vegetation plays a key role in moderating the movement of materials between the riparian area and the stream³⁶¹. Vegetation in the riparian zone has been shown to be the single most important structural element for the retention of fluvially transported organic matter during high flow events⁴⁸⁴. Similarly, riparian vegetation promotes the storage of sediment²¹⁶, that may provide germination sites for some species of riparian plants³⁷⁰. The variations in retentive capacity of different riparian areas for organic matter leads to large differences in the organic content of riparian soils, ranging from nearly all inorganic material in some locations to very high concentrations of organic matter in stream-adjacent swamps and wetlands⁷¹. This variation in substrate further contributes to riparian vegetation heterogeneity².

The area, in which water exchange between the channel and the underlying riparian soils occurs is termed the *hyporheic zone*⁴⁹¹. The extent of the hyporheic zone varies as a function of site topography and soil characteristics. In riparian areas of low relief and porous soils, the hyporheic zone may extend as far as 3 km from the edge of the channel⁴⁹¹. The riparian vegetation has an influence on the amount of water stored

in the hyporheic zones. Hicks et al.²⁰⁷ found that August stream flows following logging of a small western Cascade Mountain watershed, including the removal of riparian vegetation, increased for 8 years as a result

of reduced transpiration. Subsequently, as early successional vegetation occupied the riparian area, summer stream flows decreased to below pre-harvest levels due to high transpiration associated with this vegetation type. Riparian vegetation, through its influence on riparian soil characteristics and water movement, also can impact chemical transformations within the hyporheic zone⁵²⁰. Along low-gradient, unconstrained stream reaches, vegetation-hyporheic interactions may occur over a broad area. In areas of moderate to high topographic relief the zone of direct interaction between vegetation and the hyporheic zone will be reduced.

The attractiveness of riparian areas to wildlife likely reflects three main attributes: the presence of water, local microclimate condition, and more diverse plant assemblages found in riparian areas compared to uplands. Wildlife also congregate seasonally in riparian areas where salmon spawn, to take advantage of an abundant food supply of carcass flesh⁹¹. The high value of riparian habitats to wildlife has been recognized by naturalists³⁸⁰, and considered a bridge between upland habitats and the aquatic environment. The combination of shape, moisture, deposition soils, and disturbance regime unique to riparian areas contributes to their exceptional productivity in terms of plant growth, plant diversity, and structural complexity of the vegetation^{237, 329, 270}. Wildlife dependency and diversity peak at this terrestrial/aquatic boundary. Brown⁶⁹ reports that 359 of 414 (87%) species of wildlife in western Washington and western Oregon use riparian areas and wetlands during some season or part of their life cycle. In their detailed examination of wildlife and habitats for all of Washington and Oregon, Johnson and O'Neil²³⁶ reported that 393 of 456 (86%) of the common terrestrial and freshwater wildlife species have seasonal use of riparian areas, wetlands, and streams. Of these 393 species, 110 were found to be *closely associated* (e.g., obligates) with eastside and westside riparian habitat types.

The close association may very well have evolved from the direct or indirect exploitation of the rich vegetative habitat provided by riparian areas²⁵⁵. Quantitative studies conducted during the past several decades have supported observations and have identified biological and physical attributes of riparian habitats which enhance their value to wildlife. Brinson et al.⁶⁴ and Oakley et al.^{381 cited in 380} summarize these important biological and physical features of riparian areas:

- C presence of surface water.
- C increased humidity, high rates of transpiration, and greater air movement.
- C complexity of biological and physical habitats.
- C maximum edge effects with adjacent upland forests which is beneficial for some species.
- C food supply.
- C thermal cover.

According to O'Connell et al.³⁸⁰ stream type has a direct influence on the riparian habitat and its associated wildlife communities. In the smaller headwater streams the impacts of the upstream riparian vegetation on the streams is greater than downstream where flow volume increases, flooding is more widespread, and the impact of riparian vegetation on the stream is less. Brinson et al.⁶⁴ suggest that middle order perennial streams and associated riparian areas have the greatest wildlife use. Periodic flooding can enhance the availability of food for wildlife by creating new feeding areas⁶⁴. Flooding can also make riparian habitat unsuitable for other species. Species abundance of riparian mammal communities has been related to the timing of recent hydrologic events; impoverished mammal populations have been attributed to recent flooding whereas more abundant populations have been observed in areas not subject to recent flooding⁶⁴.

Habitat Forming Processes

Disturbance plays a major role in maintaining community diversity and productivity in many ecosystems^{105, 420}, and is a key factor in creating and maintaining diverse stream habitat in the Pacific Northwest^{38a, 54}. These disturbances range in severity from minor events, such as seasonal changes in flow, to less frequent high intensity events such as wildfire, debris torrents, and major floods. Riparian and channel conditions evolve as impacted areas recover from disturbance²¹¹. The result is a diverse set of riparian community types and stream habitat conditions that vary over both time and space².

Disturbance contributes to both diversity of aquatic fauna and productivity of these communities when considered at a watershed level¹⁷³. Aquatic communities associated with early-successional riparian areas typically exhibit low diversity, but high productivity for certain species. Removal of the channel shading canopy brings about dramatic increases in light and algal productivity; however, input of terrestrial litter decreases^{172, 46}. Invertebrates that feed on algal material (grazers) typically dominate communities at recently disturbed sites,¹³⁷. These invertebrates form a major component of the diet of some salmon and trout³³⁸ and can contribute to increased fish productivity following disturbance^{352, 52, 46}. The increased productivity is typically observed during summer, and often does not extend into winter months when the availability of shelter from high flows for juvenile salmon becomes important^{94, 401, 162}.

After forest canopy closure, primary productivity in streams decreases. The type of litter delivered to these systems and the physical characteristics of the channel differ from those at sites bordered by mature vegetation. Hardwood trees, especially red alder, often dominate the canopy at these sites. Litter from red alder trees decomposes much more rapidly than conifer litter, in part due to the higher N content⁴⁵¹. The high N content of the litter improves its nutritional value for shredding macroinvertebrates but the high rate of decomposition causes it to be scarce at some times of the year.

Alder stands begin to die-out and provide LWD to channels after about 60 years¹⁷⁵. Shade tolerant conifers, like western red cedar (*Thuja plicata*) and western hemlock (*Tsuga heterophylla*) colonize the site and begin to provide needles and other litter to the channel. Some stands of alder can persist and will repeat themselves several times (Slaney, personal communication). Woody debris amounts and average piece size increase for 100 or more years following conifer occupation of a site^{506, 44}. The morphology of the channel and the routing of sediment and organic matter evolve slowly as the riparian community changes, ultimately creating channels which are highly complex structurally and support a macroinvertebrate community dominated by shredders¹¹.

Forest practices and other land uses have accelerated the rate of occurrence of some types of disturbance. The acceleration in disturbance has led to the establishment of early successional communities in the majority of riparian areas on commercial forest land in the Pacific Northwest^{60, 86, 154}. Practices, such as splash damming of rivers to float logs to market⁵⁵⁰, and removing all trees to the channel's edge⁴³ modify the riparian successional process. Timber harvest or roads constructed on unstable slopes or road drainage systems that were improperly maintained, dramatically increase the incidence of landslides^{421, 93, 456}. Many hillslope failures enter stream channels and may move considerable distances downstream, removing streamline vegetation and soil. On the positive side, however, localized landslides also can input massive amounts of spawnable sized gravels and LWD into stream channels, where they may benefit salmon populations⁴⁵⁶. These disturbance events have affected a large proportion of the riparian areas bordering streams in the region over the last century, and have played a key role in determining channel form and habitat conditions⁵⁰⁷.

From a study of fire history in Mt. Rainier National Park, Hemstrom and Franklin ²⁰¹ reported that alluvial terraces and valley bottoms were often forested with old stands and that every major river valley contained a stream-side old-growth corridor. These observations support an inference that the moist environment of riparian areas inhibits fire and reduces the fire return interval for riparian forests. The moister environment may also advance the regeneration of more structurally complex forest following wildfire. Fires that burn across riparian areas may be less intense, and less intense fires would kill fewer trees and consume less coarse woody debris, both snags and logs. The persistence of live trees in riparian forests may also provide a local seed source that facilitates a more rapid development of a multi-layered, conifer-dominated forest ⁴⁰⁵.

Beaver

Beavers have long co-existed with salmon in the Pacific Northwest, and have had a important ecological relationship with salmon populations. The beaver created and maintained a series of beneficial aquatic conditions in many headwater streams, wetland, and riparian systems, which serves as juvenile salmon rearing habitat. Beavers have multiple effects on water bodies and riparian ecosystems that include altering hydrology, channel morphology, biochemical pathways, and stream productivity³⁸⁵. Beaver ponds were of special importance in more arid regions, but also had important roles in coastal systems ³⁶⁵. Beavers were once extremely abundant in the Pacific Northwest, but as far back as 1778, trapping expeditions into western North America began depleting their numbers. Between 1834 to 1837, pelts from 405,472 beavers from the area that would become southwest Washington and Oregon were shipped to Europe. It is difficult to imagine the amount of influence beavers have had on the landscapes, most Pacific Northwest streams have been void of beaver activity for many decades before ecologists had the opportunity to study them.

Past excessive trapping, and subsequent unregulated land- and water-use activities, significantly reduced abundance of beaver and beaver ponds. Even ponds of the surviving beavers were actively removed. Additionally, excessive livestock grazing in riparian areas has degraded habitat conditions for beaver ³⁸⁵. Severe declines of beaver in Washington and Oregon have fundamentally altered important natural aquatic ecosystem processes such as nutrient cycling, flood plain development, and stream hydrology.

Beaver dams can obstruct channels and redirect channel flow and the flooding of streambanks and side channels. By ponding water, beaver dams create enhanced rearing and over-wintering habitat that protect juvenile salmon during high flow conditions ³⁶⁵. Studies in Oregon coastal streams have suggested that where the amount of spawning is adequate, the winter survival of juvenile coho, which can be swept downstream in high winter flows, is limited by the presence of adequate slow-water habitat ³⁷⁴. Beaver dams are often found associated with riverine ponds called “wall-base channels”⁴⁰¹ along main river flood plains, and these habitats are used heavily by juvenile coho salmon ^{400, 94} and cutthroat trout ^{94, 162} during the winter. Though their dams can occasionally block upstream migration of adult and juvenile salmon, studies of trout movement indicate that fish can pass over beaver dams during all seasons ³⁸⁵. Beaver dams may temporarily keep salmon adults in the lower parts of spawning streams where flows are greater and pools are deeper, then, when dam breaching flows occur, free passage to upstream areas is provided.

Beaver foraging can cause a loss of woody riparian vegetation and an increase of fine sediments, but it also increases the input of large woody debris to streams and beaver droppings may enrich pond productivity. Bank dens and channels can increase erosion potential, but because ponds fill with sediment to become

wetlands over time, this helps to retard upstream erosion and retain sediments that otherwise could adversely alter downstream areas ⁴⁸⁵. In a Wyoming study of an area that had 10.5 beaver dams per km, each dam was found to retain 5,350 m³ of sediment. In another Wyoming study, sediment loads were reduced by 90% after flowing 8 km through an area with well developed riparian habitat and beaver dams.

Beaver ponds provide a sink for nutrients from tributary streams and create conditions that promote anaerobic decomposition and de-nitrification. These processes can cause nutrient enrichment and increased primary and secondary production downstream from the pond and increasing nutrient retention time and enhanced invertebrate production in the pond ³⁶⁵. These factors help increase salmon growth and survival, and also helps improve water quality. Beaver ponds increase the surface to volume ratio of the impounded area, which can result in increased summer temperatures ⁴⁸⁵. Beaver ponds also can cause increased storage of water in the banks and flood plains, and this increases the water table, enhances summer flows, adds cold water during summer, and causes more even stream flows throughout the year. During winter, beaver ponds in cold environments prevent anchor ice from forming and prevent super-cooling of the water. By storing spring and summer storm run-off, beaver ponds help to reduce downstream flooding and the damage from rapid increases in stream flows ³⁸⁵.

Beavers also help shape riparian habitat. Beaver ponds increase the surface area of water several hundred times and thereby enhance the overall riparian habitat development ³⁸⁵. They also enhance vegetation growth by increasing the amount of groundwater for use by riparian plants and wetland areas. The presence of beaver can have both positive and negative influence on salmon habitat, but on the whole, their presence is considered of great benefit to both water quality and salmon, particularly juvenile coho salmon and cutthroat trout, and to many other species of wildlife and invertebrates.

ESTUARY HABITAT

By definition ⁴¹³, an estuary is a region where salt water of the ocean is measurably diluted by freshwater runoff from the land within a constricted body of water. Thus, the salinity gradient that juvenile salmon encounter when migrating through estuaries depends upon the inflow of fresh water and the strength of the tides, which influences the degree of mixing, and the depth to which the juvenile salmon penetrate the water column. Because the saline seawater is more dense than fresh water from the river, the fresh water tends to lay over the seawater unless it is thoroughly mixed by tidal, river and wind energies. In the Pacific Northwest, river flow plays a strong role in the structure and dynamics of estuarine circulation ⁵¹⁵. Variation in circulation reflects the general seasonal cycles of the Pacific Northwest climate, but also the geomorphic structure of both watershed and estuary. Estuaries of lowland watersheds along the Washington and Oregon coasts tend to exhibit high peak flows associated with winter storms, but often extremely low flows associated with the dry summers. This can often cause dramatic differences in the available estuarine habitat between winter and spring-summer periods, limiting summer rearing. In some southern Oregon and northern California estuaries river, flow can decrease to the point that bars form across the estuaries' entrances, restricting juvenile salmon ocean emigration to extreme high (spring) tides.

Perhaps the most fundamental concept in understanding the estuarine ecology of juvenile salmon is that the salmon do not respond to singular habitats *per se*, but rather interact with a landscape mosaic of habitats in response to changing migratory mandates, tidal cycles and freshwater runoff events (Figure 6). River flow and tide, physiological change, prey and predator distributions, and likely metapopulation genetic structures as well, all affect the rate of movement through the estuary. But, the opportunity for juvenile salmon to exploit preferred habitats is just as likely dependent on the arrangement of key landscape features such as tidal-freshwater and brackish rearing zones, low-velocity refugia, migratory corridors, and foraging patches. Although this is a relatively new topic of research, with few definitive experiments and tests, there is some emerging evidence that the edge of marsh vegetation in dendritic tidal channel and slough systems may relate directly to juvenile salmon production ⁴⁷³.

Large watersheds with significant snow accumulations at higher elevations, and extended melting periods can create prolonged spring freshets ³⁰⁸. Spring and winter freshets, and winter "rain-on-snow" events associated with rapid snowmelt, produce flooding in tidal floodplains and estuaries that influences short-term and long-term productivity of juvenile salmon and their ecosystems ^{360, 566}. Although flood plain and estuarine wetland flooding increases flows in the main distributary channels, likely diminishing the ability of juvenile salmon to occupy them, considerable side-channel and other flood plain wetlands (i.e., ponds, relict side-channels) are inundated and become available for refuge and rearing. This flooding recruits organic detritus and dissolved nutrients from these peripheral wetlands and imports them to the estuary. While trapped in estuarine wetlands or circulation features such as estuarine turbidity maxima ⁴⁷², these materials contribute to primary and secondary production by supporting food web pathways to juvenile salmon.

The structure of the watershed and estuary, and the seasonal variability in river flow, shapes estuarine circulation, and strongly influences juvenile salmon residence time, habitat use and production. Except where the river has been extensively diked and channeled, the flood plain in the freshwater-tidal region is characterized by extreme habitat complexity, abrupt changes in water velocity and low-velocity off-channel habitats. As the "estuarine gateway," the tidal-freshwater mixing zone can be exceedingly important to juvenile salmon ⁴⁶⁹ because it:

(1) provides habitat for overwintering chinook, coho and steelhead forced downstream during high river flows; (2) contains complex low-velocity refugia such as off-channel sloughs and LWD; (3) allows migrating juveniles to adapt physiologically as they encounter brackish waters of the upper estuary; (4) drift insects are trapped and concentrated due to flow reversals, providing opportune feeding conditions⁵²³; and (5) is the first region of estuarine settling of suspended sediments and detritus, which can fuel soft-sediment habitat formation and detritus-based food webs exploited by salmon. River flow and tide, physiological change, prey and predator distributions, and likely metapopulation genetic structure as well, all affect the rate of movement through the estuary.

Estuaries are composed of both discrete and highly integrated habitat complexes and their associated plant and animal communities. Categorizing habitats to a large degree is a function of scale, as juvenile salmon can respond to habitat features (e.g., LWD or tidal channels) that are elemental to the broader habitats. Estuaries generally possess eight habitat components: (1) subtidal distributaries; (2) mud- and sand-flats; (3) gravel-cobble beaches; (4) low elevation emergent marshes; (5) high elevation emergent marshes; (6)




Figure 6. Movements and migrations of juvenile Pacific salmon across tidal-freshwater delta-estuarine landscapes (Contributed by Simenstad, unpublished diagram).

forested and shrub swamps; (7) eelgrass; and (8) kelp. Salmon communities have been shown to utilize many of these habitat types. Juvenile coho (fry, fingerling) are often found rearing during winter and early spring in the tidal flood plains of many large rivers such as the Chehalis River^{323, 472, 471, 470}. These fish are either staging for migration through the estuary or are moving back into freshwater for extended rearing. Work in British Columbia^{522, 523, 438} and Alaska⁵¹² show that certain sub-populations have a minimal juvenile

freshwater rearing phase of their life history (“ocean-type”), and spend extended periods of time either feeding in estuaries or in the ocean. Such subyearling migrant coho, may constitute significant portions (up to 50%) of the returning adult spawners⁵²³.

Natural disturbance regimes are responsible for creating and maintaining habitat complexes important to juvenile salmon. Erosive flooding, channel reconfiguration, and changes imposed by LWD all promote increased habitat complexity and heterogeneity. In the absence of disturbance, early successional habitats such as mudflats and low elevation estuarine marshes (e.g., *Carex lyngbyei* sedge) would not persist or would be relatively rare. Yet, these habitats can be some of the most productive and beneficial of salmon habitats and play unique roles for some salmon species.

Association with specific migratory and rearing habitats in estuaries tends to relate primarily to fish size, but there may be some indications of co-evolutionary habitat partitioning^{196,466}. Juvenile salmon measuring 30-60 mm long tend to occupy near shore shallow water (1-2 m deep), often irrespective of habitat type and tidal stage. The result is that juvenile salmon likely use shallow water habitats as refugia during both migration and rearing. Chum and chinook fry are particularly noted to occupy estuarine marshes and adjoining habitats for extended periods of time, up to 1 month^{273,104,460}. However, some species diverge from this habitat use: pink fry spend relatively little time in estuarine marshes and start making the transition to more offshore neritic (surface) waters after only a few days to a week in the estuary. In contrast, chum fry do not begin to disperse into neritic habitat until they have grown to 50-60 mm long, which usually involves more than two weeks in the estuary. Based on evidence of extended residence times in estuary habitats, subyearling chinook fry appear to maintain some affinity for shallow-water habitats until they are even larger (i.e., 80-100 mm long)^{104,196,197}. Information on residence time of coho fry in estuaries is more limited, but residence times for these “ocean-type” fish appear to fall between chum and chinook.

Chum and pink salmon that migrate directly into brackish and saline estuarine waters appear to require little or no interim adaptation²⁵⁶; when smoltification does occur, it occurs very rapidly upon entering brackish waters, and adaptability may actually decrease with freshwater rearing^{227,192}. Yearling chinook, coho and sockeye salmon proceed through a definitive smolt stage before entering saline waters, while subyearling chinook and coho appear to spend considerable time in the tidal freshwater-brackish zone of estuaries, perhaps in part due to an extended smoltification process.

Feeding behavior and diet of juvenile salmon passing through and rearing in estuaries is often specialized on specific types, species, and even life history stages of organisms, suggesting strong co-adaptive development of preference for bioenergetically “optimum” prey resources that through rapid growth provides a survival margin upon entry to the ocean²⁴⁷. Diet is strongly structured by size of fish and habitat occupied, and to some degree may be influenced by earlier life history stages⁴⁶⁶. The density of prey taxa may actually influence estuarine migration rates and residence times^{467,565}.

OCEAN HABITAT

Upwelling along the coast of the Pacific Northwest often results in high primary and secondary productivity, resulting in large standing stocks of fishes, seabirds, and marine mammals. The coastal upwelling domain extends from British Columbia to Baja California and is located inshore of the equatorial flowing California Current. Coastal upwelling is driven by prevailing northwesterly winds during the spring and summer. Those winds result in offshore displacement of near-shore surface waters and vertical advection of deep, cool and often nutrient-rich waters into the euphotic zone along the coast and into estuaries. Rich blooms of phytoplankton are observed along the coast following episodic upwelling events^{514, 286, 501}. Upwelling varies seasonally and over longer annual and semiannual cycles, with intensity generally increasing southward to northern California. Upwelling is most intense in regions of capes such as Cape Blanco in southern Oregon.

There was a strong correlation between the intensity of coastal upwelling and the smolt-to-adult survival of hatchery coho salmon from the Oregon Production Index region south of the Columbia River from 1960 to 1981³⁷³. During the late 1970s, however, there was a major change in ocean climate in the North Pacific Ocean, called a regime shift or the Pacific Decadal Oscillation, which was correlated with a sharp decline in the survival of Oregon coho salmon between smolt release years 1975 and 1976^{397, 32, 153}. After this regime shift the production of Oregon coho salmon has usually been low. The relationship between coastal upwelling and coho survival is no longer significant. The reason for this changed relationship is unclear, but is related to weak coastal upwelling, warm sea temperatures, high sea levels, and frequent El Niño events³⁹⁴. Upwelling has probably not been effective in injecting nutrient-laden water into the euphotic zone because of the deep lens of overlying warm, nutrient-depleted water along the coast^{195, 432}. The persistence of warm, unproductive ocean conditions is a major reason for the decline of many stocks of anadromous fishes along the west coast, and for the very large variability in survival and reproduction of marine birds.

Although the mechanisms that have resulted in poor ocean survival of salmon are speculative, one hypothesis is that weak upwelling results in low growth and poor survival of zooplankton and forage organisms, and impacts juvenile salmon during their critical first summer in the ocean. This lack of forage and the narrow band of cool waters along the coast during weak upwelling years concentrates juvenile salmon near the coast where they are more vulnerable to predation by seabirds, marine mammals and fishes¹⁴⁵. During warm years predators from southern waters, e.g., Pacific and jack mackerel, invade coastal waters and may either compete with or prey upon juvenile salmon^{397, 394}.

The principal prey of juvenile salmon off the coast of Oregon and Washington during the spring and summer are fishes and crustaceans^{83, 83a, 395}. Salmon in the open ocean forage opportunistically on a diverse assemblage of pelagic organisms. The diets of maturing salmon in the North Pacific Ocean vary among species and sizes of fish, with season and year, and with location and proximity to the coast. Fishes, squids, amphipods, copepods, and pteropods are primary prey^{268, 396, 178}.

ECOLOGICAL RELATIONSHIPS OF SALMON

Macroinvertebrates

Freshwater Macroinvertebrates And Salmon

Freshwater ecosystems are inhabited by a large variety of macroinvertebrates that play an integral part in the salmon's life history. They include insects, crustaceans, and other forms of macroinvertebrates (larger than 595 microns in their later instars or mature forms). Many species in their aquatic phase have been described from the hyporheic zone, or zone below the surface of the stream bottom⁴⁹². Given the variety of physical habitat across the region's landscape, there is an opportunity for freshwater invertebrate species to form diverse and specialized communities.

Freshwater macroinvertebrates play a significant role in energy pathways of aquatic ecosystems. The consumption of algae, detritus, and bacteria is the basis for transfer of this energy. A few invertebrate species are known to actively derive their food base from higher life forms (e.g., small fish). The food source used by an invertebrate defines what function it performs in this food web.

Structure And Function In Macroinvertebrate Communities

The type and location of food in the aquatic environment consumed by invertebrates determines their functional designation. Headwater streams or heavily canopied streams are dominated by leaf litter input, allochthonous material, which has been linked to significant shredder activity^{111,536}. Shredders comprise a group of aquatic insects that utilize coarse particulate organic matter, such as leaf litter, with a significant dependence on the associated microbial biomass⁵²⁹. Portions of a drainage where the riparian canopy opens can result in substantial autochthonous input (periphyton growth), and are consumed by scrapers like the mayfly family Heptageniidae³¹⁵. Lower in a drainage the channel can accumulate large deposits of detritus. Invertebrates distributed here are mainly collector-gathers and may constitute the bulk of juvenile salmon diets¹⁹⁶. The distribution of dominant food sources throughout a drainage are influenced by a continuum of physical changes as one travels from the steep headwater streams to the relatively low gradient flood plains⁵²⁹.

Invertebrate community structure in a stream or pond reflects physical characteristics of the living space. Numbers of species in a stream ecosystem are usually greater in physically diverse habitats. Structural attributes like species richness change along a disturbance gradient. Two investigations found that species richness was consistently higher in streams with intermediate disturbance of substrate^{116,519}. The effect of disturbance and physical change over a continuum results in species replacement and sometimes adjustments of the functional characteristics in the community³²⁵.

Physical And Chemical Influences On Macroinvertebrate Distribution

Factors that control distribution and abundance of macroinvertebrates are substrate, current velocity, temperature, predators, and food resources²²³. Substrate heterogeneity often promotes greater species richness^{326,328}. Interstitial spaces in stream gravels can serve as refuge from predators and physical disturbance, and entrap detritus. Water temperature in the interstitial microenvironment can be relatively constant and cooler than the overlying surface water⁵⁵⁷.

Early life stages of the salmon can be affected by substrate quality. Factors that favor survival of salmon egg and fry (low levels of fine sands and silts) are coincident with requirements of aquatic invertebrates that have

a narrow tolerance range to environmental fluctuations. Protection from natural physical disturbance is important for early life stages of salmon and mobile aquatic invertebrates. Stable stream bottoms during periods of flood or freshet reduce predation on dislodged animals. In some instances, salmon redd construction is a natural disturbance that reduces invertebrate density in localized areas of a stream³²⁴. This disturbance also opened niche space for other functional groups of aquatic insects, like blackflies, who feed on suspended particles and recolonized quickly along with stonefly nymphs and midge larvae³²⁴. Other invertebrates that enter the drift behaviorally or unintentionally from substrate disturbance are potential prey items for feeding salmon. Mayfly and stonefly density and richness can be reduced by physical alterations to the stream corridor. These changes may have significant implications to the salmon food base. Invertebrate drift is either voluntary, a behavioral activity, or coincides with catastrophic stream conditions, especially during floods. Taxonomic groups prominent in behavioral drift are amphipods, Ephemeroptera (mayflies), Plecoptera (stoneflies), Tricoptera (caddisflies), and Simuliidae (blackfly larvae). Later stages of the nymph and larval forms are most active in the diel (24-hour cycle) drift⁵⁴⁷. Behavioral drift occurs with a diel periodicity, typically at two peaks in a 24-hour time frame. Most invertebrates that enter the drift are night-active, with photoperiods as the major cue. Fewer invertebrates are day-active and begin drifting by cues through change in water temperature.

Drifting invertebrates are a food source for certain species of fish that forage in the stream water column. Rader⁴¹⁵ determined that the mayfly genus, *Baetis*, whose drift propensity was high, was a significant food source to juvenile and adult salmon. Other studies indicated that food preference of juvenile fish was related to their abundance and location within the stream channel. Juvenile coho salmon diet varied seasonally depending on the type and abundance of invertebrates, salmon fry, or salmon eggs in the benthos or drift²⁶⁰.

Significance Of Macroinvertebrate Life Cycles

There are two life strategies characteristic of freshwater macroinvertebrate species⁵⁵⁸. The simpler *hemimetabolous* strategy inherent in stonefly and mayfly species contain an egg, multiple nymph, and adult stages. A few of the stonefly species are long-lived (more than a year) in the aquatic nymphal form. Large-bodied stoneflies found in streams indicate adequate flow in channels that are key to survival of early salmon life stages and to some of the invertebrate fauna they will eventually consume.

The second life strategy contains representatives of the *holometabolous* invertebrates. Midges, blackflies, and caddisflies have egg, larva, pupa, and adult life stages. These types are mostly short-lived having one or many generations per year in a population. Aquatic environments that are seasonally stressed by high temperatures, low dissolved oxygen, or drought are primarily colonized by holometabolous invertebrates. These stressors increase the mortality of early life stages in salmon, but encourage dominance of holometabolous species in the aquatic invertebrate community.

Aquatic Macroinvertebrates As A Food Source For Salmon

Aquatic ecosystems are frequently inhabited by both hemimetabolous and holometabolous macroinvertebrates. The hemimetabolous species richness is greater in mid- to upper-drainage streams and play a larger role in the diet of juvenile chinook¹⁹⁷ and coho salmon⁴⁴². Although holometabolous invertebrates are dominant in lower-drainages, species in the family Chironomidae are present in all habitats and are a significant food source to salmon in early freshwater stages^{399, 440, 100}. Invertebrate consumption is based on handleable body size and abundance of individuals. Larval and adult insects are the most common forms of food found in natal areas of the freshwater habitat of salmon, with differential diet preferences exhibited by the shorter (chinook) and longer freshwater (cutthroat) life cycle salmon species. Feeding

habits gradually change during downstream migration with the addition of large-bodied prey items. However, a certain component of the diet is comprised of accustomed food sources like Chironomidae that were consumed during the early freshwater life cycle.

All species of salmon fry consume some life stages of dipterans, primarily Chironomidae, during the freshwater life phase¹⁷⁸. Stonefly and mayfly nymphs are consumed by pink, chum, and chinook salmon fry. Coho fry are suspension and surface feeders whose diet is predominately terrestrial insects. Ecologically important freshwater invertebrates in coho natal habitat are emergent and flying insects such as mayflies, stoneflies, and midges (Chironomidae). The rapid migration of chinook fry to the river estuary introduces terrestrial homopterans (leaf hoppers and aphids) into their diet. Additional details of prey items during the freshwater cycle can be found in Simenstad et al.⁴⁶⁶, Shreffler et al.⁴⁶¹, Scott and Crossman⁴⁴⁹, Chapman and Bjornn⁹⁵, Mundie³³⁸, Martin²⁹⁴, Peterson³⁹⁹, Friesen¹⁵⁵ and Groot and Margolis¹⁷⁸. The influence of riparian vegetation along streams and estuaries appears to be an important factor in determining abundance and type of terrestrial insects on which salmon are able to forage.

Salmon as a Food Source For Aquatic Macroinvertebrates

Freshwater macroinvertebrates such as caddisflies, stoneflies, and midges are involved in processing the microbially conditioned salmon carcasses. Bilby et al.⁴⁷ observed a significant contribution



Plate #12. Aquatic insects feeding on salmon carcasses. (Photo by: Jason Walter and Brian Fransen).

of nitrogen from spawning salmon to the collector-gatherer invertebrate community. Increases in aquatic invertebrate density from the introduction of salmon carcasses⁵⁶⁴ stimulated feeding by early life stages of select salmon species⁴⁸. Other stages of the salmon life history contribute to the invertebrate food base. Nicola³⁷⁶ observed the stonefly nymph, *Alloperla* (Plecoptera), scavenging dead pink and chum salmon embryos and alevins. Also, Elliott and Bartoo¹³¹ found the midge, *Polypedilum* (Diptera), associated with dead pink salmon embryos and alevins.

Freshwater invertebrate shredder abundance increases in the presence of salmon carcasses⁵⁶⁴. Non salmon-bearing streams support a limited abundance of shredders mediated through input of leaf matter.

This organic food base must first be conditioned by the microbial community to increase palatability to shredders. Cool water temperatures characteristic of coastal streams slow the microbial decay of the leaf litter food source resulting in limitations in distribution and abundance of the shredder community. The appearance of salmon and the additional influx of biomass to streams appears to be a controlling factor for shredder species. However, the role of shredders in the presence of salmon carcasses continues to be investigated. Bilby et al.⁴⁷ found no significant concentrations of carbon contributed from decaying carcasses in the shredder community. Undigestible animal tissue consumed by shredders was excreted as fine particulate organic matter. Nutritive food value for shredders may have been derived primarily from the microbial community on decaying carcasses. Aquatic insects of the collector-gatherers group typically benefit from the activity of shredders^{316, 47}.

The relationship between invertebrates and salmon can be complex. Functions of invertebrates have not yet been fully defined, but we know they are essential to salmon survival. Traditional functional groupings of invertebrates have been helpful in understanding their ecological roles; however, continuing research suggests that many genera are capable of filling other functional groups if given the opportunity and resources (Plotrikoff, personal communication). Invertebrates complete a loop beginning as recipients of food from adult salmon carcasses that, in turn, fuel the growth and survival of early stages in the salmon's life cycle.

Vertebrate Wildlife

Vertebrate Wildlife And Salmon

Anadromous salmon provide a rich, seasonal food resource that directly affects the ecology of both aquatic and terrestrial consumers, and indirectly affects the entire food-web that knits the water and land together. Wildlife species have likely had a very long, and probably co-evolutionary, relationship with salmon in the Pacific Northwest. In their *Natural History of Washington Territory and Oregon*, Suckley and Cooper⁵⁰² wrote of the California condor:

“The Californian vulture visits the Columbia river in fall, when its shores are lined with great numbers of dead salmon, on which this and the other vultures, besides crows, ravens, and many quadrupeds, feast for a couple of months.”

The "Five Mile Rapids" prehistoric archaeological site along the banks of the Columbia River, five miles east of the Dalles, Oregon, yielded bones from at least 63 individual California condors, plus remains of turkey vultures, cormorants, bald eagles, and gulls¹¹⁰. Carbon-14 dating placed materials at this site from 10,000 to 7,500 years before present^{323, 474}. Miller³²¹ suggested that these birds were attracted to the site by the presence of abundant living and dead salmon and human refuse resulting from fishing.

The life stages of salmon (i.e., eggs, fry, smolts, adults, and carcasses) all provide direct or indirect foraging opportunities for terrestrial, freshwater, and marine wildlife. While sometimes abundant and somewhat dependable from year to year, the availability of salmon to wildlife is largely seasonal in nature. The high seasonal variability in a particular food resource is reflected in the opportunistic foraging of many wildlife consumers - however, "opportunistic" is not a synonym for biological unimportance. Thus, one could hypothesize that while many wildlife species could develop important food-web relationships with salmon, few wildlife species would likely be able to form an ecological "dependence" on salmon. Only those species which are highly mobile, or are able to capture, consume, and store (in body tissues) substantial quantities of salmon biomass in a short period of time would be likely to develop a strong direct ecological dependence on salmon. It is more probable that the majority of wildlife which directly consume salmon will have flexible foraging strategies, utilizing salmon when available, and alternate food sources during other times of the year.

Indirect relationships develop when a food resource is providing foraging opportunities to a secondary consumer, an example in our case is reflected by peregrine falcons which eat gulls that feed on salmon carcasses. As salmon are a concentrated resource, this will serve to concentrate otherwise dispersed wildlife species (e.g., bears). In this scenario, there may well be competition, parasitism, or other aggressive interactions between or among wildlife species. Some of these interactions, e.g., bald eagles disturbing common mergansers, serve to benefit salmon by reducing predation. The magnitude of the salmon-wildlife interaction warrants special examination and calls attention to the pervasive occurrence of these important ecological functions linkages across the region. The loss or severe depletion of anadromous fish stocks could have major effects on the population biology (i.e., age class, longevity, dispersal ability) of many species of wildlife, and thus, on the overall health and functioning of natural communities over the majority of the region.

Research on predator-prey interactions in which anadromous fish are the prey has strongly emphasized the

effects of predation on the fish populations^{576, 570, 571, 559, 331, 430, 225}. Many existing studies describe predatory species as competitors of human harvesters and attempt to control the rate of predation to maximize human consumption. Focusing on the important interplay between salmon and wildlife populations will help to reverse this perspective. In the following sections, we discuss the relationships between salmon and their vertebrate consumers, and the salmon's role in enhancing ecological functions involving wildlife in terrestrial, freshwater, and marine systems.

Wildlife Species With A Relationship To Salmon

Johnson et al.²³³ examined the relationships between the Pacific salmon and 605 species of terrestrial and marine mammals, birds, reptiles, and amphibians currently or historically common to Washington and Oregon. They found a positive relationship between salmon and 138 species of wildlife, the relationship was “unknown” for 60 species, and a determination of “no relationship” was made for 407 species (Table 3). Where a relationship existed, they identified both the type(s) of relationship and the stage(s) of the salmon life cycle to which it applied. Of the 138 species with a relationship to salmon, 9 species were categorized as having a *Strong, Consistent* relationship (Appendix I), 58 as *Recurrent* (Appendix II), 25 as *Indirect* (Appendix III), and 65 as *Rare* (Appendix IV). This tally totals more than 138 because 19 species had more than one type of relationship with salmon.

Of the 138 wildlife species, 88 were characterized as having a routine relationship (combination of species with *Strong, Consistent, Recurrent;* and *Indirect*) with salmon. Of these 88 species, there were 25 mammals (8 of these were marine mammals), 60 birds, 2 amphibians, and 1 reptile.

The relationship categories are briefly described as follows:

1) *Strong, Consistent Relationship*. Salmon play (or historically played) an important role in this species distribution, viability, abundance, and/or population status. The ecology of this wildlife species is supported by salmon, especially at particular life stages or during specific seasons. Timing of reproductive activities, and daily or seasonal movements often reflect salmon life stages. Relationship with salmon is direct (e.g., feeds on salmon, or salmon eggs) and routine. The relationship may be regional or localized to one or more watersheds. Examples: A significant portion of the diet of killer whales is adult salmon (*Saltwater* stage); common mergansers may congregate to feed on salmon fry (*Freshwater Rearing* stage) when they are available.

2) *Recurrent Relationship*. The relationship between salmon and this species is characterized as routine, albeit occasional, and often tends to be in localized areas (thus affecting only a small portion of this species population). While the species may benefit from this relationship, it is generally not considered to affect the distribution, abundance, viability, or population status of this species. The percent of salmon in the diet of these wildlife species may vary from 5% to over 50%, depending on the location and time of year. Example: turkey vultures routinely feed on salmon carcasses, but feed on many other items as well.

3) *Indirect Relationship*. Salmon play an important routine, but *indirect* link to this species. The relationship could be viewed as one of a secondary consumer of salmon; for example, salmon support other wildlife that are prey of this species. This includes aspects such as salmon carcasses that support insect populations that are a food item for this species. Example: American dipper feed on aquatic insects that are affected by salmon-derived nutrients. The hypothesis of an *indirect* relationship between an aerial insectivore and salmon was supported by the presence of two or more of the following characteristics of the insectivore: (1) riparian obligate or associate, (2) feeds below or near the canopy layer of riparian trees, (3) known or perceived to feed on midges, blackflies, caddisflies, stoneflies, or other aquatic insects that benefit from salmon-derived nutrients, and/or (4) feeds near the water surface. While this category includes general aspects of salmon nutrient cycling in stream/river systems, we are not including or examining the role of carcass-derived nutrient cycling on lentic system riparian and wetlands vegetation, and subsequent links to wildlife.

4) *Rare Relationship*. Salmon play a very minor role in the diet of these species, often amounting to less than 1 percent of the diet. Typically, salmon are consumed only on rare occasions, during a shortage of the usual food and may be especially evident during El Niño events. As salmon are often present in large quantities, they may be consumed on rare occasions by species that normally do not consume them. Examples: red-tailed hawks are known to consume salmon carcasses in times of distress; trumpeter swans are primarily vegetarians, but on rare occasions will consume eggs, parr, as well as salmon carcass tissue.

5) *Unknown Relationship*. A relationship between this species and salmon may exist, but there is not enough information to determine the scope or scale of the relationship at this time. Example: while it is logical to speculate that riparian feeding bats may feed on salmon-derived insects, aspects of seasonality of both bats and salmon carcasses are relevant, as is the nocturnal flight behavior of the insects. Do bats and salmon carcasses coincide seasonally, and if so, are salmon-derived insects actually available to feeding bats? At this time, the evidence for this relationship is inconclusive and remains to be examined.

6) *No Relationship*. There is no recognized or apparent relationship between salmon and this species.

As part of the same study, Johnson et al.²³³ reported 60 species as having an "unknown" relationship with salmon (Appendix V), suggesting that the diets of these species *in Washington and Oregon*, were not understood well enough to characterize their relationship with salmon. Additional observations on the diets of these species will help determine whether the relationships of these species with salmon is routine, a rare and unusual event, or whether a relationship exists at all. Johnson et al.²³³ identified 407 species as having "no relationship" to salmon (Appendix VI).

Table 3. Relationships between Pacific salmon and 605 species of wildlife in Washington and Oregon. There were 137 species with a positive relationship with salmon (i.e., combined total for species with *Strong, Consistent, Recurrent, Indirect, and Rare* relationships). The total number of individual wildlife species for columns and rows are shown in parenthesis; the number of species shown in the rows and columns may not equate to the numbers shown as totals as 19 species had more than one type of relationship with salmon, and 73 species are associated with salmon at more than one life stage.

Salmon life stage	Strong, Consistent relationship	Recurrent relationship	Indirect relationship	Rare relationship	Unknown relationship	No relationship
Incubation – eggs and alevin (23)	2	10	1	10		
Freshwater rearing – fry, fingerlins, parr (49)	4	31	4	10		
Saltwater – smolts, immature adults, adults (63)	6	36	5	19		
Spawning (16)	5	10	0	1		
Carcassess (82)	5	28	22	38		
	(9)	(58)	(25)	(64)	(60)	(408)

Wildlife Response To Salmon Congregations

The numerical response of predators to salmon congregations is often substantial, sometimes spectacularly so. The ability of wildlife species to concentrate at salmon sites is more than opportunistic foraging, it has significant biological importance. Anadromous fishes (including their eggs) are a major source of high-energy food that allows for successful reproduction and enhanced survival of adults and juveniles of many wildlife species, and support for long-distance migrant birds. Wildlife movements to salmon congregations can be seasonal (e.g., bald eagles along the Skagit River in Washington), or depending on the situation (e.g., hatchery fish released during an El Niño high food-stress seabird breeding season) can occur within a matter of hours. Perhaps as noteworthy, but much harder to detect, is that some wildlife species that have been reported to group at salmon sites in other areas (e.g., black bears in southeast Alaska) do not appear to be doing so with any regularity in Washington and Oregon. This may well be reflecting the depressed nature of some salmon stocks rather than the inherent behavior of the wildlife species. Of the 88 species with a link to salmon ²³³, 43 species (37 birds, 6 mammals) concentrate or form loose aggregations at salmon sites (Table 4). Some reasons why other species do not congregate at salmon streams are: strong territoriality (e.g., great blue heron), foraging strategies which require above-water structures or perches (e.g., belted kingfisher), and limited movement capabilities (e.g., shrews).

Table 4. Wildlife species that have been observed or are perceived to aggregate at salmon congregations in Oregon and Washington.

Artic tern	Common raven	Bald eagle
Forester's tern	Turkey vulture	Bank swallow
Western grebe	Western gull	Barn swallow
American white pelican	Herring gull	Black bear (now questionable)
Brandt's cormorant	Thayer's gull	California sea lion
Common tern	American crow	Cliff swallow
Common merganser	Tufted puffin	Grizzly bear (now questionable; may not be enough salmon available to congregate at).
Brown pelican	California gull	
Caspian tern	Northwestern crow	Harbor seal
Red-breasted merganser	Common murre	Killer whale
Elegant tern	Black-billed magpie	Northern (Steller) sea lion
Double-crested cormorant	Rhinoceros auklet	Northern rough-winged swallow
Barrow's Goldeneye	Ring-billed gull	Tree swallow
Clark's grebe	Glaucous gull	Violet-green swallow
Common Goldeneye	Glaucous-winged gull	

That some wildlife species concentrate at salmon areas is well established. The fact that other wildlife species do not concentrate at salmon areas may reflect salmon declines. We offer the following examples relevant for Washington and Oregon.

Bald Eagle Suckley and Cooper ⁵⁰² state: “*This noble looking bird is exceedingly abundant in Oregon and Washington Territories, and in certain localities, especially during the salmon season, may be found in great numbers.*” The North Fork of the Nooksack River (coastal Washington) currently hosts one of the largest and most visible concentrations of wintering bald eagles in the lower forty-eight states. Peak concentrations (100 or more eagles) occur along the Nooksack ²⁶³ and Skagit ²¹⁸ rivers with December- and January-spawning chum salmon.

Caspian Tern The first breeding record of Caspian terns along the Oregon/Washington coast was a colony of 50 pairs in Grays Harbor, Washington in 1957 ⁶. This, and other nesting colonies (mid-1950's through early 1990's) along the Washington coast in Grays Harbor, Willapa Bay, and near the mouth of the Columbia River, have been abandoned or destroyed by human actions ^{429, 430, 479}. A colony of Caspian terns originally settled on Rice Island, a dredge material disposal island in the lower Columbia River, in 1987. In 1997, an estimated 14,000 terns used this island for nesting and/or roosting during the 80-100 day (April-July) breeding season ^{429, 430}. This represents the largest known colony of Caspian terns in North America, and possibly the world. In 1997, the terns appeared to be largely dependent on juvenile salmon (roughly 75% of the diet), consuming an estimated 14.5 million smolts, the majority being hatchery fish ^{429, 430}. Tern nesting success was very low (roughly 5%) in 1997; predation on adult terns by bald eagles, and gull predation on tern eggs and chicks (caused by eagle and researcher disturbance) were the primary causes ^{429, 430}.

Common Murre The common murre is a seabird that nests in large colonies along the Oregon coast; colonies in Washington have undergone significant declines in the last decade. It is only an occasional consumer of salmon, as the vast majority of its diet is other small marine fishes. A severe El Niño event occurred in 1983 during the seabird nesting season along the eastern Pacific coast, with the majority of common murre (and other seabirds species) either not attempting to nest or abandoning their nests once initiated ^{27, 210}. Adult survival was also greatly reduced ¹⁶⁹. Oregon Aqua-Foods, Inc. had released a total of 2 million or more salmon smolts into the Yaquina Estuary at roughly 2.5 day intervals between June and August since 1977 ²⁸. Murre were more numerous at the mouth of Yaquina Estuary for the first two days post-release in July of 1983 than in July of 1982 (a non-El Niño year), as they were drawn in to feed on the released salmon. First day post-release averages of murre were 3,710 in 1983 and 3,053 in 1982. In August of 1983 however, murre numbers were significantly lower than in 1982 (average 1983 = 106; 1982 = 1,860), as murre had begun moving north earlier to feed on other food resources ²⁷. In summary, although not a primary food resource, murre will make use of salmon resources during food-stress conditions.

Black Bear Contrary to popular image, Washington and Oregon black bears rarely congregate at salmon sites. Poelker and Hartwell ⁴⁰⁶ reported on three diet studies of black bears in western Washington for the time periods of 1952-54 and 1968, and found that fish represented 5.0% of the diet. In their treatise on land mammals, Verts and Carraway ⁵³⁵ describe black bears in Oregon as being largely herbivorous, and do not mention salmon as part of their diets. Cederholm et al. ⁹¹ found black bears on Washington's Olympic Peninsula to heavily consume salmon carcasses. In northern California, Kellyhouse ²⁴⁴ found evidence of salmon in 10% of black bear fecal samples analyzed from spawning areas. The California Department of

Fish and Game⁸⁴ also reported black bears taking advantage of anadromous fish runs. The strong link between black bears and salmon was demonstrated in the Anas Creek drainage of southeastern Alaska (D. Chi, personal communication). This effort studied the behavior and activity patterns of black bears (n = 40 individuals) which had established movement patterns according to salmon migrations. Bears arrived in the lower reaches of the creek in June to begin feeding on spawning salmon, and stayed through August and early September to feed on salmon carcasses. Thirteen of the bears were radio-marked and their movements indicated that they were moving in from at least eight miles (12.9 km) away. Other bears were assumed to be coming in from further away. The general lack of salmon in the published accounts of black bear diets across Washington and Oregon^{exception: see 91} is somewhat counter to the observed salmon use in adjacent regions. Radio-marked bears in Washington (G. Kohler, personal communication) and Oregon⁵²⁸ have been found to move to and congregate at higher elevations in the fall to feed on huckleberries (i.e., forming “traditional use areas”), thus, one could reasonably conclude that if salmon were to be found in substantial and predictable numbers, bears in Washington and Oregon, like those studied by Chi in Alaska, would also establish traditional use areas around salmon. Black bears in western Washington typically den by 1 November and emerge around 1 April, thus salmon runs occurring during the winter will not be available to bears. Recent bear studies in western Washington have included the Humptulips, Wishkah, Wynoochee, and Quinault Rivers and while these rivers hold low levels of hatchery-based salmon, bears do not congregate along them (G. Kohler, personal communication). D.H. Johnson (unpublished data) summarized 1990-1998 hatchery return data of adult salmon for the Humptulips river system in western Washington. These fish return to the hatchery facility as early as late-September (most begin around mid-October), and as typical with most, were done spawning by mid-December. While there are additional fish in this system, an average number of 217 (range 95-320) chinook, 6,496 (range 177-10,195) coho, and 165 (range 51-339) steelhead returned annually to the hatchery. The substantial majority of these fish species return to spawn after November 1st (the average date of bear denning) and are not available to bears; an average of 73 chinook, 1,264 coho, and 0 (zero) returning steelhead were available to black bears. Here, “available” means simply present in the river system, and not located at spawning redds. In summary, bears have a strong relationship to salmon where they have access to them, but it appears that in substantial measure, current salmon populations do not represent a predictable food supply to bears in Washington and Oregon.

Review Of Wildlife Relationships By Salmon Life Stages

For the 138 species with a relationship to salmon, Johnson et al.²³³ identified the salmon life stage(s) involved for each species. In this study, the five general life history stages of salmon were identified as: (a) *Incubation* (egg and alevin), (b) *Freshwater Rearing* (fry, fingerling, and parr), (c) *Saltwater* (smolt, subadult, adult), (d) *Spawner*, and (e) *Carcass*. The number of wildlife species associated with each (in parenthesis) were: *Incubation* (23), *Freshwater Rearing* (49), *Saltwater* (63), *Spawning* (16), and *Carcass* (83); this tally of wildlife species totals more than 138 because 73 species are associated with salmon at more than one life stage (Appendixes I, II, III). See Appendix VII for a complete list of published and unpublished observations of wildlife predators and scavengers on salmon at various stages of their life.

Incubation Stage (eggs and alevin)

Twenty-three wildlife species are linked to salmon at this stage. Twenty-two wildlife species are direct consumers of “drift eggs” (eggs not buried in redds) or alevin (2 amphibians, 1 reptile, 19 birds, and 1 mammal); and 1 bird (bald eagle) is an indirect consumer of eggs/alevin, feeding on the waterfowl that consume eggs and alevin.

Freshwater Rearing (fry, fingerling, and parr)

Forty-nine wildlife species are linked to rearing salmon, including 2 amphibians, 5 reptiles, 34 birds, and 4 mammals. Forty-five of these species are direct consumers of salmon and 4 species (bald eagle, gyrfalcon, peregrine falcon, and snowy owl) are indirect consumers, feeding on terns, waterfowl, gulls, and other animals that eat rearing salmon.

Saltwater (smolt, subadult, adult)

Sixty-three wildlife species are consumers of salmon at this stage (51 birds and 12 marine mammals). Fifty-eight of these species are direct consumers of salmon and 5 species are indirect consumers. This list is somewhat expansive due to the geography being included, that is, the estuarine and all marine water habitats.



Plate #13. Garter snake eating a salmon smolt, unknown Olympic Peninsula stream, Washington. (Photo by: Jim Rozell, deceased).

Spawner

Sixteen species of wildlife are consumers of spawning salmon (6 birds and 10 mammals). This list is relatively small, as few wildlife species are physically capable of capturing and handling live, adult fish. The gray wolf and grizzly bear are on this list, but both have undergone significant range contractions and declines in their abundance (e.g., both are extirpated from Oregon and significantly reduced in Washington).



Plate #14. River otter, a known predator of salmon. (Photo by: Charles J. Gibilisco).

Carcass

Carcasses are linked to the largest group of wildlife consumers of any salmon life stage, with 83 species (1 reptile, 50 birds, and 32 mammals) being consumers of carcasses and/or carcass-derived insects. Body sizes of these animals range from shrews to grizzly bears. Seventy-one species of wildlife (1 reptile, 38 birds, and 32 mammals) are direct consumers of carcasses; 22 species (14 birds and 8 mammals) are consumers of carcass-derived insects; and 10 species (2 birds and 8 mammals) are consumers of both carcasses and carcass-derived insects.



*Plate #15. Gull eating a chum salmon (*O. keta*) carcass at Kennedy Creek, Washington. (Photo by: Jeff Cederholm).*



*Plate #16. Pacific Giant Salamander, a known predator of juvenile salmon.
(Photo by: William Leonard).*

Continued documentation of wildlife species-salmon interactions, especially of the 60 species having an “unknown” relationship, will provide vital information for ongoing developments in ecologically-based salmon spawner escapement research and prescriptions for riparian management practices. As Key Ecological Functions (KEFs) are identified through such research ²⁹², tools for informed decisions will be made available to fish and land managers operating under an ecosystem context.

See Appendix VII for a complete list of published and unpublished observations of wildlife predation and scavenging on salmon at various stages of their life.

Key Ecological Functions (KEFs) Provided To The Ecosystems Through Salmon-Wildlife Interactions

In striving to manage for healthy and sustainable ecosystems, we simultaneously are striving to provide for the full range of ecological functions that these systems provide. Key ecological functions (KEFs) refer to the main ecological roles of a species (or group of species) that influence diversity, productivity, or sustainability of ecosystems³³⁴. A given KEF can be provided by a single species or shared by many species, and a given species can have several KEFs. Main categories of KEFs include trophic relations; herbivory; nutrient cycling; interspecies relations; disease; pathogen and parasite relations; soil relations; wood relations; water relations; and vegetation structure and composition relations. Building upon work by Marcot et al.²⁹¹, Marcot and Vander Heyden²⁹² characterized the key ecological functions for each of the 605 common wildlife (i.e., terrestrial and marine birds, mammals, reptiles, and amphibians) species in Washington and Oregon. Several questions can be thus posed:

- *In what way does providing for salmon also provide for a wider array of ecological functions of wildlife species associated with salmon?*
- *What are those functions?*
- *How do different kinds of salmon-wildlife relations, and different salmon life stages, provide for an array of ecological functions?*

This somewhat innovative analysis describes the functional links among fish and wildlife species across aquatic and terrestrial communities. To conduct this analysis, we queried the database matrixes on salmon-wildlife relations, key ecological functions of wildlife species, and habitats used by wildlife²³⁶. The general conclusion is that salmon provide a causal mechanism for movement behaviors and a nutrient source for a variety of wildlife species, which in turn perform a surprisingly broad array of ecological functions²⁹² across a wide span of habitats. For this analysis, one can think of the array of ecological functions performed by these wildlife species as a “functional web”. It focuses on salmon in their various life stages, and extends well beyond the aquatic realm to influence the diversity, productivity, and ultimately sustainability of habitats and ecosystems throughout Washington and Oregon.

Wildlife With Strong Consistent Links to Salmon

The 9 species of wildlife with strong consistent links to salmon (bald eagle, American black bear, Caspian tern, common merganser, grizzly bear, harlequin duck, killer whale, osprey, and river otter) comprise a functional group of “salmon-eaters” with close affinities to salmon. There are 32 primary wildlife-habitats across Washington and Oregon²³⁶; Figure 7 summarizes the occurrence of these 9 wildlife species by habitat. Not surprisingly, most of these 9 species inhabit freshwater and marine habitats, but some of them also occur across the range of inland forest, woodland, shrubland, and grassland habitats. It is of interest that from 1-7 of these 9 species can be found in each of the 32 habitats (Figure 7). In this way, salmon provide for a set of wildlife species that occur well beyond just salmon-inhabited aquatic systems.

In addition, the full set of key ecological functions performed by these 9 species also extends beyond the aquatic system. Each of these species provides a set of ecological functions to the various array of habitats that they occur within. Figure 8 depicts the collective range of ecological functions that these 9 wildlife species provide to the number of habitats that they occupy. The functions range from various trophic, organismal, and wood and soil relations. Some functions are more widespread (occur in more habitats)

Species Richness by Habitat
Strong Consistent Salmon Links




Figure 7: Occurrence by number of vertebrate wildlife species in the 32 wildlife habitats in Washington and Oregon, as used by the nine wildlife species with a strong consistent relationship to salmon.

than are other functions. Examples of some widespread functions are potential control of vertebrate populations (through predation), carrion feeding, piscivory (fish-feeding), invertebrate feeding (including insectivory), omnivory, transportation or dispersal of seeds and animals, creation of terrestrial runways used by other species, and secondary use of burrows created by other species.

Species Richness by Function Strong Consistent Salmon Links




Figure 8: The array of key ecological functions performed by the nine vertebrate wildlife species with a strong, consistent relationship to salmon, across the 32 wildlife habitats in Washington and Oregon.

All Wildlife With Links to Salmon

What of the full set of species showing either strong consistent, recurrent, and/or indirect links to salmon? (Some species have more than one type of relation because they use more than one salmon life stage). Table 5 lists key ecological functions of wildlife more or less unique to each type of salmon-wildlife link. Each of the 3 types of relations provides for some unique set of ecological functions. For example, wildlife species indirectly linked to salmon can provide the following ecological functions: fungivory (fungus-eating), tertiary consumption or secondary predation, prey source; regulate insect populations through predation; serve as interspecific host for avian nest parasites, and create primary small ground burrows. These functions are performed not at all, or by far fewer wildlife species, by the wildlife species with strong consistent links or occasional links to salmon.

Table 5. IMPORTANCE OF TYPES OF SALMON-WILDLIFE RELATIONS TO KEY ECOLOGICAL FUNCTIONS.

1. STRONG CONSISTENT RELATIONSHIPS ARE IMPORTANT FOR:

- trophic relations:
- primary consumption:
 - spermivory
 - grazing
 - frugivory
 - root feeding
- organismal relations:
 - controlling vertebrate populations
 - dispersing seeds, fruits, invertebrates, vascular plants
 - creating feeding opportunities for other species
 - primary cavity excavation in trees and snags
 - primary creation of large ground burrow
 - primary creation and secondary use of ground runways
- wood relations:
 - fragments standing and down wood
 - kills standing trees (creates snags)

2. OCCASIONAL RELATIONSHIPS ARE IMPORTANT FOR:

- trophic relations:
 - pirating food
- organismal relations:
 - secondary use of aerial and aquatic structures created by other spp.
- disease relations:
 - carrier of domestic animal disease
- soil relations:
 - improves soil structure and aeration by digging and burrowing

3. INDIRECT RELATIONSHIPS ARE IMPORTANT FOR:

- trophic relations:
 - primary consumption:
 - fungivory
 - tertiary consumption
 - prey relations:
 - providing prey for predators
 - organismal relations:
 - controlling insect populations
 - serves as interspecific host for avian nest parasite
 - primary creation of small ground burrows
-
-

Table 6. IMPORTANCE OF SALMON LIFE STAGES TO KEY ECOLOGICAL FUNCTIONS.
(Listed are functions unique to each life stage category).

1. INCUBATION STAGE IS IMPORTANT FOR:

1. organismal relations
secondary cavity use

2. FRESHWATER REARING STAGE IS IMPORTANT FOR:

(no specific function is mostly supported by this stage)

3. SALTWATER STAGE IS IMPORTANT FOR:

2. organismal relations:
creates aerial structures used by other species
creates aquatic structures used by other species

4. SPAWNING STAGE IS IMPORTANT FOR:

3. trophic relations:
4. primary consumption:
spermivory
grazing
frugivory
root feeding
bark/cambium/bole feeding
5. organismal relations:
controlling vertebrate populations
creating feeding opportunities for other species
primary cavity excavation
primary excavation of large ground burrows
primary creation of ground runways
6. wood relations:
fragments standing and down wood
kill standing trees (creates snags)

5. CARCASS STAGE IS IMPORTANT FOR:

7. trophic relations
 8. primary consumption
 9. organismal relations:
controlling insect populations
serves as interspecific host for avian nest parasite
-
-

What this means is that different degrees of salmon-wildlife relations provide for some unique kinds of wildlife ecological functions. Only the full set of all wildlife-salmon link relations can provide for all collective functions. Thus, to manage the full set of all ecological functions, one should not focus solely on those few wildlife species with strong consistent links to salmon, but on all types of links.

How Salmon Life Stages Provide for Ecological Functions

In a similar way, most of the 5 life stages of salmon provide for a unique set of wildlife species and their ecological functions (Table 6). For example, wildlife associated with the incubation stage of salmon include secondary cavity users and primary excavators of small ground burrows; these two ecological functions are not provided, or only poorly provided, by wildlife species associated with any of the other salmon life stages. Thus, to manage for the full set of ecological functions, one should focus on providing all life stages of salmon.

Managing the Functional Web

So what is the manager to do with this information? For one, be aware that salmon can be viewed as the center of a broad “functional web” of wildlife and their ecological roles. Such roles extend well past the salmon populations and aquatic habitats themselves, and likely influence the structure and processes of the communities and ecosystems in which they reside, thus a “keystone” species ⁵⁵⁹.

Second, one can use the information presented here and in the species data matrixes to list the collective set of habitat elements and conditions used by wildlife species associated with salmon. For example, one can link the list of wildlife associated with salmon life stages likely to be found in low order headwater streams, and determine the set of habitat elements used by this set of wildlife species, by habitat type, and then establish habitat-specific management guidelines to provide for such habitat elements over time. Maintaining such habitat elements and conditions would help maintain the full salmon-wildlife functional web.

Third, one can begin to predict – or, at least pose tentative management hypotheses about – which ecological functions may be in jeopardy if the wildlife that performs such functions are not maintained. That is, one can now determine which wildlife species may be influenced by altering salmon populations and habitats that imperil specific salmon life stages, and the set of ecological functions associated with such wildlife species. In some cases, other wildlife species not associated with salmon may also perform some ecological function, but never in exactly the same manner and in the same set of habitats and habitat elements.


SALMON FISHERIES

Washington and Oregon salmon fisheries includes commercial, recreational, and treaty harvests that occur in the states' rivers, inland lakes, inland marine waters, coastal embayments, and at sea. Fishing is an important source of mortality, both for immature fish in the ocean and mature fish on their return to freshwater to spawn (Figures 9, 10, 11, 12, 13, and 14). Understanding past fishery effects is important because man's increasing efficiency as a predator augments the natural mortality rates that salmon encounter in natural situations. Because of the presence of humans, salmon are significantly less available to a wide variety of natural consumers in the freshwater, terrestrial, and marine



Plate #17. Commercial salmon fishing vessel. (Photo by: Washington State Historical Society, Tacoma, WA. Curtis. Photo Negative No. 63811).

Plate #18. Steelhead sport catch on the Hoh River, Washington. (Photo by: Jeff Cederholm)



environments, and thus these ecosystems are suffering. According to the NRC ³⁶⁵, salmon mortality caused by human activities and natural factors usually exceed fishing mortality. Thus, although factors other than fishing have a major effect on the production of adult fish, fishing is still the easiest salmon mortality factor to control ³⁶⁵.




Figure 9. Total commercial catch, in millions of pounds, of Lower Columbia River Salmon from 1870's to 1990's ^{390, 379, 387b, 540}.




Figure 10. Estimated and reported total catch, in numbers, of Puget Sound salmon from 1890's to 1990's ^{390, 57a, 538b}.




Figure 11. Washington total ocean commercial troll catch, in numbers, of salmon from 1930's to 1990's ^{390, 538b, 388a}.







Figure 12. Washington ocean sport catch, in numbers, of salmon from 1950's to 1990's ^{390, 538b, 388a}.

The salmon species complexes in the Columbia River, Puget Sound, and Coastal Washington and Oregon, except for Puget Sound chum, have never recovered to the numbers that existed when commercial harvest was initiated. Increased hatchery production did enhance some fisheries, notably the rise of the coho salmon runs in the 1960s, and chum salmon runs in Puget Sound. In general, however, artificial propagation has failed to rebuild the runs to former levels³⁹⁴, and in some instances likely contributed to the further decline of wild stocks²⁰⁹. Hatchery production has also contributed to the harvest of wild salmon by creating socioeconomic incentives that maintain mixed-stock fisheries.




Figure 13. Estimates of Oregon coastal coho salmon production and harvest from 1970's to 1990's^{387b, 388a}.



Figure 14. Total commercial catch, in pounds, of steelhead in Oregon coastal streams from 1920's to 1940's ^{387a}.

Admittedly, managing salmon fisheries is a challenge, and past management approaches were generally commodity/extraction-based; however, this approach has significantly contributed to the decline of wild stocks ⁵⁴⁴. Wild salmon life history characteristics that contribute to this challenge include wide geographic distribution with extensive feeding and spawning migrations; complexity of life history forms, ages, and sizes; and the death of most adult fish after spawning only once. Fishing activities that contribute to the problem include: indirect mortality due to catch and release of undersized fish (bycatch), out-of-state domestic and foreign interception, conflicts among user groups, and the mixed-stock fisheries.

A major dilemma that fishery resource agencies find themselves in is how to selectively harvest hatchery salmon, while still meeting spawning escapement goals of wild stocks of salmon. Hatchery-produced salmon co-mingle with wild salmon in ocean waters, and as a result, a mixed-stock fishery is created. If harvests are allowed in such mixed-stock fisheries, then wild and hatchery fish will be caught at rates that only hatchery fish can sustain. Wild salmon cannot withstand the high hatchery exploitation rate because they are exposed to a full range of natural and human-caused selection pressures and mortalities. Hatchery fish are sheltered from mortality factors that normally occur during incubation and freshwater rearing. At smolt migration, many more progeny are still alive per hatchery female than per wild female. Thus, hatchery populations can maintain smolt output at a consistent level with far fewer spawning adult fish than can wild

populations. This condition enables hatchery stocks to withstand higher rates of harvest than wild stocks; however, even hatchery stocks eventually succumb to the high exploitation rates through changes to smaller adult size ⁵¹⁷, or different time of return ¹⁹⁰. Therefore, high fishing exploitation rates are associated with fishery resource management agency policies, and agency policies also determine hatchery policies and practices. The current demand by certain interests to protect wild stocks can be in direct conflict with the agency mandate to enhance or supplement current stocks with hatchery-produced fish to ensure sustainable harvest for the major user groups. This conflict will have an important bearing on future management of salmon fisheries and hatchery practices. Live capture selective fishing, including live release of wild unmarked fish and retention of marked hatchery fish, is potentially an option ²⁷⁹.

Spawning Escapement Goals

Spawning escapement goals (the number spawners required to perpetuate the population ²⁵⁴) are set by fishery managers to determine the portion of the estimated returning adult population that can be harvested. Recently Knudsen ²⁵⁴ reviewed the methods used along the west coast to establish wild salmon escapement goals, and found that of 854 management units 8 (1%) were set by methods that were rated excellent (i.e., using methods that combined information in a way that most effectively characterized the management unit's production potential), 142 (16%) were rated as good, 499 (58%) were rated as fair, 13 (2%) were rated as poor, and 192 (22%) had no goals established at all. Analysis of annual spawning escapement data by Konkel and McIntyre ²⁵⁷, collected for naturally spawning salmon populations in the Pacific Northwest between 1969 and 1984, suggests that escapements are down for coho and chum; but up for chinook, sockeye, and pink salmon. In general, escapement trends have been downward since 1970 for all populations, even for those that have achieved their annual escapement goals.

According to the Pacific Fisheries Management Council (PFMC) ³⁸⁸ the state of Washington has established annual escapement goals for coho, chum, and chinook salmon and steelhead, and include wild and hatchery fish. Some escapement goals exist for pink and sockeye salmon that are mostly for wild fish. No determination has been made of the spawning escapement needs of sea-run cutthroat trout. Spawning escapement goals have been established for 98 wild salmon stocks in Washington ^{538a}. These stocks include 30 coho salmon, 29 chum salmon, 27 chinook salmon, 9 pink salmon, and 3 sockeye salmon populations. Fifteen wild stocks of Washington steelhead have established annual spawning escapement goals ¹⁰⁶. Overall, for 113 wild salmon stocks in Washington with established spawning escapement goals, only 46 (41%) met these goals as of the early 1990's ³⁹⁰. Escapements may have improved for some stocks in recent years due to fishery restraints.

In Oregon, escapement goals have been established primarily for chinook and coho salmon and include wild and hatchery fish. No determination of the spawning escapement needs has been made for wild steelhead, chum salmon, pink salmon, or sea-run cutthroat. In the early 1990s, spawning escapement goals were met for only 1 of the 2 populations of wild anadromous salmon in Oregon ³⁹⁰, being met for coastal chinook, but not for coastal coho salmon. Escapements may have improved for some stocks in recent years, due to fishery restraints.

The result of heavy exploitation in the fishery, along with major habitat loss over the past century is that the loadings of marine derived nutrients have been vastly diminished throughout Washington and Oregon rivers. A recent analysis of historical salmon cannery records from west coast rivers by Gresh et al. ^{174a}, indicates that the number of salmon now returning to Washington and Oregon rivers is only 3.3 percent of the historical biomass (132-228 million kg down to 5-7 million kg). These authors conclude that: "This nutrient

deficit may be one indication of ecosystem failure that has contributed to the downward spiral of salmonid abundance and diversity in general, further diminishing the possibility of salmon population recovery to self-sustaining levels.”

The critical factor that salmon harvest managers need to face is how to reduce the annual salmon harvest, and achieve stock-by-stock ecosystem-based spawning escapement goals. With the exception of carcass supplementation programs, there has not been a concerted effort to manage salmon populations for the benefits they provide to the recovery of listed wildlife species (e.g., grizzly bears) or to the broader ecological systems. Salmon spawning escapement goals should not only replace a stock of salmon with sufficient numbers of high quality recruits, but also meet the needs of the broader aquatic and terrestrial ecosystems that depend on salmon for nutrients and carbon influx^{47, 48, 319, 92}. Salmon harvest managers could take a lesson from the worldwide conventions for herring harvest managers where ecosystem function has been explicitly recognized through “...a precautionary, conservative approach to fisheries management.”

541a

For a more thorough review of the magnitude and characteristics of Northwest Pacific Coast salmon fisheries and habitat issues, we recommend the following readings:

Cone, J., and S. Ridlington. 1996. The northwest salmon crisis - A documentary history. Oregon State University Press. Corvallis, OR. 374 pp.

National Research Council. 1996. Upstream - Salmon and society in the Pacific Northwest. Committee on Protection and Management of Pacific Northwest Anadromous Salmonids. Board of Environmental Studies and Toxicology. Commission on Life Sciences. National Academy Press, Washington, D. C. 452 pp.

The Oregon Coastal Salmon Restoration Initiative - Volumes 1, 2, and 3. 1997. The Oregon Plan - Submitted to: The National Marine Fisheries Service March 1997. Salem, OR.

Stouder, D.J., P.A. Bisson, and R..J. Naiman (eds.). 1997. Pacific salmon and their ecosystems: Status and future options. I.T.P. Chapman and Hall International Thomson Publishing. New York, N.Y.

Kaczynski, V.W., and J.F. Palmisano. 1992. A review of management and environmental factors responsible for the decline and lack of recovery of Oregon’s wild anadromous salmonids. Oregon Forest Industries Council. Salem.

Palmisano, J.F., R.H. Ellis, and V.W. Kaczynski. 1993. The impact of environmental and management factors on Washington’s wild anadromous salmon and trout. Prepared for: Washington Forest Protection Association and The State of Washington Department of Natural Resources, Olympia, WA. 371 pp.

Knudsen, E.E., C.R. Steward, D.D. MacDonald, J.E. Williams, and D.W. Reiser. 1999. Sustainable fisheries management - Pacific salmon. CRC Lewis Publishers, Boca Raton, FL. 724 pp.

Lichatowich, J.A. 1999. Salmon without rivers - A history of the Pacific salmon crisis. Island Press, Covelo, Calif. 317 pp.

UNDERSTANDING SALMON RELATIONSHIPS

Salmon As A Key Linkage In Biodiversity And Productivity

Ecological processes have been so altered by human activities, especially in the more densely populated regions, that natural resource and environmental management will need to expand from current site and case-specific methods, to landscape and ecosystem scale approaches. The struggle to develop the tools required for these scales of management has only just begun^{269, 460, 365, 305}. The documents: *From the Forest to the Sea*²⁹⁵, *Forest Ecosystem Management: An Ecological, Economic, and Social Assessment*¹⁵¹, *Pacific Salmon and Their Ecosystems: Status and Future Options*⁴⁹⁹, *An Ecosystem Approach to Salmonid Conservation*⁴⁸⁵, *Fish Habitat Rehabilitation Procedures*⁴⁷⁷ and *River Ecology and Management - Lessons From Pacific Coastal Ecoregion*³⁶² are recommended readings that describe the understanding of natural systems and processes and take a holistic approach to rehabilitation and restoration of watersheds and Pacific Northwest ecosystems.

Anadromous salmon play an important role in maintaining an ecosystem's productivity. The seasonal migrations of millions of salmon between Pacific rim streams and the subarctic Pacific Ocean appear to



Plate #19. Chum salmon carcasses in Kennedy Creek, Washington. (Photo by: Jeff Cederholm).

increase overall terrestrial productivity. Key processes discussed here are the transport of materials, energy and nutrients between marine, aquatic, and terrestrial ecosystems with emphasis on salmon as a transport vector. From a broad ecosystem management perspective, the status of salmon metapopulations is a powerful indicator of human adaptation to boreal biomes^{348, 464, 419}. Sibatani⁴⁶⁴ has suggested that salmon are the “canary in the mineshaft”; the mineshaft in this instance being entire subarctic ocean basin ecosystems. Cederholm et al.⁹² review and discuss the mechanisms of salmon nutrient transport and the significance to terrestrial and freshwater ecosystems. The following discussion suggests a need to understand and apply information on the exchange of materials, energy, and nutrients, between the aquatic and terrestrial ecosystems of the northern Pacific Basin. Such an application would occur at multiple geographic and temporal scales, examining both healthy and depressed salmon populations under varying conditions. Effectively applied, managers would be able to define and achieve long-term ecosystem management success not just for salmon, but for numerous other fish and wildlife resources and the overall health of the environment.

Biomass of Salmon Runs As An Energy Source

Organic matter that supports the trophic system of fresh water ecosystems is provided from both autochthonous and allochthonous sources. Common types of autochthonous sources are: algae, mosses, vascular plants, and phytoplankton. All of these factors are found in freshwater, and generate organic matter through the process of photosynthesis. Common types of allochthonous input include leaves, needles, wood and insects from the terrestrial environment and dissolved organic matter carried in groundwater that enters the water body. Salmon provide an important source of allochthonous organic matter for Pacific Northwest fresh water ecosystems^{248, 47, 239}. Salmon spawning runs transport organic matter and nutrients from the northern Pacific Ocean to their natal spawning grounds. The organic matter and nutrients carried in the biomass of the salmon runs is input to the trophic system through multiple levels and pathways including direct consumption, excretion, decomposition, and primary production. Direct consumption may occur in the form of predation, parasitism, or scavenging on the live spawner, carcass, egg or fry life stages. Carcass decomposition and the particulate and dissolved organic matter released by spawning fish (e.g., eggs and milt, excrement) delivers nutrients to primary producers. Potential nutrient or energy pathways and factors influencing biomass cycling of spawning salmon is graphically depicted in Figure 15.

Freshwater and estuarine ecosystem productivity depends upon nutrient inputs and retention. Larkin and Slaney²⁶⁶ and Munn et. al.³³⁹ discuss nutrient cycling and the nutrient spiraling concept, whereby nutrients spiraling downstream can influence aquatic system functions for considerable distances. Larkin and Slaney²⁶⁶ and Munn et. al.³³⁹ also discuss the importance of instream habitat complexity (wood debris complexes) for increasing productivity by increasing salmon carcass retention; citing Cederholm and Peterson⁹⁰ and Cederholm et. al.⁹¹. Retention of nutrients also occurs at smaller scales and through chemical and physical processes. Organic molecules in water are rapidly absorbed onto the biofilm that covers most aquatic surfaces⁴⁷. Transport of salmonid organic matter and nutrients across mosaics of inchannel, riparian, floodplain, and estuarine habitats in a watershed may then occur as water, sediments, and organic debris are redistributed; as in freshets. The discussion that follows emphasizes the necessity of retention mechanisms and how the physical and biological complexity of the aquatic, riparian, and wetland zones enhances this function.

Sportsmen and naturalists have long recognized the importance of salmon runs to the natural economy of streams, as this quotation by Haig-Brown¹⁸⁰ reveals:

“The death of a salmon is a strange and wonderful thing, a great gesture of abundance. Yet the dying salmon are not wasted. A whole natural economy is built on their bodies. Bald eagles wait in the trees, bears hunt in the shallows and along the banks, mink and marten and coons come nightly to the feast. All through the winter mallards and mergansers feed in the eddies, and in freshet time, the herring gulls come in to plunge down on the swifter water and pick up the rotting drift. Caddis larvae and other carnivorous insects crawl over the carcasses that are caught in the bottoms of the pools or against the rocks in the eddies. The stream builds its fertility on this death and readies itself to support a new generation of salmon.”




Figure 15. Factors of stream complexity and marine derived nutrient pathways that influence biological activity⁹².

The scientific community has also recognized the contribution of nutrients and organic matter from spawning salmon for some time. Juday et al.²⁴¹ estimated that sockeye salmon transported in excess of 2 million kg of organic matter and 5000 kg of phosphorus to the Karluk River system in Alaska in an average year. This recognition resulted in sockeye lake fertilization programs in British Columbia and Alaska, to replenish lost nutrients caused by fish harvest^{497, 496}. Over the last 10 years stable isotope analysis has enabled direct measurement of marine-derived nutrients in stream^{248, 47, 239} and lake²⁴⁹ ecosystems. These studies have firmly established the need to consider the importance of salmon biomass as a flow of energy and nutrients into the freshwater and estuarine food webs of the Pacific Northwest.

Nutrient levels in fresh water and estuarine systems can be substantially enriched by the organic inputs of spawning runs. The majority of material transported to freshwater by some species of anadromous salmon is of marine origin. Mathisen et al.²⁹⁸ demonstrated that over 95% of the body mass of some salmon species is produced during ocean residence; the remainder represents mass accumulated in freshwater prior to migration to

the sea. The species of salmon that spawn at high densities (e.g., chum, pink, sockeye) significantly alter nutrient loadings and budgets in the freshwater systems where they spawn. In Kamchatka, Krokhin ²⁶² reported that 35-40% of the yearly total phosphorus input to a lake was transported by spawning sockeye salmon, as was much of the nitrogen input to the system. Similar results have been reported for the Iliamna Lake system in Alaska ^{122, 297, 298, 249} and for the Paratunka River basin, Kamchatka ^{261, 262}. Nutrients from spawning pink and chum salmon have been shown to not only enrich the freshwater and estuarine habitats where they spawn, but also the estuarine habitats downstream ^{62, 503}. Munn et. al. ³³⁹ consider changes in nutrient loading and cycling and ecosystem productivity that could result from restoration of historic salmonid populations to the Elwha River system in Washington state. The study indicates a potential 65-fold increase in nitrogen and phosphorus loadings from salmon returns. They concluded that restoration of the Elwha River system salmon runs would have a profound effect on the primary and secondary production in the system.

Table 7 shows calculations of the annual input of organic matter, nitrogen and phosphorus to freshwater catchments within the Puget Sound Basin of Washington, resulting from recent peak spawning escapement levels. These overall inputs of 189 metric tons (mt) of nitrogen and 22 mt of phosphorus equate to approximately 1.9 and 1.2 percent, respectively, for the 10,000 mt and 1,909 mt recent annual outputs of nitrogen and phosphorus, respectively, from freshwater as calculated by Inkpen and Embrey ²²⁶. While the percentages in the table indicate that current salmon escapements are generally a minor source of nitrogen and phosphorus to freshwater habitats within the basin, further examination of information on nutrient loading sources, including salmon runs, indicates that pre-commercial exploitation era escapements were a significant component of the nutrient budgets for the freshwater habitats accessible to anadromous salmon.

Available data on calculated present nutrient loading and projected salmon return loads for a restored Elwha River system were used in comparison with the above values for Puget Sound to illustrate the likely order of magnitude of the difference between the pre-European settlement inputs and present conditions. Annual nutrient inputs of 29.8 mt of nitrogen and 3.5 mt of phosphorus are projected for a 980 mt biomass of restored salmon returns based on Munn et. al. ³³⁹. The nutrient levels are approximately 40% and 13%, respectively, of the present 74.5 mt of nitrogen and 27.3 mt of phosphorus loads to the Elwha watershed ²²⁶. The Elwha River ratios of salmon origin nutrients to total nutrients are respectively 21 and 11 times the percentages of 1.9 and 1.2 for Puget Sound. This is in part due to human induced enrichment of some Puget Sound systems from agricultural and urban land uses several times those in the relatively undisturbed Elwha River system. These preliminary estimates as to the levels of nutrient loadings to the water column and sediments, while crude, should be sufficient cause to trigger some rethinking of what constitutes healthy baseline water and sediment quality for Puget Sound and Pacific Coast streams and estuaries.

Productivity of freshwater ecosystems may be substantially affected by the nutrient and organic matter contributions of spawning salmon. In Lake Dalnee, Kamchatka, chronically low returns of sockeye salmon over a period of several years brought about: (1) a decrease in annual primary production of 20 percent; (2) a 30 percent decrease in total annual production of zooplankton; and (3) a decrease in total annual production of plankton-eating fish (including juvenile sockeye) of 45 percent ²⁶². The number of returning sockeye salmon can alter the productivity of lake ecosystems ^{241, 297, 249}. Richey et al. ⁴²⁵ found that kokanee salmon (*O. nerka*) carcasses added 44.6 kg of phosphorus to a small tributary of Lake Tahoe, California, raising the phosphate concentration of the water by 4-6 µg/L. Algal productivity was stimulated as a result of the increased availability of nutrients.

Table 7. A first order approximation of body biomass and nitrogen and phosphorus imports in metric tons (mt) to the freshwater catchments within the Puget Sound Basin resulting from recent peak wild spawning salmon escapements.

Watershed	Species							
	Chinook	Chum	Coho	Pink	Sockeye	Totals	Nitrogen	Phosphorus
Nooksack	0	334	20	294	0	648	19.7	2.3
Samish	0	25	35	n/a	n/a	60	1.8	0.2
NS Independents	0	0	0	0	0	0	0.0	0.0
Skagit	127	797	26	214	0	1,164	35.4	4.2
Stillaguamish	8	453	13	38	0	512	15.6	1.8
Snohomish	36	413	53	150	0	651	19.8	2.3
Lake Washington	48	0	0	0	215	263	8.0	0.9
Duwamish/Green	74	8	2	0	0	85	2.6	0.3
Puyallup	0	8	19	3	0	30	0.9	0.1
Nisqually	16	349	2	1	0	367	11.2	1.3
South Sound	190	1,345	6	0	0	1,541	46.8	5.5
Hood Canal	27	791	3	13	0	834	25.3	3.0
Strait of Juan de Fuca	42	24	2	4	0	73	2.2	0.3
Totals	567	4,548	180	717	215	6,227	189.3	22.4

The loadings in metric tons were derived by multiplying recent escapements for each species by corresponding average weights and body composition proportions for nitrogen and phosphorus. Escapements data are from Washington Department of Fisheries, Washington Department of Fish and Wildlife, and Western Treaty Indian Tribes^{542, 539}. Average sizes of salmon by species are from Ricker⁴²⁶ as quoted in Larkin and Slaney²⁶⁶. Proportions of body composition for nitrogen and phosphorus are from Larkin and Slaney²⁶⁶. This table is a furthering of the concepts discussed in Lichatowich.²⁷⁷ This is an initial approximation of nutrient imports, more complete data may be available from multiple fish management agencies i.e., Washington Department of Fish and Wildlife, Western Washington Treaty Indian Tribes, Oregon Department of Fish and Wildlife, etc.

Even low spawning densities can provide significant contributions of nutrients to the system. Juvenile coho, steelhead and cutthroat in a small stream in western Washington obtained from 25% to 40% of their N and C from dead coho salmon that spawned in the stream⁴⁷. Aquatic insects also contained high levels of marine-derived N and C, and the foliage of plants growing along the streams also contained nitrogen of marine origin⁴⁷. The direct feeding of salmon fry on salmon carcasses has been known for a long time in the Amur River of Asia³⁷⁷, however, the growth benefits for juvenile salmon has only recently been documented in North America⁴⁷.

Nitrogen and carbon contained in the nonanadromous rainbow trout residing in a southeast Alaska stream was derived almost entirely from the large numbers of pink salmon which spawned at the site ²⁴⁸. Growth rate, condition factor and population density of juvenile coho and steelhead increased dramatically after addition of coho salmon carcasses to two small streams in southwestern Washington ⁴⁸. Additional work by Michael ³¹⁸ and Adams ¹ indicate that juvenile coho spawned in year N reap the benefits provided by all species of salmon spawning in year N+1. Johnston et al. ²³⁹ found a relationship between the proportion of marine-derived nitrogen in insects and the density of spawning sockeye salmon in tributaries of the Stuart River in interior British Columbia. Therefore, the productivity of freshwater habitats may be influenced by the abundance of spawning fish using the system. Also Shuldt and Hershey ⁴⁶² showed direct effects of salmon carcasses decomposition on elevating periphyton accrual and dissolved nutrients in Lake Superior tributaries.

Macroinvertebrate communities in streams receiving salmon runs can change in response to spawning activity and nutrient enrichment. In a Snoqualmie River, Washington tributary and in Kennedy Creek, Washington, Minakawa ³²⁴ found the presence of salmon carcasses and eggs produced a two-fold or greater increase in total insect densities and biomass compared to control reaches. Piorkowski ⁴⁰⁴ found insect taxa richness and diversity to increase in response to nutrient enrichment from salmon carcasses in southeast Alaska, and suggested that insect colonization of carcasses facilitated decomposition and subsequent nutrient release. Bilby et al. ⁴⁷ found all functional feeding groups except insect shredders to be enriched with marine origin isotopes of nitrogen and carbon in western Washington streams after coho salmon spawning. Some aquatic invertebrates such as stoneflies (Plecoptera: Alloperla) ³⁷⁵ and Dipteran flies (Chironomidae) ¹³¹ will scavenge for dead salmon eggs and alevins within the gravel. Limnephilid caddisfly larvae are attracted to recently expired salmon and have been observed feeding directly on fish flesh ^{404, 324}.

Terrestrial insects including fly maggots (Diptera) have also been observed feeding heavily on salmon carcasses in streams in the Queen Charlotte Islands of British Columbia ⁴²², but generally little work has been done to systematically document these activities. Maggot larvae have commonly been observed consuming beached salmon carcasses during the warmer months of the spawning season along the spawning reaches of several Washington streams. Dead chum salmon along Kennedy Creek in South Puget Sound often have their heads filled with maggots (Cederholm, personal observation). Chinook carcasses along Puget Sound Basin rivers can be reduced to skeletal remains by maggots within a two week period; fall freshets frequently have been observed to wash the carcasses and masses of larvae back into the stream where they are then available as food for juvenile salmon and other organisms (Graeber, personal observation). Hornets have also been observed to feed on carcass remains during warm fall weather periods in the same areas; they are especially attracted to exposed fresh flesh or blood (Graeber, personal observation).

Quantitative measurements of salmon carcass consumption in the terrestrial environment has focused on their utilization by high profile species like Bald eagles along the Skagit River, Washington ^{490, 187} and grizzly bears along the Columbia River ²⁰⁵. But Cederholm et al. ⁹¹ recorded 43 taxa of mammals and birds present on small Olympic Peninsula streams at a time when coho salmon carcasses were present, and found that 51% of those taxa had fed on carcasses. Skagen et al. ⁴⁷⁵ in their study of human disturbance on an avian scavenging guild, observed significant bird scavenging of chum and coho salmon carcasses along the North Fork of the Nooksack River, Washington. The primary bird scavengers were eagles, crows, and glaucous-winged gulls. Additional information on observations of wildlife predation and scavenging on salmon is presented in Appendix VII.



Plate #20. Fly maggots eating a chum salmon carcass at Kennedy Creek, Washington. (Photo by: Jeff Cederholm).

Cederholm et al.⁹¹ also reported that black bears, raccoons, and river otters increase food availability for terrestrial species incapable of removing carcasses from the stream. The larger animals rarely completely consumed the carcasses they removed from the stream, and were often followed by an array of other smaller birds and animals who fed on the “leavings”. A similar interaction occurred at McDonald Creek, Glacier National Park, Montana, where kokanee salmon captured by grizzly bears were incompletely consumed, leaving remains for birds and small mammals⁴⁸⁶.

As the above studies indicate, spawning salmon provide a source of carbon, nitrogen and phosphorus essential to maintaining the production of salmon juveniles and other trophic levels of the stream. Accumulating evidence suggests that spawning salmon populations are an important link to the adjacent riparian and terrestrial communities, and indeed, fortifies the role of salmon as a keystone species, wherein the integrity and persistence of the entire community is contingent upon the population’s actions and abundance⁵⁵⁹.

Pacific Salmon Provide Key Ecological Functions

The key ecological functions that spawning salmon play within certain freshwater ecosystems may be illustrated with a well-documented case study from McDonald Creek in Glacier National Park, Montana. This stream is a principal spawning tributary for the Flathead Lake/Flathead River ecosystem. The triggering event in the series of changes that cascaded through this ecosystem was caused by the introduction of an exotic species, the opossum shrimp (*Mysis relicta*), from 1968 and 1975⁴⁸⁶. The shrimp were added to the lake as a food source for kokanee salmon, but behavioral patterns made them unavailable for consumption. Opossum shrimp are voracious predators of zooplankton, the principal food of the kokanee. The shrimp decimated the zooplankton in the lake and by the late 1980s the lake and McDonald Creek spawning kokanee population had collapsed. These fish served as an important food source for various birds and mammals that had fed upon them in the spawning tributaries. One of the most prominent predator and scavenger utilizing this resource were bald eagles that gathered by the hundreds during the kokanee spawning period. In 1981, spawning kokanee in excess of 100,000, attracted 639 eagles, the densest eagle concentration south of Canada. Beginning in 1987 eagle numbers declined along with the kokanee, reaching a low of just 25 birds in 1989. It is feared that loss of the kokanee spawning

run could lead to higher eagle mortality during migration or during winter unless the birds can find alternate food resources, a prospect that is not likely in that ecosystem⁴⁸⁶. A number of other bird and mammal species that used the McDonald Creek kokanee also have been displaced⁴⁸⁶. Gulls, mergansers and mallards commonly fed on kokanee carcasses, while Barrows and common goldeneyes and dippers fed on loose eggs. Mammals that fed on spawning kokanee or carcasses along McDonald Creek, including grizzly bears, coyotes, mink, and river otters, are now less common along the creek.

Estuaries, where rivers and streams meet tidal influence and enter the ocean, act as traps for sediments, organic materials, and nutrients washed from watersheds. Some species of Pacific salmon typically spawn near saltwater or even beginning within the upper reaches of estuaries and often spawn at very high densities. The effect of salmon carcasses on the nutrient dynamics and trophic productivity of estuarine systems is just beginning to be examined. Kline et al.²⁴⁸ reported that approximately 30,000 pink salmon spawned within 1.2 km of the estuary of Sashin Creek in southeast Alaska. In southwestern Washington, the 5 km of Kennedy Creek accessible to anadromous fishes has supported as many as 80,000 spawning chum salmon (WDFW unpub. data). Using the size and body composition information previously cited (under the Elwha River discussion), we estimate that this peak escapement to Kennedy Creek delivered approximately 398 mt of salmon flesh containing 12 mt of nitrogen and 1.4 mt of phosphorus to 0.075 km² of stream channel area (5 km with an average channel width of 15 m). The nutrient loading per unit of channel area would be 160 mt/km² nitrogen and 18.7 mt/km² phosphorus. The salmon carcass materials, or their nutrients, in dissolved phase or sorbed to sediment particles may be carried to the estuary through various physical, biological, and chemical processes^{47, 266, 339}. Therefore, a nutrient link may function between adult salmon carcasses and juvenile salmon rearing in the estuary. For example, Fujiwara and Highsmith¹⁵⁹ found elevated stable isotope ratios of nitrogen in *Ulva* sp., an estuarine macroalga, following the decomposition of salmon carcasses in Seldovia Bay, Alaska. *Ulva* sp. are a major food source for harpacticoid copepods, which in turn are a preferred prey of juvenile chum salmon fry in the estuary. Thus, the contribution of nutrients and organic matter from salmon carcasses may be a substantial source in some systems and may be a key factor in promoting estuarine productivity. The importance of estuaries as nursery zones for anadromous salmon along the Pacific Northwest coast is well documented^{196, 355, 289, 397, 454, 194}, the role carcasses play in maintaining productivity of these systems may be critical in supporting the health of salmon populations¹⁵⁹.

The role salmon populations play as a key vector in the recycling of energy and nutrients inland from the North Pacific Ocean to aquatic and terrestrial ecosystems is now gaining recognition as a critical component of ecosystem function^{350, 348, 463, 464}. River and lake fertilization with inorganic nutrients has been undertaken with ecosystem restoration in mind in some British Columbia systems^{476, 496, 16, 15}, however, artificially supplementing inorganic nutrients may not fully mitigate for the loss of the multiple pathway flow of energy and materials provided by naturally spawning salmon.

Salmon As Vectors In Broader Nutrient Cycling

The flux of nutrients is essential for the continuity and stability of any living system⁴¹², and nutrients provide a link between aquatic and terrestrial ecosystems^{61, 317}. Biological vectors through which materials and energy are transported include migration of animals (i.e., mammals, birds, fish) that carry nutrients across ecosystem boundaries^{61, 379, 463, 464, 349}. Therefore, the role and importance of salmon in the freshwater and terrestrial ecosystems can be recognized within the context of broader nutrient cycling, and spawning migrations of salmon represent an obvious example of this process. Other means of moving nutrients upstream, such as the emergence of the adult stages of insects and meteorological vectors, are considered

to be relatively insignificant compared to anadromous fish²⁷⁸. The nutrient-subsidy contributed by salmon in the North Pacific could potentially serve as a model for a more general, and global, aspect of nutrient circulation. Such studies may be supplemented by an assessment of the extent to which migratory birds, which travel long distances between boreal/subboreal zones and temperate and/or tropical regions, contribute to similar effects.

Using the discipline known as *Resource Physics*, Tsuchida⁵²⁴ has explained how the hydrologic cycle and the convection of air are able to keep the earth in a low-entropy state^{see also 350}. The hydrologic cycle is in turn driven by solar thermal energy and the earth's gravity. Subsequently, inorganic salts (especially nitrates and phosphates), which are essential to formation and activities of animals, plants, and microorganisms, are eventually washed downstream⁵²⁵. Ultimately, these salts are dissolved in river water and transported to the ocean, where they attain the highest and most uniform concentration below the depth of 1,000 m, largely free from biological consumption in the absence of photosynthesis. However, due in part to ocean currents (local upwelling), these nutrients eventually find their way back to the surface waters. This occurs in northern temperate or subpolar oceans by the effective vertical mixing of seawater due to the approximation of water temperature between deep and shallow water, primarily during colder seasons. Finally, uptake of these salts by marine plants near the surface, where photosynthesis is possible during the warmer seasons, allows a means through which other animals are able to derive and transport the nutrients inland. For example, Tsuchida⁵²⁵ speculates that in coastal regions, some birds will carry nutrients back to the land after deriving them from the consumption of marine organisms. Bird excrement is a fertilizer rich in inorganic matter, especially phosphates, as evidenced by the material deposits (Peruvian bird guano) on tropical sea islands. Sibatani^{463, 464} points out that another, arguably more significant way that nutrients are transported back onto the land is by anadromous fish swimming up, spawning, and dying in the many rivers of the Asian and North American continents. Murota and the Faculty of Environmental Studies³⁵¹ discussed how migratory fish move ocean nutrients inland and benefit the Siberian forest and its various wildlife inhabitants (Figure 16).

*“In the Edo era, some people in Japan started to notice that forests along seashores or rivers attracted fish towards them. It was considered that a forest could give benefits to fish in the forms of shadow as shelter, nutrients, and so on. This consideration remained in the minds of people living near waterfronts or forests after the Meiji Restoration (1868). When the first forest act was introduced at the beginning of the twentieth century, it contained the article ordering the conservation of uo-tsuki-rin, which literally meant ‘fish-attaching forest.’ This article is still valid in the present-day forest act of Japan.”*³⁴⁸.

With regard to this *uo-tsuki-rin*, Sibatani^{463 cited in 348}, the Japanese scientist raises an interesting question:

“He thinks that it may be the fish that helps forests to grow rather than forests providing fish with comfortable spaces. This hypothesis comes from his research on the forests in Maritime Territories of Eastern Siberia, specifically along the Ussuri River, a tributary of the Amur River. This area is subject to cold temperatures and receives very little sunshine, but there has been forest growth for a long period of time. Nutrients must have been carried from somewhere. But as there are no significant mountains in the upstream areas, the nutrients must have come from downstream, or more exactly, from the Northern Pacific, and the salmon as well as some other kinds of fish are likely to play a significant role as their carriers.”




Figure 16. Anadromous fish and the Siberian forest³⁵¹.

The theories of Sibatani^{463, 464}, Murota^{348, 349}, and Tsuchida^{524, 525} cause one to reflect on the once bountiful salmon runs of the Columbia River. Before Europeans settled the Pacific Northwest, salmon and steelhead had access to over 20,000 km of main river and tributaries in the Columbia River basin³⁷⁹. The annual Columbia River salmon and steelhead run size has been estimated to range between ten and sixteen million fish³⁷⁹. Applying an average fish weight range (5.51 to 6.17 kg.), compiled as a species composition weighted average from associated fishery data reported in Table 12 of the NPPC report³⁷⁹, to the run size estimates yields annual potential spawning biomass contributions in the Columbia River basin of between 55,140 and 98,706 metric tons (mt). This amounts to 2.51 to 4.93 mt per kilometer of anadromous habitat.

But what of the native people of the Columbia River? Many traveled long distances to partake in the catch and consumption of salmon, and in doing so participated in the further cycling of nutrients over this vast watershed, and beyond. Some tribes of the upper Columbia were known to cross the Continental Divide to trade dried salmon for buffalo hides³⁷⁹, thus providing an additional mechanism for transfer of marine-derived nutrients to the inland land mass. Salmon, wildlife, and humans, therefore, may be the most prominent carriers of ocean nutrients to inland ecosystems.

Salmon are a pervasive influence on the productivity of the otherwise generally oligotrophic ecosystems of coastal freshwater catchments of the entire Pacific rim of North America, including the Pacific Northwest. A coast wide estimate of the baseline biomass comparable to the above Columbia Basin estimate may be useful to further discussions as to the scope of that influence. However, as noted above, run reconstruction is a problematic impediment, both because various impacts preceded any documentation of the resources base and because the available documentation is not consistent in content or time frame. For the above reasons, the following estimate is offered only as an approximation of the historic baseline biomass for the eastern rim of the Northern Pacific, by major subareas. The substantial historical research required to refine these approximations for finer scale applications is left for subsequent efforts.

Peak cannery pack to total run biomass ratios for the Columbia River are used here to estimate the approximate biomass for pre-European settlement era salmon runs. The peak historic commercial salmon cannery pack for the eastern Pacific rim, adapted from Cobb^{99a}, are shown in Table 8 as a starting point. The cannery pack records are a well documented, but very minimal, representation of baseline run size and biomass. Expansion factors for the estimate are calculated from the above noted Columbia River historic run biomass estimate range (55,140 to 98,706 mt.), divided by the peak Columbia River cannery pack weight (13,819 mt.). The expansion factors (approximately 4.0 and 7.1 times the peak cannery pack) were then applied to the cannery pack weight data (column 3) to generate the biomass estimates reported in Table 8 (columns 4 and 5). The resulting coast wide total run biomass range estimate is 1.2 to 2.1 million metric tons.

Maintaining the Salmon Link In Nutrient Cycling

As the decline of wild salmon continues throughout the Pacific Northwest³⁶⁷, it is logical to assume that the productivity of some freshwater and terrestrial ecosystems will also decline. Decreased production could be self-perpetuating, as salmon stocks already in decline are likely to decrease further in a negative feedback loop^{47,266,339}. The impending listing of many salmon stocks as endangered or threatened has forced federal and state government agencies to take aggressive action for salmon protection and recovery^{366,544}.

There are numerous implications for fisheries management and the stream and riparian ecosystems that salmon inhabit. An obvious scenario includes allowing sufficient salmon to spawn in as many streams that were historically used by salmon as possible. A diversity of species that utilize several different stream orders will distribute marine nutrients throughout the entire watershed. Larkin and Slaney²⁶⁶ have made a case for the need to consider nutrients loading levels and distribution in harvest management and hatchery production planning to sustain stream productivity for salmon. Munn et al.³³⁹ have provided an estimate on the expected increase in nutrient levels and system productivity resulting from fully restored salmonid escapements under the proposed Elwha River restoration. Resource managers are beginning to consider the implications of changes in the flows of salmonid organic matter and nutrients may have on the levels of sustainable fish and wildlife management^{339,266}.

Numerous implications for water quality management exist within the native range of the Pacific salmon. The keystone role that salmon play in nutrient cycling needs to be recognized as an essential component to background water quality levels for healthy watersheds. The water quality regime of the high biodiversity of inputs from human activities. We will need to learn more about the range of historic nutrient loadings to freshwater and estuarine systems to understand how the declines in salmon runs and how disruptions to the sub-ecosystems of the salmon may be altering the biotic communities. From there we can better answer questions on appropriate allowable loadings and the water quality standards that will best sustain healthy communities.

Table 8. Peak Cannery Packs and Estimated Total Run Biomass for all Species of Pacific Salmon in the Northwest and North American Pacific Coast Prior to 1930

Area	Peak Year	Cannery Pack ^{1/}		Total Run Biomass Estimates ^{3/}	
		Number of Cases	Metric Tons ^{2/}	~4.0 x Cannery Pack Metric Tons (x 1,000)	~7.1 x Cannery Pack Metric Tons (x1,000)
Noyo River	1919	7,500	163	1	1
Sacramento River	1882	200,000	4,354	17	31
Eel River	1883	15,000	327	1	2
Klamath River	1912	18,000	392	2	3
Smith River	1925	7,700	168	1	1
Coastal Oregon rivers	1911	138,146	3,008	12	21
Columbia Basin	1895	634,696	13,819	55	99
Willapa Harbor	1902	39,492	860	3	6
Grays Harbor	1911	75,941	1,653	7	12
Coastal WA Rivers	1915	31,735	691	3	5
Puget Sound	1913	2,583,463	56,248	224	402
Fraser River	1901	998,913	21,749	87	155
Outlying Districts	1928	1,265,522	27,553	110	197
Rivers Inlet	1925	197,087	4,291	17	31
Skeena River	1922	482,305	10,501	42	75
Nass River	1918	143,908	3,133	13	22
Alaska	1926	6,652,882	144,849	578	1,035
Total		13,486,994	293,760	1,172	2,098

1. Data on number of 48 pound cases of canned salmon packed is adapted from Cobb ^{99a}.
2. Weight of the 48 pound cases was converted to metric tons using the conversion factor of 0.45359237 kilograms per pound currently recognized by the American National Standards Institute (IEEE/ASTM SI 10-1997) ^{xxx}.
3. Total run biomass estimates were calculated for the major areas reported by Cobb ^{99a}, using expansion factors calculated from historic run size and associated fish weight data reported for the Columbia River by the Northwest Power Planning Council³⁷⁹. Expansion factors of approximately 4.0 and 7.1 were calculated as the lower and upper ends of the historic Columbia River run biomass (55,140 to 98,706 mt.) estimated in this paper, divided by the peak Columbia River cannery pack weight (13,819 mt.), as adapted from Cobb ^{99a}.

In order to ensure nutrient cycles from the ocean back to the watersheds, the major vector of this process, wild anadromous salmon, must recover from their current status. Identifying and securing channels for the recycling of nutrients is an important component for maintaining biological diversity, at least in the Northern Pacific^{463, 464, 348, 349}. The key to the sustainability of the human economy also may lie in these material cycles to some degree, as our economy relies heavily on healthy ecosystems to sustain the production of food and other resources^{350, 349, 525}. The importance of the nutrient cycling link provided by anadromous salmon, therefore, illustrates the need for an uncompromising and all-encompassing plan to protect and recover wild salmon populations before the system is unrecoverable.



Plate #21. Jeff Cederholm holding a chum salmon at Kennedy Creek. (Photographer unknown).

WITH AN ECOSYSTEM PERSPECTIVE IN MIND - WHERE DO WE GO FROM HERE?

The need for an ecosystem approach to salmon management has inevitably grown in Washington and Oregon ^{365, 499, 305}. Terrestrial ecologists have recognized the influence human uses (and thereby disturbances) have had on terrestrial and aquatic ecosystems of large scales ^{295, 151, 545}. Some investigators have explored the significance of the salmon as a key ecological process vector on the broadest scales of energy and nutrient transport ^{559, 464, 348}. The use of salmon as an indicator of ecosystem health, or “the canary” is complicated by the fact that this canary is a food resource in high demand by humans ⁴⁶⁴.

The magnitude of the role of salmon populations as keystone vectors in energy and nutrient cycling inland from the Northern Pacific to freshwater and terrestrial ecosystems is now gaining recognition as a critical component to an overall understanding of ecosystem functions ^{464, 348, 349, 92}. Application of this knowledge to understand the cumulative impacts of human land use practices and fisheries exploitation on ecosystem functions is only just beginning. New tools are necessary to make management actions toward regaining lost productivity and biodiversity objectives ^{499, 305, 365}.

Greater understanding of the hydrologic cycle has helped us to improve land and water uses to better adapt to our environment. Better understanding of geologic processes including the role of hydrogeology in shaping the landscape and controlling the rates at which sediments, and small and large organic debris cycle through watersheds is also leading to changes in views on land and water uses ^{434, 295}.

Understanding nutrient cycling processes, pathways and the effects of nutrient loading has helped in managing water and sediment quality problems ^{435, 538, 245}. In much the same manner, an understanding of nutrient spiraling and cycling in streams ³⁶⁹ and the keystone role of anadromous salmon ^{266, 339} will be valuable, if not essential, to understand how we have affected coastal ecosystems and the processes that support them. Such an understanding will lead us to better identification of those management action options we may take to achieve desired future conditions.

Maser et al. ²⁹⁵ have added substantially to the literature on downstream and seaward movement of materials, energy and nutrient transport processes to and through aquatic systems. Their report compiled and presented a wealth of information on the inputs, fates, and effects of forest debris, particularly LWD, in freshwater, estuarine, and marine ecosystems. In essence, physical and chemical processes were well described, now the biological is considered. Adding estimates of the upstream flow of energy and nutrients via salmon to existing watershed processes literature, will provide a more complete picture of the large scale and long-term energy and nutrient cycles for entire watersheds. An understanding of the overall cycles and levels of productivity at this scale will provide the context for interpretation of local level trends in production and materials transport, utilization and storage. It has been well established that aquatic systems have metabolisms that function based upon physical processes, rates of loadings of materials, energy, and nutrients, rates of primary and secondary production, and resulting changes in standing stocks of fishes.

As a keystone species to the productivity and biodiversity of the ecosystems of the North Pacific basin, anadromous salmon closely link the management issues of the forests, the floodplains and

lowlands, the estuaries and nearshore areas, and the ocean domains as a continuum. Materials, energy and nutrient budget analyses on an appropriate time scale will be necessary to estimate the potential effects that past land use practices have had on production and discern which ones persist. Budgets for the North Pacific Basin can theoretically be calculated in a similar manner to that already used on smaller systems. Then we can begin to use resulting information to provide a context for management decisions in various disciplines and forums that will affect materials, energy, and nutrient flows and stocks at various scales. To address the disconnects apparent in the current state of terrestrial and aquatic systems will require some estimate of many factors, including:

1. What is the status of the nutrient capital and rates of transport within the domains and the basin as a whole (nutrient budget)?
2. What is the range of the nutrient and materials capital and rates of transport (how does the current budget relate to the known ranges of standing stocks and rates of metabolism and transport)?
3. How have humans altered the nutrient budget?
4. What adaptive management actions might be warranted and feasible to push the terrestrial and aquatic systems toward the identified goals?
5. Are there some measures to employ in the interior until stocks of salmon can be restored?
6. What are the desired future conditions?

Early European settlement of the eastern North Pacific Rim territories provides many accounts of heavy extractions of forest resources^{295, 276, 348, 464, 554, 110a, 70, 365}. Logging of the forests had the obvious effect of short circuiting the prior cycles that supplied woody debris to freshwater, estuarine, and marine ecosystems. In recent decades mechanized logging equipment combined with highly efficient (“clean”) logging practices and slash burning to prevent wildfires and accelerate re-growth of planted conifers has further resulted in very little debris left on the site or entering streams. The store of vast quantities of nutrients in the form of the decayed woody debris and trapped detritus that serves as substrate for long-term nitrogen fixation and retention no longer exists. Large woody debris may continue to enter stream corridors, but not necessarily in the amount, size, and quality that it did in the past, thereby decreasing potential to provide stream structure and organic matter to food-webs^{416, 299}.

The long-term loss of the function of LWD as a primary component of the floodplain waterways, resulting from land and channel clearing, may well be more significant than the loss of the wood material itself. The resistance of the abundant woody material slowed the flows of water, sediment, and smaller debris; resulting in very complex valley floor stream-ways composed of multiple highly sinuous channels that were generally well connected to off-channel wetland systems by sloughs and high water channels. Put this liquidation of natural resource capital into the context of long-term climate cycles¹⁸⁹ and Maser et al.’s²⁹⁵ long-term geologic and successional cycles, and one can begin to formulate management goals for ecological processes.

A combination of development activities have diverted water, shortened, straightened, cleared, dammed, diked, drained, filled and polluted the habitats of salmon. Early in settlement, logging and splash damming, land clearing for agriculture, and channel clearing for navigation appear to have had the most pronounced effects. The development and consumptive utilization of natural resources was highly dependent upon water-borne transportation and patterns of impacts are reflective of navigation-centered commerce. Continuing development for agriculture, industry, and urban growth has resulted in further losses through conversions to other uses. Impacts have become more pervasive throughout the landscape as

transportation infrastructure and vehicle capabilities have increased. Releases of persistent toxins has contaminated coastal sediments. Thus available freshwater and estuarine salmon habitats also continue to be degraded by ongoing land uses ^{365, 295, 454, 174, 526, 554, 245}.

Desired Future Condition

The inability of the various interest groups to resolve conflicting and agreed-to goals or conditions has been identified as the fatal gap in salmon management ^{365, 54}. Describing the desired future condition of the terrestrial and aquatic ecosystems and identifying and defining human actions can influence movement of ecosystems toward those conditions. The institutional changes suggested by Lichatowich ²⁷⁷ will be necessary to implement the long-term management approach required. Some of those institutions are known and have been in use during other eras where human land and resource uses appeared to have been indefinitely sustainable ^{464, 554}.

In spite of the high potential commodity value of the harvest and our knowledge that the Pacific salmon is a key driver for the biodiversity and productivity of the northern Pacific basin, we have not developed the strategies for effective long-term management of the resource's health. The widespread use of salmon hatcheries has also significantly reduced the amount of salmon carcass nutrients available for the aquatic food webs. Modern human culture has not fully adapted to the environment of the northern Pacific Basin. The environment of our region is showing the signs of stress all around us which indicates the failure of our past and current approaches. If humans are to thrive at present population levels within portions of the basin, we will need to look more at multiple scales of ecosystem management and at integrating that management to sustain productivity over very long time frames.

The challenge, then, for the people of Washington and Oregon and the whole North Pacific Rim is to recognize that the character and the health of the northern Pacific Basin sub-ecosystems depend upon the material and energy flow processes that link them. The flows of energy and of organic and inorganic materials among the various aquatic and upland ecosystems determines the productivity of each component and of the whole. In short, the whole is greater than the sum of the parts, symbiosis on a grand scale. Understanding the biological processes and the impacts of our management practices upon them will be necessary for our long-term (measured in generations) success in the region. To do that we will need to look beyond the plants, animals, and the habitats of a given smaller scale ecosystem to the processes that link them together, perhaps into the large scale ecosystem of the anadromous salmon. The flow of sediments, woody debris, detritus, and nutrients through a watershed determines the character and productivity of the entire watershed, estuary, the near shore zone and even the domains of the northern Pacific Ocean. The flow of energy and nutrients back upstream via the Pacific salmon and the ability of the watershed to retain them, in large measure, determines the productivity of the entire watershed.

IMPLICATIONS FOR FISHERIES MANAGEMENT

Harvest management of anadromous salmon stocks along the Pacific coast has generally been governed by strictly density dependent policies, in which escapement of spawning fish is maintained at the level which is predicted to generate the greatest number of harvestable fish. This management approach is referred to as maximum sustained yield (MSY). The MSY level is estimated from a relationship between spawning fish and the number of recruits (offspring in the next generation) they produce.

Two models are most commonly employed to describe the relationship between spawning escapement and the population of recruits produced in the next generation for the stock being managed: the Ricker Model^{425a} (Figure 17-A) and the Beverton-Holt Model^{41a} (Figure 17-B). Both these models assume that the shape of the spawner-recruit relationship is determined primarily by density-dependent interactions. The Ricker model is derived by assuming that the mortality rate of eggs and juveniles is proportional to the initial cohort size and the Beverton-Holt model by assuming that the mortality rate is linearly dependent on the number of fish alive in the cohort at the time. The Ricker model is most often applied to species that compete for spawning space in streams, and the Beverton-Holt model to species that compete for rearing space or food in streams. However, exceptions to this general rule commonly have been found. Chilcote^{97a} found that 26 steelhead populations in Oregon, a species for which the Beverton-Holt model would typically be selected as most appropriate because steelhead juveniles compete for rearing space and food in




Figure 17. Spawner-recruit curves: (A) Ricker model, (B) Beverton-Holt model.

streams, were better fit by a Ricker model. In cases where variable environmental conditions from generation to generation or other factors not related to density-dependent interactions have a major effect on survival; spawner-recruit data may not be adequately described by any model. Because so many physical and biological processes are averaged across the life cycle from spawner to recruit, Hilborn and Walters ^{204a} have cautioned that it is better to think of spawner-recruit curves as general statistical descriptions rather than something determined from any fundamental biological principle.

Regardless of which model is selected, a so-called “replacement line” with a slope = 1 (i.e., one recruit per spawner) is drawn from the origin to intersect and extend beyond the curve. Theoretically, for any level of spawning where the predicted number of recruits is greater than the number needed for spawning, the “surplus” recruits are available for harvest. At the high levels of spawning, the number of recruits produced is less than the spawners required to maintain the population at that level and there are no surplus recruits. The MSY level is defined as the point where the difference between the spawner-recruit curve and the replacement line is greatest, (labeled B and C on the curves in Figures 17-A and 17-B) thus providing the maximum number of harvestable fish. The point where the replacement line intersects the spawner-recruit curve (labeled A on the curves in Figures 17-A and 17-B) represents the equilibrium population level for the stock. Given no harvest of fish and a system with stable environmental conditions over time, the population would migrate towards this point where the number of returning recruits would just replace the number of spawners that produced them. One might think of point A as the natural carrying capacity of the undisturbed system for returning adults.

Although the concept of managing a stock of salmon based on MSY is very appealing, there have been many serious problems encountered in attempting to apply this theory. There are often very high levels of variability in spawner-recruit data. This variability makes selection of the appropriate model to fit the data uncertain making the estimation of key model parameters problematic. As these model parameters are key determinants in the estimation of the MSY level, the setting of escapement goals for stock maintenance based on these models is very risky indeed.

There are well-developed statistical methods that attempt to deal with the variability in spawner-recruit data sets. These procedures can lead to more risk-averse management decisions when using spawner-recruit models (see for example ^{286a, 514a, 424a}). Even so, the simplifying assumptions underlying the MSY approach cause it to be fraught with other pitfalls. If those making management decisions do not recognize these deficiencies, the true condition of a stock may not be understood until the population suffers a precipitous decline.

Perhaps the greatest source of error in spawner-recruit model estimates is caused by the assumption that density-dependent factors are primarily responsible for determining survival (i.e., number of recruits per spawner). There is ample evidence that environmental factors have a substantial effect on survival of salmon. In the freshwater environment survival may be affected by floods, droughts or declining habitat quality caused by numerous anthropogenic influences ^{365, 91a}. There is growing evidence that productivity of the northern Pacific Ocean varies cyclically ^{32, 189}, greatly influencing mortality rates in the marine environment ^{537a}. Thus, controls on mortality rate may be greatly altered by numerous factors that vary in their intensity over time and the effects of which may not be density-dependent.

Considering the above discussion research has revealed that organic matter and nutrients transported to streams by spawning salmon are important for maintaining the productivity of these systems. Nitrogen and

carbon contained in rainbow trout residing in a southeast Alaska stream was derived almost entirely from the large numbers of pink salmon which spawned at the site ²⁴⁸. Juvenile coho salmon, cutthroat trout and steelhead in a tributary of the Snoqualmie River, Washington, obtained as much as 40% of the carbon and nitrogen in their muscle tissue from the carcasses of coho salmon⁴⁷. Johnston et al. ²³⁹ found a relationship between the proportion of marine-derived nitrogen in insects and the density of spawning sockeye salmon in tributaries of the Stuart River in interior British Columbia. Therefore, the productivity of freshwater habitats may be influenced by the abundance of spawning fish using the system.

All age classes and species of fish in the stream utilize the materials transported by spawning salmon ^{47, 48}. For example, juvenile coho spawned in year N reap the benefits provided by all species of salmon spawning in year N+1 ^{318, 1}. Conventional spawner-recruit models cannot capture this relationship.

Managing at the MSY level may have the effect of substantially reducing the delivery of marine-derived nutrients to freshwater habitats. Let us assume that a coho population is undisturbed (no harvest) and at equilibrium, and, since coho juveniles compete for rearing space and food in streams, can be modeled by the Beverton-Holt ^{41a} relationship shown in Figure 18-A. Since this population is undisturbed and at equilibrium, reading off the spawner axis below point A gives the number of spawners whose carcasses would decompose in the stream each cycle to provide nutrients to support the next generation of juveniles. Harvesting this population to the MSY level would reduce the number of spawners, and associated nutrients, to point S on Figure 18-A. This level of spawners represents a reduction of about two-thirds in the number of adult spawners allowed to return to the stream and a corresponding decrease in the level of marine nutrients returned to the stream.

The decreased availability of carcasses has been shown to impact the growth rate of juvenile fishes. Artificially increasing availability of marine-derived materials by adding the carcasses of hatchery-spawned coho salmon to a small stream in southwestern Washington accelerated growth of juvenile coho salmon in this system relative to a nearby stream reach with low availability of carcasses ⁴⁸. Coho at the enriched site grew twice as fast as fish at the site without carcasses and achieved a body size nearly 50% greater by early winter. Body size of juvenile salmon has been positively correlated with overwinter survival in freshwater ^{190a, 414}. Increased smolt size provides a survival advantage in the marine environment ^{49a, 49b, 215a, 537a, 516a, 214a}. Therefore, if harvest of fish causes a reduction in nutrient delivery to the stream sufficient to impact growth, survival will be negatively impacted. This impact will decrease recruitment to the next generation of spawners, further depleting the nutrient capital of the system and potentially further depressing survival. The effect is a progressive downward shift of the stock-recruitment relationship for each successive cycle. This shift is illustrated in Figure 18-B with the adjusted curve labeled NEW. The possibility that the spawner-recruit relationship could be fundamentally altered due to management decisions is never taken into account in setting harvest levels.

If it were only a matter of stream productivity loss, restoring lost stream productivity might eventually reproduce the same run size. But unanticipated over harvest in any cycle is tantamount to reducing the capacity of the system as well, which down-shifts the stock-recruitment relationship in the manner illustrated in Figure 18-C, so that eventual run size also declines.

Another important factor ignored by spawner-recruit models is the loss of stock productivity related to loss of genetic variability. Loss of genetic variability can reduce survival and mean fitness of a population ^{141a}. Geiger et al. ^{165a} reported that there is a link between genetic variability and the level of exploitable




Figure 18. Spawner-recruit curves showing the undisturbed populations (A); the same population with a loss in productivity (B); and the same population with an additional loss in capacity (C).

production in Alaskan pink salmon populations. They showed that a relatively small proportion of the breeding population is the most productive in any one generation and that the genetic composition of this productive segment changes between generations. This process appears to be true for coho salmon as well^{304a}, and Geiger et al.^{165a} suggest that it may be true for salmon in general. As Ricker and Beverton-Holt models only incorporate total population size, and not population-segment specific contributions to recruitment, they cannot predict decreases in overall stock productivity that result from decreased genetic variability. Harvest policies and practices that inadvertently reduce or eliminate small population segments would decrease genetic variability and could impact stock productivity.

There is increased interest in the Pacific Northwest in developing harvest management strategies that address nutrient delivery to freshwater ecosystems^{266, 544}. However, there is relatively little information on which to base escapement targets that will meet this objective. Ideally, estimates of the number of spawners needed to fulfill this function would be determined by experimentally altering escapement levels for each stock and evaluating the impact on system productivity. However, conducting such an experiment on hundreds or thousands of stocks over a sufficient length of time is a daunting prospect and would not provide usable results for many years. Several other approaches to determining appropriate escapement levels are currently being investigated. One option being considered attempts to determine the amount of food required to support a population of rearing fish that fully utilizes the habitat available in a stream. Escapement levels would be established which ensure sufficient nutrients and organic matter are returned to the stream to produce this level of food. Another alternative is to develop a relationship between spawner density and the proportion of marine-derived nutrients in the tissues of juvenile fish. This type of relationship may enable a “saturation level” for marine nutrients to be established and escapement goals set accordingly. However, these approaches do not account for impacts associated with land use that drive down stock productivity (reduce survivals) and reduced habitat capacity, and decreased genetic diversity, nor do they incorporate any consideration of temporal variability in environmental conditions. Nonetheless, these approaches do represent a shift from MSY to more ecologically-based stock management objectives.

LITERATURE CITED

1. Adams, C. 1998. Determining the effects of marine derived nutrients on growth of juvenile coho salmon (*Oncorhynchus kisutch*) using stable isotopes. The 1998 Annual General Meeting of the North Pacific International Chapter of American Fisheries Society Meeting, held March 18-20, 1998 at Alderbrook, WA.
2. Agee, J. K. 1988. Successional dynamics of forest riparian zones, pages 31-43. *In*: K. J. Raedeke, editor. Streamside Management: Riparian Wildlife and Forestry Interactions. Contrib. No. 59, Institute of Forest Resources, University of Washington, Seattle.
3. Ainley, D. G., C. S. Strong, T. M. Penniman and R. J. Boekelheide. 1990. The feeding ecology of Farallon seabirds, pages 51-127. *In*: D. A. Ainley and R. J. Boekelheide, editors. Seabirds of the Farallon Islands, Ecology, Dynamics, and Structure of an Upwelling-System Community. Stanford University Press, Stanford, CA.
4. Ainley, D. G., and D. W. Anderson. 1981. Feeding ecology of marine cormorants in southwestern North America. *Condor* 83:120-131.
5. Akande, M. 1972. The foods of feral mink (*Mustela vison*) in Scotland. *Journal of Zoology*. 167:475-479.
6. Alcorn, G. D. 1958. Nesting of the Caspian Tern in Gray's Harbor, Washington. *Murrelet* 39:19-20.
7. Alexander, G. R. 1979. Predators of fish in coldwater streams, pages 153-170. *In*: Predator-Prey Systems in Fisheries Management. Sport Fishing Institute, Washington DC.
8. Alexander, G. 1977. Food of vertebrate predators on trout waters in north-central lower Michigan. *Michigan Academician* 10:181-195.
9. Allee, B. A. 1982. The role of interspecific competition in the distribution of salmonids in streams, pages 111-122. *In*: E. L. Brannon and E. O. Salo (eds.). Proceedings of the Salmon and Trout Migratory Behavior Symposium. June 3-5, 1981, First International Symposium. School of Fisheries, University of Washington, Seattle.
10. Allen, K. R. 1951. The Horokiwi Stream: a study of a trout population. *New Zealand Dept. Fisheries Bulletin* 10:1-238.
11. Anderson, N. H. 1992. Influence of disturbance on insect communities of Pacific Northwest streams. *Hydrobiologia* 248:79-92.
12. Anderson, M. E. 1977. Aspects of the ecology of two sympatric species of *Thamnophis* and heavy metal accumulation within the species. M. S. Thesis, University of Montana, Missoula.
13. Antonellis, G. A. Jr., and M. A. Perez. 1984. Estimated annual food consumption by northern fur seals in the California current. *CalCOFI Rep.* 25:135-145.
14. Antonelli, A. L., R. A. Nussbaum and S. D. Smith. 1972. Comparative food habits of four species of stream-dwelling vertebrates (*Dicamptodon ensatus*, *D. copei*, *Cottus tenuis*, *Salmo gairdneri*). *Northwestern Sci.* 46:277-289.
15. Ashley, K., L. C. Thompson, D. C. Lasenby, L. McEachern, K. E. Smokorowski, and D. Sebastian. 1997. Restoration of an interior lake ecosystem: the Kootenay Lake fertilization experiment. *Vol. 32: 295-323.*
16. Ashley, K., and P. A. Slaney. 1997. Accelerating recovery of stream, river and pond productivity by low-level nutrient replacement, pages 13-1 to 13-24. *In*: P. A. Slaney and D. Zaldokas, editors. Fish Habitat Rehabilitation Procedures. Watershed Restoration Program. Ministry of Environment, Lands and Parks. Vancouver, British Columbia.
17. Ayles, G. B., J. G. I. Lark, J. Barica and H. Kling. Seasonal mortality of Rainbow trout (*Salmo gairdneri*) planted in small eutrophic lakes of central Canada. *Canadian Journal of Fisheries Research Board.* 33:647-655.

18. Baird, P. H. 1990. Influence of abiotic factors and prey distribution on diet and reproductive success of three seabird species in Alaska (USA). *Ornis Scandinavica* 21:224-235.
19. Baird, R. 1991a. W. The Risso's dolphin in British Columbia. *Victoria Naturalist (Victoria B.C.)* 47:6-7.
20. Baird, R. 1991b. Optimal foraging and intraspecific competition in the tufted puffin. *Condor* 93:503-515.
21. Baker, R. C., F. Wilke and C. H. Baltzo. 1970 The Northern fur seal. United States Dept. of the Interior. U. S. Fish and Wildlife Service. Circular No. 336. Washington D.C.
22. Balcomb, K. C., J. R. Boran and S. L. Heimlich. 1982. Killer whales in Greater Puget Sound. Pages 681-685 in *International Whaling Commission Report of the Commission* 32.
23. Balcomb, K. C., J. R. Boran J. R. Osborne and N. J. Haenel. 1980. Observations of killer whales (*Orcinus orca*) in Greater Puget Sound, State of Washington. INTIS PB80-224728, U. S. Dept. of Comm., Springfield, Virginia.
24. Baltz, D. M., and G. V. Morejohn. 1977. Food habits and niche overlap of seabirds wintering on Monterey Bay, California. *Auk* 94:526-543.
25. Banci, V. 1987. Ecology and Behaviour of Wolverine in Yukon. M.S. Thesis. Simon Fraser University. 178 pp.
26. Banfield, A. W. F. 1974. The mammals of Canada. University of Toronto Press. Toronto. 438 pp.
27. Bayer, R. D. 1986a. Seabirds near an Oregon estuarine salmon hatchery in 1982 and during the 1983 El Niño. *Fish Bull.* 84:279-286.
28. Bayer, R. D. 1986b. Nearshore flights of seabirds past Yaquina Estuary, Oregon, during the 1982 and 1983 summers. *Western Birds* 16:169-173.
29. Bayer, R. D. 1986c. Breeding success of seabirds along the mid-Oregon coast concurrent with the 1983 El Niño. *Murrelet* 67:23-26.
30. Beach, U. S. 1937. The destruction of trout by fish ducks. *Transactions of American Fisheries Society* 66:338-342.
31. Beach, R., A. C. Geiger, S. J. Jeffries, S. D. Treacy and B. L. Troutman. 1985. Marine mammals and their interactions with fisheries of the Columbia River and adjacent waters, 1980-1982. Third Annual Report. Washington Department of Wildlife, Wildlife Management Division. Olympia.
32. Beamish, R. J., and R. Boullion. 1993. Pacific salmon production trends in relation to climate. *Canadian Journal of Fisheries and Aquatic Sciences* 50:1002-1016.
33. Beechie, T. J., and T. H. Sibley. 1990. Evaluation of the TFW stream classification system: stratification of physical habitat area and distribution. Timber/Fish/Wildlife Final Report TFW-16B-89-006, Washington Department of Natural Resources. Olympia.
34. Bell, D. T., and S. K. Sipp. 1975. The litter stratum in the streamside forest ecosystem. *Oikos* 26:391-397.
35. Ben-David, M. T., R.W. Flynn, and D.M. Schell. 1997. Annual and seasonal changes in diets of martens: evidence from stable isotope analysis. *Oecologia* 111:280-291.
36. Ben-David, M. T. 1997. Timing of reproduction in wild mink: the influence of spawning Pacific salmon. *Canadian Journal of Zoology* 75:376-382.
37. Ben-David, M. T., A. Hanley, D. R. Klein, and D. M. Schell. 1997. Seasonal changes in diets of coastal and riverine mink: the role of spawning Pacific salmon. *Canadian Journal of Zoology* 75:803-811.

38. Benda, L., T. J. Beechie, R. C. Wissmar, and A. Johnson. 1992. Morphology and evolution of salmonid habitats in a recently deglaciated river basin, Washington State, USA. *Can. J. Fish. Aquat. Sci.* 49:1246-1256.
- 38.a Benda, L. E., K. J. Miller, T. Dunne, G. H. Reeves, and J. K. Agee. 1998. Dynamic landscape systems. Pages 261-288 *in* R. J. Naiman, and R. C. Bilby, editors. *River Ecology and Management: lessons from the Pacific coastal ecoregion*. Springer New York, NY.
39. Bent, A. C. 1921. Life histories of North American gulls and terns. United States National Museum. Bulletin No.1131:65-71.
40. Berman, C., and T. P. Quinn. 1991. Behavioural thermoregulation and homing by spring chinook salmon, *Oncorhynchus tshawytscha* Walbaum, in the Yakima River. *J. Fish. Biol.* 39:301-312.
41. Beschta, R. L., R. E. Bilby, G. W. Brown, L. B. Holtby, and T. D. Hofstra. 1987. Stream temperature and aquatic habitat: fisheries and forestry interactions, pages 191-232 *in*: E. O. Salo and T. W. Cundy, editors. *Streamside Management: Forestry and Fishery Interactions*. Institute of Forest Resources Contribution Number 57. University of Washington, Seattle.
- 41a. Beverton, R. J. H., and S. J. Holt. 1957 (Reprinted 1993). On the dynamics of exploited fish populations. Reprinted by Chapman and Hall, distributed in U. S. by American Fisheries Society. Bethesda, MD.
42. Bilby, R. E. 1988. Interactions between terrestrial and aquatic systems, pages 13-29 *in*: K. J. Raedeke, editor. *Streamside Management: Riparian Wildlife and Forestry Interactions*. Contrib. No. 59, Inst. For. Res., University of Washington, Seattle, WA.
43. Bilby, R. E. 1984. Post-logging removal of woody debris affects stream channel stability. *J. For.* 82:609-613.
44. Bilby, R. E., and J. W. Ward. 1989. Changes in characteristics and function of woody debris with increasing size of streams in western Washington. *Transactions of American Fisheries Society* 118:368-378.
45. Bilby, R. E., and J. W. Ward. 1991. Characteristics and function of large woody debris in streams draining old-growth, clear-cut, and second-growth forests in southwestern Washington. *Canadian Journal of Fisheries and Aquatic Sciences* 48:2499-2508.
46. Bilby, R. E., and P. A. Bisson. 1992. Allochthonous versus autochthonous organic matter contributions to the trophic support of fish populations in clear-cut and old-growth forested streams. *Canadian Journal of Fisheries and Aquatic Sciences* 49:540-551.
47. Bilby, R.E., B.R. Fransen and P.A. Bisson. 1996. Incorporation of nitrogen and carbon from spawning coho salmon into the trophic system of small streams: evidence from stable isotopes. *Canadian Journal of Fisheries and Aquatic Sciences* 53:164-173.
48. Bilby, R. E., B. R. Fransen, P. A. Bisson, and J. W. Walter. 1998. Response of juvenile coho salmon (*Oncorhynchus kisutch*) and steelhead (*Oncorhynchus mykiss*) to the addition of salmon carcasses to two streams in Southwestern Washington, U.S.A. *Can. J. Fish. Aquat. Sci.* 55:1909-1918.
49. Bilby, R. E., and P. A. Bisson. 1998. Function and distribution of large woody debris. In: R. J. Naiman and R. E. Bilby, editors. *River Ecology and Management: Lessons From The Pacific Coastal Ecoregion*. Springer Verlag. New York City, NY.
- 49a. Bilton, H. T., D. F. Alderdice and J. T. Schnute. 1982. Influence of time and size at release of juvenile coho salmon (*Oncorhynchus kisutch*) on returns at maturity. *Canadian Journal of Fisheries and Aquatic Sciences* 39:426-447.
- 49b. Bilton, H. T. 1984. Returns of chinook salmon in relation to juvenile size at release. *Canadian Technical Report Fisheries and Aquatic Sciences* 1245.

50. Bisson, P.A., R.E. Bilby, M.D. Bryant, C.A. Dolloff, G.B. Grette, R.A. House, M.L. Murphy, K V. Koski, and J.R. Sedell. 1987. Large woody debris in forested streams in the Pacific Northwest: past, present, and future. Pages 143-190 in E.O. Salo and T.W. Cundy, editors. *Streamside Management: Forestry and Fishery Interactions*. Contribution Number 57, Institute of Forest Resources, University of Washington, Seattle, WA.
51. Bisson, P. A., and G. E. Davis. 1976. Production of juvenile chinook salmon, *Oncorhynchus tshawytscha*, in a heated model stream. *NOAA Fisheries Bulletin* 74:763-774.
52. Bisson, P. A., and J. R. Sedell. 1984. Salmonid populations in streams in clearcut vs. old-growth forests of western Washington. Pages 121-130 in W. R. Meehan, T. R. Merrell, Jr., T. A. Hanley, editors. *Proceedings of a Symposium: Fish and Wildlife Relationships in Old-growth Forests*. Held April, 1982, Juneau, AK.
53. Bisson, P.A., T. Quinn, G. H. Reeves, and S. V. Gregory. 1992. Best management practices, cumulative effects, and long-term trends in fish abundance in Pacific Northwest river systems. Pages 223-265 in R. J. Naiman, editor. *Watershed management, Balancing Sustainability and Environmental Change*. Springer-Verlag. New York City, N.Y.
54. Bisson, P. A., G. H. Reeves, R. E. Bilby, and R. J. Naiman. 1997. Watershed management and Pacific salmon: Desired future conditions. Pages 447-474 in *Pacific Salmon and Their Ecosystems: Status and Future Options*. Chapman and Hall, New York, N.Y.
55. Bisson, P. A., and R. E. Bilby. 1998. Organic matter and trophic dynamics. In: R. J. Naiman and R. E. Bilby, editors. *River Ecology and Management: Lessons From the Pacific Coastal Ecoregion*. Springer Verlag. New York, N.Y.
56. Bjornn, T. C., and D. W. Reiser. 1991. Habitat requirements of salmonids in streams. Pages 83-138 in W. R. Meehan, editor. *Influences of Forest and Rangeland Management on Salmonid Fishes and Their Habitats*. American Fisheries Society Special Publication No. 19.
57. Bjornn, T. C. 1971. Trout and salmon movements in two Idaho streams as related to temperature, food, stream flow, cover, and population density. *Transactions of American Fisheries Society* 100:423-438.
- 57a. Bledsoe, L. J., et al. 1989. The Puget Sound runs of salmon: An examination of the changes in run size since 1896. Pages 50-61 in C. D. Levings, et al, editors. *Proceedings of the national workshop on effects of habitat alteration on salmonid stocks*. Canadian Special Publication Fisheries and Aquatic Sciences. No. 105.
58. Bollen, W. B., and K. C. Lu. 1968. Nitrogen transformations in soils beneath red alder and conifers. Pages 141-148 in J. M. Trappe, J. F. Franklin, R. F. Tarrant, and G. M. Hansen, editors. *Biology of Alder*. USDA Forest Service. Portland, OR.
59. Booth, D. B., and C. R. Jackson. 1997. Urbanization of aquatic systems: Degradation thresholds, stormwater detection, and the limits of mitigation. *Journal of American Water Resources Association* 33:1077-1090.
60. Booth, D. E. 1991. Estimating prelogging old-growth in the Pacific Northwest. *Journal of Forestry* 89:25-29.
61. Bormann, F. H., and G. E. Likens. 1967. Nutrient cycling. *Science* 155:424-429.
62. Brickell, D. C., and J. J. Goering. 1970. Chemical effects of salmon decomposition on aquatic ecosystems. Pages 125-138 in R. S. Murphy, editors. *Proceedings of the Symposium on Water Pollution Control in Cold Climates*. U. S. Government Printing Office, Washington, DC.
63. Briggs, D. T., and C. W. Davis. 1972. A study of predation by sea lions on salmon in Monterey Bay. *California Fish Game* 58:37-43.
64. Brinson, M. M., B. L. Swift, R. C. Plantico, and J. S. Barclay. 1981. *Riparian ecosystems: Their ecology and status*. USDI Fish and Wildlife Service, Biological Services Program. Kearneysville, W.V.

65. Brittell, J., D. Sweeney and S. T. Knick. 1979. Washington bobcats: diet, population dynamics and movement. 1979: 107-110.
66. Brooks, A. 1928. Does the marbled murrelet nest inland? Murrelet 9:68.
67. Brown, R. F., S. Riemer and S. J. Jeffries. 1995. Food of pinnipeds collected during the Columbia River area salmon gillnet observation program, 1991-1994. Oregon Dept. of Fish and Wildlife, Wildlife Diversity Program, Technical Report #95-6-01 (Available From Oregon Dept. of Fish and Wildlife, 2501 SW First, Portland, OR 97201). 16 pp.
68. Brown, R. F., and B. R. Mate. 1983. Abundance, movements, and feeding habits of harbor seals, *Phoca vitulina*, at Netarts and Tillamook Bays, Oregon. U. S. Fish and Wildlife Service Fish. Bull. 81(2):291-301.
69. Brown, E. R. 1985. Riparian zones and freshwater wetlands. Management of wildlife and fish habitats in forests of western Oregon and Washington, Part I - Chapter Narratives. USDA, Forest Service.
70. Brown, B. 1982. Mountain in the clouds: A search for wild salmon. Simon and Schuster, New York, N.Y.
71. Brown, S., M. M. Brinson, and A. E. Lugo. 1979. Structure and function of riparian wetlands. Pages 17-31 in R.R. Johnson and J.F. McCormick, editors. Strategies for protection and management of floodplain wetlands and other riparian ecosystems. USDA For. Ser. Gen. Tech. Rep. WO-12.
72. Browne, P., R. L. DeLong H. R. Huber, and J. L. Laake. 1997. Pinniped predation on endangered salmonids in Washington and Oregon: harbor seal food habits on the Columbia River. Pages 109-118 in P. S. Hill, and D. P. DeMaster, editors. Marine Mammal Protection Act and Endangered Species Act Implementation Program 1996. AFSC Processed Report 97-10.
73. Burcham, J. S. 1904. Notes on the habits of the water ousel (*Cinclus mexicanus*). Condor. 6:50.
74. Burger, A. E., R. P. Wilson, D. Garnier and M. P. T. Wilson. 1993. Diving depths, diet, and underwater foraging of Rhinoceros Auklets in British Columbia. Can. J. Zoo. 71:2528-2540.
75. Burgess, S. A., and J. R. Bider. 1980. Effects of stream habitat improvements on invertebrates, trout populations, and mink activity. Journal of Wildlife Management 44:871-880.
76. Burger, A. E. 1993 (1994). Mortality of seabirds assessed from beached-bird surveys in southern British Columbia. Canadian Field Naturalist 107:164-176.
77. Burgner, R. L. 1991. The life history of sockeye salmon (*Oncorhynchus nerka*). Pages 1-118 in C. Groot and L. Margolis. Pacific Salmon Life Histories. University of British Columbia Press, Vancouver.
78. Bury, R. B., and P. S. Corn. 1988. Responses of aquatic and streamside amphibians to timber harvest: A review. Pages 165-188 in K.J. Raedeke, editor. Streamside Management: Riparian Wildlife and Forestry Interactions. Contrib. No. 59, Inst. of Forest Resources, University of Washington, Seattle, WA.
79. Busby, P. J., T. C. Wainwright, G. J. Bryant, L. J. Lierheimer, R. S. Waples, F. W. Waknitz, and I. V. Lagomarsino. 1996. Status review of west coast steelhead from Washington, Idaho, Oregon, and California. U.S. Dept. of Commerce. NOAA Tech. Memo. NMFS-NWFSC-27. 261 pp.
80. Bustard, D. R., and D. W. Narver. 1975. Aspects of the winter ecology of juvenile coho salmon (*Oncorhynchus kisutch*) and steelhead trout (*Salmo gairdneri*). Canadian Journal of Fisheries Research Board 32:667-80.
81. Butler, R. L. 1991. Trout as predator. Pages 65-72 in J. Stolz and J. Schnell, editors. Trout, the Wildlife Series. Stackpole Books. Harrisburg, PA.
82. Butler, R. W. 1973. Trumpeter swans eating salmon. Vancouver National Historical Society (New Series.). 2:120.

83. Brodeur, R. D. 1989. Neutonic feeding by juvenile salmonids in coastal waters of the Northeast Pacific. *Canadian Journal of Zoology* 67:1995-2007.
- 83a. Brodeur, R. D., and W. G. Pearcy. 1990. Trophic relations of juvenile Pacific salmon off the Oregon and Washington coasts. *Fisheries Bulletin* 88:617-630.
84. California Department of Fish and Game. 1993. Final environmental document regarding bear hunting. Resource Agency, California Department of Fish and Game, Sacramento. 594 pp.
85. Campbell, R. W., and N. K. Dawe. 1990. *The Birds of British Columbia, Volume 1, Nonpasserines: Loons through waterfowl*. Royal British Columbia Museum, Victoria.
86. Carlson, A. 1991. Characterization of riparian management zones and upland management areas with respect to wildlife habitat.. Washington Timber/Fish/Wildlife Rep. T/F/W-WLI-91-001. Washington Department of Natural Resources, Olympia.
87. Carman, R. E., C. J. Cederholm, and E. O. Salo. 1984. A baseline inventory of juvenile salmonid populations and habitats in streams in Capitol Forest, Washington 1981-1982. Progress Report for Sampling During 1981 and 1982, University of Washington, Fisheries Research Institute. Report No. FRI-UW-8416, Seattle, WA. 102 pp.
88. Carter, H. R., and S. G. Sealy. 1984. Marbled Murrelet mortality due to gill-net fishing in Barkley Sound, British Columbia. Pages 212-220 in D.N. Nettleship, G.A. Sanger, and P.F. Springer, editors. *Marine Birds: Their Feeding Ecology and Commercial Fisheries Relationships*. Canadian Wildlife Service.
89. Case, D. J., and D. R McCullough. 1987. White-tailed deer forage on alewife. *Journal of Mammalogy* 68:195-197.
90. Cederholm, C. J., and N. P. Peterson. 1985. The retention of coho salmon (*Oncorhynchus kisutch*) carcasses by organic debris in small streams. *Canadian Journal of Fisheries and Aquatic Sciences* 42:1222-1225.
91. Cederholm, C. J., D. B. Houston, D. L. Cole, and W. J. Scarlett. 1989. Fate of coho salmon (*Oncorhynchus kisutch*) carcasses in spawning streams. *Canadian Journal of Fisheries and Aquatic Sciences* 46:1347-1355.
- 91a. Cederholm, C. J., R. E. Bilby, P. A. Bisson, T. W. Bumstead, B. R. Fransen, W. J. Scarlett, and J.W. Ward. 1997. Response of juvenile coho salmon and steelhead to placement of large woody debris in a coastal Washington stream. *North American Journal of Fisheries Management* 17:947-963.
92. Cederholm, C. J., Kunze, M. D., Murota, T., and A. Sibatani. 1999. Pacific salmon carcasses: Essential contributions of nutrients and energy for aquatic and terrestrial ecosystems. *Fisheries* 24(10):6-15.
93. Cederholm, C. J., and E. O. Salo. 1979. The effects of logging road landslide siltation on the salmon and trout spawning gravels of Stequaleho Creek and the Clearwater River Basin, Jefferson County, Washington. University of Washington, Fisheries Research Institute. Report No. FRI-UW-7915. Seattle, WA. 99 pp
94. Cederholm, C. J., and W. J. Scarlett. 1982. Seasonal immigrations of juvenile salmonids into four small tributaries of the Clearwater River, Washington, 1977-198. Pages 98-110 in E. L. Brannon and E. O. Salo, editors. *Proceedings of the Salmon and Trout Migratory Behavior Symposium, June 3-5, 1981*. University of Washington, School of Fisheries, Seattle.
95. Chapman, D. W., and T. C. Bjornn. 1969. Distribution of salmonids in streams, with special reference to food and feeding. Pages 153-176 in T. G. Northcote, editor. *Symposium On Salmon and Trout In Streams, H. R. MacMillan Lectures In Fisheries*. Held February 22-24, 1968, at the University of British Columbia. Vancouver.
96. Chapman, D. W. 1962. Aggressive behavior in juvenile coho salmon as a cause of emigration. *Canadian Journal of Fisheries Research Board* 19:1047-1080.

97. Chi, D. K. 1999. The effects of salmon availability, social dynamics, and people on black bear (*Ursus americanus*) fishing behavior on an Alaskan salmon stream. Ph.D. Dissertation. Utah State University, Logan, Utah. 137 pp.
- 97a. Chilcote, M. W. 1998. Conservation status of steelhead in Oregon. Portland, Oregon Department of Fish and Wildlife, Fish Division, Information Report 98-3, January, 1998
98. Clark, E. E. 1984. Indian legends of the Pacific Northwest. University of California Press. Berkeley. 225 pp.
99. Clemens, W. A., and G. V. Wilby. 1933. Food of the fur seal off the coast of British Columbia. Journal of Mammalogy 14:43-46.
- 99a. Cobb, J. N. 1930. Pacific salmon fisheries. U. S. Department of Commerce, Bureau of Fisheries Document No. 1092. Washington, D. C.
100. Coffman, W. P., and L. C. Ferrington, Jr. 1996. Chironomidae. Pages 635-754 in R. W. Merritt, and K. W. Cummins, editors. Kendall/Hunt Publishing. Dubuque, IA.
101. Commencement Bay Natural Resource Trustees. 1997. Commencement Bay Natural Resource Damage Assessment Restoration Plan and Final Programmatic Environmental Impact Statement. Prepared by the U. S. Fish and Wildlife Service and the National Oceanic and Atmospheric Administration for the Commencement Bay Natural Resource Trustees and Cooperating Agencies. Olympia and Seattle, WA.
102. Conaway, C. H. 1952. Life history of the water shrew (*Sorex palustris navigator*). The American Midland Naturalist 48:219-248.
103. Cone, J., and S. Ridlington. 1996. The Northwest Salmon Crisis - A documentary history. Oregon State University Press, Corvallis. 374 pp.
104. Congleton, J. L., S. K. Davis, and S. R. Foley. 1982. Distribution, abundance and outmigration timing of chum and chinook salmon fry in the Skagit salt marsh. Pages 153-163 in E. L. Brannon and E. O. Salo, editors. Proceedings of the Salmon and Trout Migratory Behavior Symposium, June 3-5, 1981. University of Washington, School of Fisheries, Seattle.
105. Connell, J. H. 1978. Diversity in tropical rain forests and coral reefs. Science 199:1302-1310.
106. Cooper, R., and T. H. Johnson. 1992. Trends in steelhead abundance in Washington and along the Pacific coast of North America. Report #92-90. Fisheries Management Division. Washington Department of Wildlife, Olympia.
107. Cottam, C. 1939. Food habits of North American diving ducks. U.S. Department of Agriculture Technical Bulletin No. 643. 140 pp.
108. Cottam, C., and Uhler F. M. 1937. Birds in relation to fishes. Wildlife Research and Management Leaflet BS-83. 16 pp.
109. Craig, J. A., and R. L. Hacker. 1940. The history and development of the fisheries of the Columbia River. Bulletin of the Bureau of Fisheries, Vol. XLIX.
110. Cressman, L.S. 1960. Cultural sequences at the Dalles, Oregon. A contribution to Pacific Northwest prehistory. Transactions of the American Philosophical Society, Vol. 50, Part 10. 103 pp.
- 110a. Crutchfield, J. A., and G. Pontecorvo. 1969. The Pacific salmon fisheries: A study of irrational conservation. The John Hopkins Press, Baltimore, MD.
111. Cummins, K. W., M. A. Wilzbach, D. M. Gates, J. B. Perry, and W. B. Taliaferro. 1973. Shredders and riparian vegetation. Bioscience 39:24-30.

112. Dahm, C. C. 1981. Pathways and mechanisms for removal of dissolved organic carbon from leaf leachate in streams. *Canadian Journal of Fisheries and Aquatic Sciences* 38:68-76.
113. Day, J. W., Jr., C. A. S. Hall, W. M. Kemp, and A. Yáñez-Arancibia. 1989. *Estuarine Ecology*. John Wiley & Sons, New York, N.Y. 558 pp.
114. Dalquest, W. W. 1948. *Mammals of Washington*. University of Kansas Publication Volume 2, Museum of Natural History, University of Kansas, Lawrence. 444 pp.
115. Dawson, W. L. 1909. *The Birds of Washington. A Complete, Scientific and Popular Account of the 372 Species of Birds Found in the State*. Seattle. 2.
116. Death, R. G., and M. J. Winterbourn. 1995. Diversity patterns in stream benthic invertebrate communities: the influence of habitat stability. *Ecology* 76:1446-1460.
117. DeGange, A. R. 1996. The marbled murrelet: a conservation assessment. Gen. Tech. Rep. PNW-GTR-388. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 72 pp.
118. DeShazo, J. J. 1980. Sea-run cutthroat trout management in Washington - An overview. Fisheries Management Division, Washington State Game Dept., Olympia.
119. DeVoto, B. 1981. *The journals of Lewis and Clark*. Houghton Mifflin Company. Boston, Mass. 504 pp.
120. Dolloff, C. A. 1993. Predation by River otters (*Lutra canadensis*) on juvenile coho salmon (*Oncorhynchus kisutch*) and Dolly Varden (*Salvelinus malma*) in southeast Alaska. *Canadian Journal of Fisheries and Aquatic Sciences* 50:312-315.
121. Donaldson, I. J., and F. K. Cramer. 1971. *Fishwheels of the Columbia*. Binfords and Mort, Publishers. Portland, OR.
122. Donaldson, J. R. 1967. The phosphorus budget of Iliamna Lake, Alaska as related to the cyclic abundance of sockeye salmon. Ph.D. Dissertation, University of Washington, School of Fisheries, Seattle. 141 pp.
123. Drummond, H. 1983. Aquatic foraging in garter snakes: a comparison of specialists and generalists. *Behaviour* 86:1-30.
124. Dunstone, N. 1993. *The mink*. T & A. D. Poyser Ltd, London. 232 pp.
125. Dzinbal, A., and R. L. Jarvis. 1982. Coastal feeding ecology of harlequin ducks in Prince William Sound, Alaska, during summer. Pages 6-10 in D. N. Nettleship, G. A. Sanger and P. F. Springer, editors. *Marine Birds: Feeding Ecology and Commercial Fisheries Relationships*. Proceedings of the Pacific Seabird Group Symposium, Seattle, Washington. In English With French Summ. WR 199.
126. Eagle, T. C., and J. S. Whitman. 1987. Mink. Pages 614-624 in M. Novak, J. A. Baker, M. E. Obbard and B. Malloch, editors. *Wild Furbearer Management and Conservation in North America*. Ministry of Natural Resources, Ontario.
127. Eastman, D. E. 1996. Response of freshwater fish communities to spawning sockeye salmon (*Oncorhynchus nerka*). M. S. Thesis, University of Washington, School of Fisheries, Seattle, WA.
128. Edie, B. G. 1975. A census of the juvenile salmonids of the Clearwater River Basin, Jefferson County, Washington, in relation to logging. M. S. Thesis. University of Washington, School of Fisheries, Seattle, WA. 86 pp.
- 128a. Edmonds, R. L. 1980. Litter decomposition and nutrient release in Douglas-fir, red alder, western hemlock, and Pacific silver fir ecosystems in western Washington. *Canadian Journal of Forestry Research* 10:327-337.
129. Ehinger, C. E. 1930. Some studies of the American dipper or water ouzel. *Condor* 47:487-498.

130. Eipper, A. W. 1956. Differences in the vulnerability of the prey of nesting kingfishers. *Journal of Wildlife Management* 20:177-183.
131. Elliott, S. T., and R. Bartoo. 1981. Relation of larval *Polypedium* (Diptera: Chironomidae) to pink salmon eggs and alevins in an Alaska stream. *Prog. Fish-Cult.* 43:220-221.
132. Elmore, W. 1992. Riparian responses to grazing practices. Pages 442-457 in R. B. Naiman, editor. *Watershed Management: Balancing Sustainability and Environmental Change*. Springer-Verlag. New York, N. Y.
133. Elphick, C.S., and T.L. Tibbitts. 1998. Greater Yellowlegs (*Tringa melaoleuca*). *The Birds of North America*, No. 355. A. Poole and F. Gill, editors. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists Union, Washington D.C.
134. Elson, P. F. 1962. Predator-prey relationship between fish-eating birds and Atlantic salmon. *Fish. Res. Board Can. Bull.* No 133. 87 pp.
135. Emmett, R. L. 1997. Estuarine survival of salmonids: The importance of interspecific and intraspecific predation and competition. Pages 147-158 in R.L. Emmett and M.H. Schiewe, editors. *Estuarine and Ocean Survival of Northeastern Pacific Salmon*. NOAA Technical Memorandum NMFS-NWFSC-29.
136. Emmett, R. L., and M. H. Schiewe. 1997. Estuarine and ocean survival of Northeastern Pacific salmon: Proceedings of the workshop. U.S. Dept. Commerce., NOAA Tech. Memo. NMFS-NWFSC-29.
137. Erman, D. C., J.D. Newbold, and K. B. Roby. 1977. Evaluation of streamside bufferstrips for protecting aquatic organisms. California Water Resources Center, Contribution Number 165, University of California, Davis.
138. Erickson, J. 1988. Competition for fish. *Headlight Herald*. June 4th(A-4).
139. Eriksson, M. O. G. D. Blomqvist M. Hake and O. C. Johansson. 1990. Parental feeding in the Red-throated Diver *Gavia stellata*. *Ibis*.132:1-13.
140. Everest, F. H., N. B. Armantrout, S. M. Keller, W. D. Parante, J. R. Sedell, T. E. Nickelson, J. M. Johnston, and G. N. Haugen. 1985. Salmonids, Pages 199-226 in R. Brown, editor. *Management of Wildlife and Fish Habitats in Forests of Western Oregon and Washington*. Volume 1. USDA, Forest Service. Pacific Northwest Region.
141. Everitt, R. D., P. J. Gearin, J. S. Skidmore and R. L. DeLong. 1981. Prey items of harbor seals and California sea lions in Puget Sound, Washington. *Murrelet* 62:83-86.
- 141a. Falconer, D. S. 1989. *Introduction to quantitative genetics*. 3rd edition. New York, Longman Scientific and Technical.
142. Farley, L. C. 1980. *The Behavioral-Ecology of Swans Wintering in Southeast Alaska*. MS Thesis. Idaho State University. 73 pp.
143. Felleman, F. L., J. R. Heimlich-Boran and R. W. Osborne. 1991. The feeding ecology of killer whales (*Orcinus orca*) in the Pacific Northwest. Pages 113-147 in K. Pryor, and K. S. Norris, editors. *Dolphin Societies, Discoveries and Puzzles*. University of California Press, Berkeley, Los Angeles & Oxford.
144. Fiscus, C. H. and G. A. Baines. 1966. Food and feeding behavior of Steller and California sea lions. *Journal of Mammalogy* 47:195-200.
145. Fisher, J. P., and W. G. Pearcy. 1988. Growth of juvenile coho salmon (*Oncorhynchus kisutch*) in the ocean off Oregon and Washington, USA, in years of differing coastal upwelling. *Canadian Journal of Fisheries and Aquatic Sciences*. 45:1036-1044.

146. Foerster, R. E. 1968. The sockeye salmon (*Oncorhynchus nerka*). Fish. R. Bd. Can. Bull. 162. Ottawa, Canada. 422 pp.
147. Fitch, H. S. 1984. *Thamnophis couchii*. Catalogue of American Amphibians and Reptiles. 351.1-351.3.
148. Fitch, H. S. 1941. The feeding habits of California garter snakes. California Dept. Fish and Game 27:2-32.
149. Fontaine, P. M., M. O. Hammill, C. Barrette, and M. C. Kingsley. 1994. Summer diet of the harbour porpoise (*Phocoena phocoena*) in the estuary and the Northern Gulf of St. Lawrence. Canadian Journal of Fisheries and Aquatic Sciences 5:172-178.
150. Forbes, L. C. and K. 1982. Simpson. Behavioral studies of great blue herons at Pender Harbor and Sechelt, British Columbia, in 1980. Unpublished Report., Canadian Wildlife Service, Delta, B.C.
151. Forest Ecosystem Management Assessment Team (FEMAT). 1993. Forest Ecosystem Management: An Ecological, Economic, and Social Assessment. USDA, Forest Service.
152. Frame, G. W. 1974. Black bear predation on salmon at Olsen Creek, Alaska. Zeitung der Tierpsychologie 35:23-38.
153. Francis, R. C., and S. R. Hare. 1994. Decadal-scale regime shifts in the large marine ecosystems of the Northeast Pacific: A case for historical science. Fish. Oceanogr. 3:279-291.
154. Franklin, J. F. 1992. Scientific basis for new perspectives in forests and streams. Pages 25-72 in R. J. Naiman, editor. Watershed Management: Balancing Sustainability and Environmental Change. Springer-Verlag. New York, N.Y.
155. Friesen, W. 1990. Winter dietary studies of juvenile coho salmon (*Oncorhynchus kisutch*) utilizing two enhanced wall-base channels along the Clearwater River in Jefferson County, Washington. Master of Environmental Studies, The Evergreen State College, Olympia, WA. 114 pp.
156. Fraser, J. M. 1972. Recovery of planted brook trout, splake, and rainbow trout from selected Ontario lakes. Canadian Journal of Fisheries Research Board 29:129-142.
157. Fraser, J. M. 1974. An attempt to train hatchery reared brook trout to avoid predation by the common loon. Transactions of American Fisheries Society 103:815-818.
158. French, J. M., and J. R. Koplín. 1977. Distribution, abundance, and breeding status of ospreys in northwestern California. Pages 223-240 in Transactions of North American Osprey Research Conference. U.S. Natl. Park Serv. Trans. Proc. Ser. No. 2., WR 179.
159. Fujiwara, M., and R. C. Highsmith. 1997. Harpacticoid copepods: Potential trophic link between inbound adult salmon and outbound juvenile salmon. Marine Ecology Progress Series 158:205-216.
160. Fuss, H. J. 1982. Age, growth, and instream movement of Olympic Peninsula coastal cutthroat trout (*Salmo clarki clarki*). M. S. Thesis. University of Washington, School of Fisheries, Seattle, WA. 126 pp.
161. Gabrielson, I. N., and S. G. Jewett. 1940. Birds of Oregon. Oregon State College, Corvallis. 650 pp.
162. Garrett, A. M. 1998. Interstream movements of coastal cutthroat trout (*Oncorhynchus clarki clarki*) in the Clearwater River, Jefferson County, Washington. Master of Environmental Studies. The Evergreen State College, Olympia, WA. 212 pp.
163. Gaston, A. J., and S. B. C. Dechesne. 1996. Rhinoceros Auklet, *Cerorhinca monocerata*. In: A. Poole and F. Gill, editors. The Birds of North America No. 212. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologist's Union, Washington, D.C.

164. Gearin, P. J., S. R. Melin, R. L. DeLong, H. Kajimura and M. A. Johnson. 1994. Harbor porpoise interactions with a chinook salmon set-net fishery in Washington state. Pages 427-438 in W. F. Perrin, G. P. Donovan, and J. Barlow, editors. Gillnets and Cetaceans: Incorporating the Proceedings of the Symposium and Workshop on the Mortality of Cetaceans in Passive Fishing Nets and Traps. La Jolla, CA. October 1990. Whaling Commission Special Issue No.15. WR 251.
165. Gearin, P. J., R. Pfeifer, S. J. Jeffries, R. L. DeLong and M. A. Johnson. 1988. Results of the 1986-1987 California sea lion - steelhead trout predation control program at the Hiram M. Chittenden Locks. U. S. Dept. Comm. NWAFC Processed Report 88-30. 111 pp.
- 165a. Geiger, H. J., W. W. Smoker, L. A. Zhivotovsky, and A. J. Gharrett. 1998. Variability of family size and marine survival in pink salmon (*Oncorhynchus gorbuscha*) has implications for conservation biology and human use. Canadian Journal of Fisheries and Aquatic Sciences 54: 2684-2690.
166. Glova, G. J. 1986. Interactions for food and space between experimental populations of juvenile coho salmon (*Oncorhynchus kisutch*) and coastal cutthroat trout (*Salmo clarki*) in a laboratory stream. Hydrobiologia 132:155-168.
167. Gould, V. E. 1934. A monograph of the belted kingfisher *Megaceryle alcyon* (Linnaeus). 311 pp.
168. Graf, W. 1949. Observations on the salamander *Dicamptodon*. Copeia 1:79-80.
169. Graybill, M.R., and J. Hodder. 1985. Effects of the 1982-1983 El Niño on reproduction of six species of seabirds in Oregon. Pages 205-210 in W.S. Wooster and D.L. Fluharty, editors. El Niño North. Washington Sea Grant Program, University of Washington, Seattle, WA.
170. Graybill, M. R. 1981. Haul out patterns and diet of harbor seals, *Phoca vitulina*, in Coos County, Oregon. M. S. Thesis, University of Oregon, Eugene. 55 pp.
171. Gregory, S. V., F. J. Swanson, and W. A. McKee. 1991. An ecosystem perspective of riparian zones. BioScience 40:540-551.
172. Gregory, S. V. 1980. Effects of light, nutrients, and grazing on periphyton communities in streams. Ph. D. Dissertation. Oregon State University, Corvallis, OR.
173. Gregory, S. V., G. A. Lamberti, D. C. Erman, K. V. Koski, M. L. Murphy, and J. R. Sedell. 1987. Influences of forest practices on aquatic production. Pages 233-255 in E. O. Salo, and T. Cundy, editors. Streamside Management - Forestry and Fishery Interactions. College of Forest Resources, University of Washington. Contribution No. 57. Seattle.
174. Gregory, S. V., and P. A. Bisson. 1997. Degradation and loss of anadromous salmonid habitat in the Pacific Northwest. Pages 277-314 in D. J. Stouder, P. A. Bisson, and R. J. Naiman, editors. Pacific Salmon and Their Ecosystems. Chapman and Hall Publishers, New York, N.Y.
- 174a. Gresh, T, J. Lichatowich, and P. Schoonmaker. 2000. An estimation of historic and current levels of salmon production in the Northeast Pacific ecosystem: Evidence of a nutrient deficit in the freshwater systems of the Pacific Northwest. Fisheries 25(1):15-21.
175. Grette, G. 1985. The role of large organic debris in juvenile salmonid rearing habitat in small streams. M.S. Thesis, University of Washington, School of Fisheries. Seattle, WA. 105 pp.
176. Grinnell, J., J. Dixon, and J. M. Linsdale. 1937. Fur-bearing mammals of California. Their natural history, systematic status, and relations to man. University of California Press, Berkeley.
177. Griswold, G. 1953. Aboriginal patterns of trade between the Columbia Basin and the northern plains. M. A. Thesis, Montana State University. Missoula.

178. Groot, C., and L. Margolis. 1991. Pacific Salmon Life Histories, University of British Columbia Press, Vancouver, BC, Canada. 564 pp.
179. Gross, M. R., R. M. Coleman, and R. M. McDowell. 1988. Aquatic productivity and the evolution of diadromous fish migration. *Science* 239:1291-1293.
180. Haig-Brown, R. 1946. A river never sleeps. Crown Publishers. New York, N.Y.
181. Hall, J. D., P. A. Bisson, and R. E. Gresswell. 1997. Sea-run cutthroat trout-biology, management, and future considerations. Proceedings of a Symposium, Reedsport, Oregon. October 12-14, 1995. Published by the Oregon Chapter of the American Fisheries Society. Corvallis. 183 pp.
182. Hall, J. D. 1986. Notes on the distribution and feeding behavior of killer whales in Prince William Sound, Alaska. Pages 69-83 in B. C. Kirkevold, and J. S. Lockard, editors. Behavioral Biology of killer whales. Alan R. Liss, Inc., New York. 457 pp.
183. Hamilton, A. N., and W. R. Archibald. 1985. Grizzly bear habitat in the Kimsquit River Valley, coastal British Columbia: evaluation. Pages 50-57 in G. P. Contreras, and K. E. Evans, editors. Proceedings - Grizzly Bear Habitat Symposium, Missoula, Montana, April 30 - May 2, 1985. Intermountain Research Station, Ogden.
184. Hamilton, A. N., and F. L. Bunnell. 1987. Foraging strategies of coastal grizzly bears in the Kimsquit River Valley, British Columbia. *Int. Conf. Bear Res. Manage.* WR 213. 7:187-197.
185. Hampton, P. D. 1981. The wintering and nesting behavior of the trumpeter swan. M.S. Thesis. University of Montana. 60 pp.
186. Hansen, W. R. 1980. Western aquatic garter snakes in central California: an ecological and evolutionary perspective. M.S. Thesis, California State University, Fresno. 78 pp.
187. Hansen, A. J., E. L. Boeker, J. I. Hodges, and D. R. Cline. 1984. Bald eagles of the Chilkat Valley, Alaska: ecology, behavior, and management. National Audubon Society. New York, N.Y.
188. Hanson, L. C. 1993. The foraging ecology of harbor seals, *Phoca vitulina*, and California sea lions, *Zalophus californianus*, at the mouth of the Russian River, California. M. A. Thesis, Sonoma State University, Rohnert Park, CA. 70 pp.
189. Hare, S. R., N. J. Mantua and R. C. Francis. 1999. Inverse production regions: Alaska and West Coast Pacific salmon. *Fisheries* 24: 6-14.
190. Hartman, W., and R. Raleigh. 1964. Tributary homing of sockeye salmon at Brooks and Karluk lakes, Alaska. *Canadian Journal of Fisheries Research Board* 21:485-504.
- 190-a. Hartman, G. F., and J. C. Scrivener. 1990. Impacts of forestry practices on a coastal stream ecosystem, Carnation Creek, British Columbia. *Canadian Journal of Fisheries and Aquatic Sciences* 223: 148p.
191. Harvey, J. T., and M. J. Weise. 1997. Impacts of California sea lions and Pacific harbor seals on salmonids in Monterey Bay, California. MLML Technical Publication No.97-03. Moss Landing Marine Laboratories. 31 pp.
192. Hasegawa, S., T. Hirano, T. Ogasawara, M. Iwata, T. Akiyama, and S. Arai. 1987. Osmoregulatory ability of chum salmon, *Oncorhynchus keta*, reared in fresh water for prolonged periods. *Fish Physiol. Biochem.* 4:101-110.
193. Hatler, D. F. 1976. The coastal mink on Vancouver Island, British Columbia. Ph. D. Dissertation, University British Columbia, Vancouver.

194. Hayman, R. A., E. M. Beamer, and R. E. McClure. 1996. Fiscal Year 1995 Skagit River chinook restoration research. Final project performance report, National Marine Fisheries Service, Contract# 3311 for FY 1995. Skagit System Cooperative. La Conner, WA.
195. Hayward, T. L. 1993. Preliminary observations of the 1991-1992 El Niño in the California Current. California Cooperative Oceanic Fisheries Investigations Reports 34:21-29.
196. Healey, M. C. 1982. Juvenile Pacific salmon in estuaries: The life support system. Pages 315-341 in V. S. Kennedy, editor. Estuarine Comparisons. Academic Press. New York, N.Y. 709 pp.
197. Healey, M. C. 1991. Life history of chinook salmon (*Oncorhynchus tshawytscha*). Pages 311-393 in C. Groot and L. Margolis, editors. Pacific Salmon Life Histories, University of British Columbia Press. Vancouver.
198. Heard, W. R. 1991. Life history of pink salmon (*Oncorhynchus gorbuscha*). Pages 121-230 in C. Groot and L. Margolis, editors. Pacific Salmon Life Histories, University of British Columbia Press. Vancouver.
199. Heimlich-Boran, J. R. 1986. Fishery correlations with the occurrence of killer whales in Greater Puget Sound. Pages 113-131 in B. C. Kirkevold, and J. S. Lockard, editors. Behavioral biology of killer whales. Alan R. Liss, Inc., New York, N.Y.
200. Heimlich-Boran, J. R. 1988. Behavioral ecology of killer whales (*Orcinus orca*) in the Pacific Northwest. Canadian Journal of Zoology 66:565-578.
201. Hemstrom, M. A., and J. F. Franklin. 1982. Fire and other disturbances of the forests in Mount Rainier National Park. Quaternary Research 18:32-51.
202. Henny, C. J., and M. R. Bethers. 1971. Population ecology of the great blue heron with special reference to western Oregon. Canadian Field-Naturalist 85:205-209.
203. Herder, M. J. 1983. Pinniped fishery interactions in the Klamath River system, July 1979 to October 1980. Southwest Fish. Cent., Admin. Rep. LJ-83-12C. (Available From Southwest Fisheries Science Center, Natl. Mar. Fish. Serv., NOAA, P.O. Box 271, La Jolla, CA 92038). 71 pp.
204. Hewson, R. 1995. Use of salmonid carcasses by vertebrate scavengers. Journal of Zoology (London). 235:53-65.
- 204a. Hilborn, R., and C. J. Walters. 1992. Quantitative fish stock assessment: choice, dynamics & uncertainty. Chapman and Hall. New York, New York.
205. Hildebrand, G. V. S. D. Farley C. T. Robbins T. A. Hanley K. Titus and C. Servheen. 1996. Use of stable isotopes to determine diets of living and extinct bears. Can. J. Zoo. 74:2080-2088.
206. Hildebrand, G. V., S. D. Farley, C. T. Robbins, T. A. Hanley, K. Titus, and C. Servheen. 1996. Use of stable isotopes to determine diets of living and extinct bears. Canadian Journal of Zoology 74:2080-2088.
207. Hicks, B.J., R. L. Beschta, and R. D. Harr. 1991b. Long-term changes in streamflow following logging in western Oregon and associated fisheries implications. Water Resources Bulletin 27:217-226.
208. Hicks, B. J., J. D. Hall, P.A. Bisson, and J. R. Sedell. 1991a. Response of salmonids to habitat changes. Pages 483-518 in W.R. Meehan, editor. Influences of Forest and Rangeland Management On Salmonid Fishes and Their Habitat. Amer. Fish. Soc., Spec. Pub. 19. Bethesda, MD.
209. Hilborn, R. 1992. Hatcheries and the future of salmon in the northwest. Fisheries 17:5-8.
210. Hodder, J., and M.R. Graybill. 1985. Reproduction and survival of seabirds in Oregon during the 1982-1983 El Niño. Condor 87:535-541.

211. Hogan, D.L., and J. W. Schwab. 1991. Stream channel response to landslides in the Queen Charlotte Islands, B.C.: changes affecting pink and chum salmon habitat. Pages 222-236 in B. White and I. Guthrie, editors. Proceedings Of the 15th Northeast Pacific Pink and Chum Workshop. Canada Department of Fisheries and Oceans. Vancouver, B.C.
212. Hoffman, T., and T. Hall. 1988. Tillamook trip report regarding smolt/cormorant problem. U. S. Dept. of Agriculture, Animal Damage Control Program Interoffice Memorandum About Their Cormorants Collection on 27 April 1988.
213. Holland, D. C. 1985a. *Clemmys marmorata* (western pond turtle). Feeding. Herpetol. Review, WR 203.16:112-113.
214. Holland, D. C. 1985b. An ecological and quantitative study of the Western Pond Turtle (*Clemmys marmorata*) in San Luis Obispo County, California. M.S. Thesis, California State University, Fresno. 181 pp.
- 214a. Holtby, L. B., and M. C. Healey. 1986. Selection for adult size in female coho salmon (*Oncorhynchus kisutch*). Can. J. Fish. Aquat. Sci. 43:1946-1959.
215. Holtby, L. B. 1988. Effects of logging on stream temperatures in Carnation Creek, British Columbia, and associated impacts on the coho salmon (*Oncorhynchus kisutch*). Can. J. Fish. Aquat. Sci. 45:502-515.
- 215a. Holtby, L. B., B. C. Andersen and R. K. Kadowaki. 1990. Importance of smolt size and early ocean growth to interannual variability in marine survival of coho salmon (*Oncorhynchus kisutch*). Canadian Journal of Fisheries and Aquatic Sciences 47: 2181-2194.
216. Horner, R.R., and B. W. Mar. 1982. Guide for water quality impact assessment of highway operations and maintenance. Report to Washington Dept. of Transp. FHWA WA-RD-39.14. Dept. of Civil Engineering, University of Washington, Seattle, WA.
217. Hunt, W. G., B. S. Johnson and R. E. Jackman. 1992. Carrying capacity for bald eagles wintering along a northwestern river. Journal of Raptor Research 26:49-60.
218. Hunt, W. G., J. M. Jenkins, R. E. Jackman, C. G. Thelander, and A. T. Gerstell. 1992. Foraging ecology of bald eagles on a regulated river. Journal of Raptor Research 26:243-256.
219. Hunt, W. A. 1993. Jasper National Park harlequin duck research project: 1992 pilot projects - interim results. Jasper Warden Service Biological Report Series, No. 1, Heritage Resource Conservation, Parks Canada, Jasper, Alberta. 67 pp.
220. Hunt, W. G., B. S. Johnson, and R. E. Jackman. 1992. Carrying capacity for bald eagles wintering along a northwestern river. Journal of Raptor Research 26(2):49-60.
221. Huntington, C. W., W. Nelhsen, and J. Bowers. 1996. A survey of healthy native stocks of anadromous salmonids in the Pacific Northwest and California. Fisheries 21(3).
222. Huntsman, S. G. 1941. Cyclical abundance of birds versus salmon. Canadian Journal of Fisheries Research Board 5:227-235.
223. Hynes, H. B. N. 1970. The ecology of running waters. University of Liverpool Press. Liverpool, England. 555 pp.
224. Imler, R. H., and H. R. Sarber. 1947. Harbor seals and sea lions in Alaska. U. S. Fisheries Wildlife Services Special Scientific Report No.28. 28 pp.
225. Science Team. 1998. Pinniped and seabird predation: implications for recovery of threatened stocks of salmonids in Oregon under the Oregon Plan for Salmon and Watersheds. Technical Report 1998-2 to the Oregon Plan for Salmon and Watersheds. Governor's Natural Resources Office, Salem, OR. 27 pp.

- 225a. IEEE/ASTM SI 10-1997 (Revision and redesignation of ANSI/IEEE Std. 268-1992 and ASTM E 380-93). 1997. Standard for Use of the International System of Units (SI): The Modern Metric System. Institute of Electrical and Electronics Engineers, Inc., New York, N.Y.
226. Inkpen, E. L., and S. S. Embrey. 1998. Nutrient transport in the major rivers and streams of the Puget Sound Basin, Washington. U. S. Department of the Interior-U. S. Survey, National Water-Quality Assessment Program.
227. Iwata, M., T. Hirano, and S. Hasegawa. 1982. Behavior and plasma sodium regulation of chum salmon fry during transition into seawater. *Aquaculture* 28:133-142.
228. Jameson, R. J., and K. W. Kenyon. 1977. Prey of sea lions in the Rogue River, Oregon. *J. Mamm.* 58: 672.
229. Jeffries, S. 1985. Marine mammals of the Columbia River estuary. Wash. State Dep. Game; 95p. 1984. From U.S. Government Reports 85(2):85. 1985. Available From NTIS As PB85-107050/GAR. WR 197.
230. Jewett, S. G., W. P. Taylor, W. T. Shaw and J. W. Aldrich. 1953. Birds of Washington State. University of Washington Press, Seattle. 768 pp.
231. Johnson, W. E., and A. D. Hasler. 1954. Rainbow trout production in dystrophic lakes. *J. Wildl. Manag.* 18:113-134.
232. Johnson, J. H., and A. A. Wolman. 1984. The Humpback Whale, *Megaptera novaeangliae*. U. S. National Marine Fisheries Service Marine Fisheries Review 6:30-37.
233. Johnson, D.H., M. M. Hoover, E. L. Greda, and C.J. Cederholm. (In prep.). Relationships between Pacific Salmon and 605 species of birds, mammals, reptiles, and amphibians in Oregon and Washington.
234. Johnson, O. W., M. H. Ruckelshaus, W. S. Grant, F. W. Waknitz, A. M. Garrett, G. J. Bryant, K. Neely, and J. J. Hard. 1999. Status review of coastal cutthroat trout in Washington, Oregon, and California. U.S. Dept. Commerce, NOAA Tech. Memo. NMFS-NWFSC-37. 292 pp.
235. Johnson, R. R., and S. W. Carothers. 1982. Riparian habitats and recreation: Interrelationships and impacts in the Southwest and Rocky Mountain region. Eisenhower Consortium for West. Environ. For. Res. Bull. 12: 1-31.
236. Johnson, D.H., and T.A. O'Neil. 2001). Wildlife-Habitats Relationships in Oregon and Washington. Oregon. University Press, Corvallis, 768 pp.
237. Johnson, R. R., and S. W. Carothers. 1982. Riparian habitats and recreation: Interrelationships and impacts in the Southwest and Rocky Mountain region. Eisenhower Consortium for West. Environ. For. Res. Bull. 12:1-31.
238. Johnston, J. M. 1982. Life histories of anadromous cutthroat with emphasis on migratory behavior, pages 123-127. In: E. L. Brannon and E. O. Salo (eds.). Proceedings of the Salmon and Trout Behavior Symposium. School of Fisheries, University of Washington, Seattle.
239. Johnston, N. T., J. S. MacDonald, K. J. Hall, and P. J. Tschaplinski. 1997. A preliminary study of the role of sockeye salmon (*Oncorhynchus nerka*) carcasses as carbon and nitrogen sources for benthic insects and fishes in the 'Early Stuart' stock spawning streams, 1050 km from the ocean. Fisheries Project Report No. RD55, Fisheries Branch, Ministry of Environment, Lands, and Parks, Province of British Columbia, Canada.
240. Jones, R. E. 1981. Food habits of smaller marine mammals from northern California. Proceedings of the California Academy of Sciences 42:409-433.
241. Juday, C., W. H. Rich, G. I. Kemmerer, and A. Mean. 1932. Limnological studies of Karluk Lake, Alaska 1926-1930. Bulletin of the U. S. Bureau of Fisheries 47:407-436.

242. Jurek, R. M. 1974. Special wildlife investigations.: American River Green Heron study, 1974. California Department of Fish and Game. 25 pp.
243. Kajimura, H. 1983. Food of the Pacific White-sided Dolphin, *Lagenorhynchus obliquidens*, Dall's Porpoise, *Phocoenoides dalli*, and Northern Fur Seal, *Callorhinus ursinus*, off California and Washington with appendices of size and food of Dall's porpoise from Alaskan waters. U.S. Natl. Mar. Fish. Serv. Tech. Mem.; F/NWC-2. Ii + 30p. June 1980. From Monthly Catalog of U.S. Government Publications 83-6731. WR 189.
244. Kellyhouse, D.G. 1975. Habitat utilization by black bears in northern California. International Conference on Bear Research and Management 4:221-227.
245. Kennish, M. J. 1997. Practical handbook of estuarine and marine pollution. (CRC Press, Marine Science Series). CRC Press. Boca Raton, FL.
246. Kingery, H. E. 1996. American Dipper. The Birds of North America, No. 229. In: A. Poole and F. Gill, (eds.). The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologist's Union, Washington, D.C.
247. Kjelson, M. A., P. F. Raquel, and F. W. Fisher. 1982. Life history of fall-run juvenile chinook salmon, *Oncorhynchus tshawytscha*, in the Sacramento-San Joaquin estuary, California, pages 393-411. In: V. S. Kennedy (ed.). Estuarine Comparisons. Academic Press, New York, N.Y.
248. Kline, T. C., Jr., J. J. Goering, O. A. Mathisen, P. H. Poe, and P. L. Parker. 1990. Recycling of elements transported upstream by runs of Pacific salmon: I. ^{15}N and ^{13}C evidence in Sashin Creek, southeastern Alaska. Can. J. Fish. Aquat. Sci. 47:136-144.
249. Kline, T. C., Jr., J. J. Goering, O. A. Mathisen, P. H. Poe, P. L. Parker, and R. S. Scalan. 1993. Recycling of elements transported upstream by runs of Pacific salmon: II. ^{15}N and ^{13}C evidence in the Kvichak River watershed, Bristol Bay, southwestern Alaska. Can. J. Fish. Aquat. Sci. 50: 2350-2365.
250. Knick, S. T., S. J. Sweeney, J. R. Alldredge and J. D. Britnell. 1984. Autumn and winter food habits of Bobcats in Washington State. Great Basin Naturalist 44:70-74.
251. Knight, R. L., P. J. Randolph, G. T. Allen, L. S. Young and R. J. Wigen. 1990. Diets of nesting Bald Eagles, *Haliaeetus leucocephalus*, in western Washington. Can. Fld-Nat. 104:545-551.
252. Knight, R. L., and D. P. Anderson. 1990. Effects of supplemental feeding on an avian scavenging guild. Wildl. Soc. Bull. 18:388-394.
253. Knight, S. K, and R. L. Knight. 1983. Aspects of food finding by wintering Bald Eagles. Biology and Management of Bald Eagles and Ospreys, pages 28-29. In: D. M. Bird (ed.). Proceedings of First International Symposium on Bald Eagles and Ospreys Montreal. October, 1981. Harpell Press, Ste Anne De Bellevue, Quebec.
254. Knudsen, E. E. 2000. Managing Pacific salmon escapements: the gaps between theory and reality, pages 237-272. E. E. Knudsen, C. S. Steward, D. D. MacDonald, J. E. Williams, and D. W. Reiser (eds.). Sustainable Fisheries Management: Pacific Salmon. C. R. C. Lewis Publishers, Boca Raton, FL.
255. Knutson, K. L., and V. L. Naef. 1997. Management recommendations for Washington's Priority habitats. Washington Department of Fish and Wildlife. Olympia. 181 pp.
256. Kojima, H., M. Iwata, and T. Kurokawa. 1993. Development and temporal decrease in seawater adaptability during early growth in chum salmon, *Oncorhynchus keta*. Aquaculture 118:141-150.
257. Konkel, G. W., and J. D. McIntyre. 1987. Trends in spawning populations of Pacific anadromous salmonids. Technical Report 9, USDI - Fish and Wildlife Service.

258. Koplín, J. R., D. S. MacCarter, D. P. Garber and D. L. MacCarter. 1977. Food resources and fledgling productivity of California and Montana Ospreys. Trans. North American Osprey Res. Conf. U.S. Natl. Park Serv. Trans. Proc. Ser.; No. 2. 1977. WR 179. Pages 205-214.
259. Koski, K V. 1975. The survival and fitness of two stocks of chum salmon (*Oncorhynchus keta*) from egg deposition to emergence in a controlled-stream environment at Big Beef Creek. Ph.D. Dissertation, University of Washington, School of Fisheries. Seattle. 212 pp.
260. Koski, K V., and D. Kirchofer. 1984. A stream ecosystem in an old-growth forest in Southeastern Alaska. Part IV: Food of juvenile coho salmon in relation to abundance of drift and benthos, pages 81-88. In: W. R. Meehan, T. R. Merrell, Jr., J. W. Matthews (eds.). Proceedings of a Symposium on Fish and Wildlife Relationships in Old-Growth Forests, Held April, 12-15, 1982. Juneau, AK.
261. Krokhn, E. M. 1967. Effect of the size of sockeye migration on the phosphate regime of spawning lakes. Izvestia, Pacific Scientific Institute of Fisheries and Oceanography 64: 353-364 (in Russian).
262. Krokhn, E. M. 1975. Transport of nutrients by salmon migrating from the sea into lakes, pages 153-156. In: A. D. Hasler, (ed.), Coupling of Land and Water Systems. Springer-Verlag. New York, N.Y.
263. La Tourrette, J. 1992. Washington Wildlife Viewing Guide. Falcon Publishing, Inc. Helena, MT. 96 pp.
264. Lagler, K. F., and J. C. Salyer II. 1945. Influence of availability on the feeding habits of the common garter snake. Copeia. 1945. Pages 100-107.
265. Lampman, B. H. 1947. A note on the predaceous habits of water shrews . 1947. J. Mamm. 28:181.
266. Larkin, G. A., and P. A. Slaney. 1997. Implications of trends in marine-derived nutrient influx to south coastal British Columbia salmonid production. Fisheries 22 (11):16-24.
267. Leatherwood, S., and R. R. Reeves. 1983. The Sierra Club handbook of whales and dolphins. Tien Wah Press, Singapore. 302 pp.
268. LeBrasseur, R. J. 1966. Stomach contents of salmon and steelhead trout in the northeastern Pacific Ocean. J. Fish. Res. Bd. Can. 23:85-100.
269. Lee, K. N. 1993. Compass and gyroscope. Integrating science and politics for the environment. Island Press. Washington D. C.
270. Lee, L. C., T. A. Muir, and R. R. Johnson. 1987. Riparian ecosystems as essential habitat for raptors in the American West, pages 15-26. In: Proceedings of the Western Raptor Management Symposium and Workshop. Nat. Wildl. Fed. Washington, D. C.
271. Leider, S. A., M. W. Chilcote, and J. J. Loch. 1986. Movement and survival of presmolt steelhead in a tributary and the main stem of a Washington river. N. Amer. J. Fish. Manag. 6:526-531.
272. Leonard, W. P., H. A. Brown, L. L. C. Jones, K. R. McAllister and R. M. Storm. 1993. Amphibians of Washington and Oregon. Seattle Audubon Society, Seattle, WA. 68 pp.
273. Levy, D. A., and T. G. Northcote. 1982. Juvenile salmon residency in a marsh area of the Fraser River estuary. Can. J. Fish. Aquat. Sci. 39: 270-276.
274. Lichatowich, J. A., and L. E. Mobrand. 1996. Chinook salmon (*Oncorhynchus tshawytscha*) in the Columbia River: The components of decline. In: Applied Ecosystem Analysis - Background - Mobrand Biometrics, Inc., Prepared for US Department of Energy Bonville Power Administration, Environmental Fish and Wildlife. Portland, OR.

275. Lichatowich, J. A. 1999. Salmon without rivers - A history of the Pacific salmon crisis. Island Press, Covelo, Calif. 317 pp.
276. Lichatowich, J. A. 1998. Habitat alteration and changes in habitat of coho (*Oncorhynchus kisutch*) and chinook (*O. tshawytscha*) in Oregon's coastal streams, pages 92-99. In: C. C. Levings, L. B. Holtby, and M. A. Henderson (eds.). Proceedings of the National Workshop on Effects of Habitat Alteration on Salmonid Stocks. Can. Spec. Publ. Fish. Aquat. Sci. 105.
277. Lichatowich, J. A. 1997. Evaluating salmon management institutions: The importance of performance measures, temporal scales, and production cycles, pages 69 to 90. In: D. J. Stouder, P. A. Bisson, and R. J. Naiman (eds.). Pacific Salmon and Their Ecosystems. Chapman and Hall Publishers.
278. Likens, G. E., and F. H. Bormann. 1974. Linkages between terrestrial and aquatic ecosystems. BioScience 24: 447-456.
279. Link, R. M., and K. K. English. 1999. Long-term, sustainable monitoring of Pacific salmon populations using fish wheels to integrate harvesting, management, and research, pages 667-674. In: E. E. Knudsen, C. S. Steward, D. D. MacDonald, J. E. Williams, and D. W. Reiser (eds.). Sustainable Fisheries Management: Pacific Salmon. C. R. C. Lewis Publishers, Boca Raton, FL.
280. Lind, G. S. 1976. Production, nest site selection, and food habits of Ospreys on Deschutes National Forest, Oregon. M.S. Thesis, Oregon State University, Corvallis, OR. 53 pp.
281. Lingle, G. R. 1977. Food habits and sexing-aging criteria of the white pelican at Chase Lake National Wildlife Refuge, North Dakota. M.S. Thesis, Michigan Technological University. 27 pp.
282. Lister, D. B., and H. S. Genoe. 1970. Stream habitat utilization by cohabiting underyearlings of chinook (*Oncorhynchus tshawytscha*) and coho (*Oncorhynchus kisutch*) salmon in the Big Qualicum River, British Columbia. J. Fish. Res. Board Can. 27:1215-1224.
283. Loch, J. J., S. A. Leider, M. W. Chilcote, R. Cooper, and T. H. Johnson. 1988. Differences in yield, emigration-timing, size, and age structure of juvenile steelhead from two small western Washington streams. Calif. Fish and Game 74:106-118.
284. Lowrance, R., R. Todd, J. Fail Jr., O. Hendriksen Jr., R. Leonard, and L. Asmussen. 1984. Riparian forests as nutrient filters in agricultural watersheds. BioScience 34:374-377.
285. Loegering, J. P. 1997. Abundance, habitat association, and foraging ecology of American dippers and other riparian-associated wildlife in the Oregon Coast Range. Ph.D. Dissertation, Oregon State University, Corvallis. 151 pp.
286. Landry, M. R., J. R. Postel, W. K. Peterson, and J. Newman. 1989. Broad-scale distributional patterns of hydrographic variables on the Washington/Oregon shelf, pages 1-40. In: M. R. Landry and B. M. Hickey (eds.). Coastal Oceanography of Washington and Oregon. Elsevier Oceanography Series 47, Amsterdam, Netherlands.
- 286a. Ludwig, D., and C. J. Walters. 1982. Optimal harvesting with imprecise parameter estimates. Ecolog. Model. 14:273-292.
287. MacCarter, D. L. 1972. Food habits of Ospreys at Flathead Lake, Montana. M. S. Thesis. California State University. Humboldt, Arcata. 80 pp.
288. Macdonald, J. S., C. D. Levings, C. D. McAllister, U. H. M. Fagerlund and J. R. McBride. 1988. A Field Experiment to Test the Importance of Estuaries for Chinook Salmon (*Oncorhynchus tshawytscha*) Survival: Short-Term Results. Can. J. Fish. Aquat. Sci. 45:1366-1377.
289. MacDonald, J. S., I. K. Birtwell, and G. M. Kruzynski. 1987. Food and habitat utilization by juvenile salmonids in the Campbell River estuary. Can. J. Fish. Aquat. Sci. 44:1233-1246.

290. Mace, P. M. 1983. Bird Predation on Juvenile Salmonids in the Big Qualicum Estuary, Vancouver Island. *Can. Tech. Rep., Fish. Aquat. Sci.* 1176:1-77.
291. Marcot, B.G., M.A. Castellano, J.A. Christy, L.K. Croft, J.F. Lehmkuhl, R.H. Naney, R.E. Rosentreter, R.E. Sandquist, and E. Zieroth. 1997. Terrestrial ecology assessment, pages 1497-1713. In: T. M. Quigley, S. J. Arbelbide, and S. F. McCool (eds.). *An assessment of ecosystem components in the interior Columbia Basin and portions of the Klamath and Great Basins*, USDA Forest Service General Technical Report PNW-GTR-405. Pacific Northwest Research Station, Portland, Oregon. 1713 pp.
292. Marcot, B. G., and M. Vander Heyden. 2000. Ecological functions of wildlife species. In: D. H. Johnson and T. A. O'Neil (eds.), *Wildlife-Habitat Relationships in Oregon and Washington*. Oregon State University Press, Corvallis.
293. Marquiss, M., and K. Duncan. 1993. Variation in the abundance of Red-breasted Mergansers *Mergus serrator* on a Scottish river related to season, year, river hydrography, salmon density and spring culling. *Ibis*. 135:33-41.
294. Martin, D. J. 1985. Production of cutthroat trout (*Salmo clarki*) in relation to riparian vegetation in Bear Creek, Washington. Ph. D. Dissertation. University of Washington, School of Fisheries. Seattle.
295. Maser, C., R. F. Tarrant, J. M. Trappe, and J. F. Franklin. 1988. From the forest to the sea: A story of fallen trees. USDA- Forest Service, and USDI- Bureau of Land Management. General Technical Report PNW-GTR-229. 153 pp.
296. Maser, C., and J. R. Sedell. 1994. From the forest to the sea - The ecology of wood in streams, rivers, estuaries, and oceans. St. Lucie Press. Delray Beach, Florida. 200 pp.
297. Mathisen, O. A. 1972. Biogenic enrichment of sockeye salmon lakes and stock productivity. *Verh. Int. Ver. Limnol.* 18:1089-1095.
298. Mathisen, O. A., P. L. Parker, J. J. Goering, T. C. Kline, P. H. Poe, and R. S. Scalan. 1988. Recycling of marine elements transported into freshwater systems by anadromous salmon. *Verh. Int. Ver. Limnol.* 23:2249-2258.
299. McHenry, M. L., E. Shott, R. H. Conrad, and G. B. Grette. 1998. Changes in the quantity and characteristics of large woody debris in streams of the Olympic Peninsula, Washington, U. S. A. (1982-1993). *Can. J. Fish. Aquat. Sci.* 55:1395-1407.
300. Matkowski, S. M. D. 1989. Differential susceptibility of tree species of stocked trout to bird predation. *N. Amer. J. Fish. Manag.* 9:184-187.
301. Matkowski, S. M. D. 1984. Angler harvest and other causes of mortality of stocked salmonids in Duck Mountain Provincial Park, Maniotba. M. S. Thesis. University of Manitoba, Winnipeg, Canada.
302. Matthews, D. R. 1983. Feeding ecology of the common murre, *Uria aalge*, off the Oregon coast. M. S. Thesis, University of Oregon, Eugene. 108 pp.
303. Mattson, D. J., B. M. Blanchard and R. R. Knight. 1991. Food habits of Yellowstone Grizzly Bears, 1977-1987. *Can. J. Zoo.* 69:1619-1629.
304. McClelland, B. R. 1973. Autumn concentrations of Bald Eagles in Glacier National Park. *Condor* 75:121-123.
- 304a. McIntyre, J. D., A. R. Hemmingsen and R. C. Simon. 1988. Selection to increase survival of smolts in four successive broods of coho salmon. *Trans. Amer. Fish. Soc.* 117:90-92.
305. McMurray, G. R., and R. J. Bailey. 1998. Change in Pacific Northwest Coastal Ecosystems: Science for Solutions, Pacific Northwest Coastal Ecosystems Regional Study (PNCERS). U. S. Department of Commerce. NOAA Coastal Ocean Program Decision Analysis Series No. 11.

306. McNeil, W. J., J. R. Gowan, and R. Severson. 1991. Offshore release of salmon smolts. American Fisheries Society symposium, 10. Pages 548-553.
307. Meffe, G. K. 1992. Techno-Arrogance and halfway technologies: Salmon hatcheries on the Pacific coast of North America. *Conserv. Biol.* 6(3):350-354.
308. Melone, A. M. 1985. Flood producing mechanisms in coastal British Columbia. *Can. Wat. Res. J.* 10:46-64.
309. Melquist, W. E., and A. E. Dronkert. 1987. River Otter. Wild Furbearer Management and Conservation in North America, pages 626-641. In: M. J. A. Novak, M. E. Obbard, and B. Malloch, (eds.). Ministry of Natural Resources, Ontario.
310. Melquist, W. E., J. S. Whitman and M. G. Hornocker. 1981. Resource partitioning and coexistence of sympatric Mink and River Otter populations. *Proceedings of the Worldwide Furbearer Conference* 1:187-220.
311. Melquist, W. E., and M. G. Hornocker. 1983. Ecology of River Otters in west central Idaho. *Wildlife Monographs* No. 83. Pages 1-60.
312. Melquist, W. E., and D. R. Johnson. 1984. Additional comments on the migration of northern Idaho and eastern Washington Ospreys. *J. Fld. Ornithol.* 55:483-485.
313. Mendall, H. L. 1944. Food of hawks and owls in Maine. *Journal of Wildlife Management.* 8:198-208.
314. Mendall, H. L. 1939. Food habits of the herring gull in relation to freshwater game fishes in Maine. *Wilson Bull.* 41(223-226).
315. Merritt, R. W., and K. W. Cummins. 1996. An introduction to the aquatic insects of North America. Kendall/Hunt Publishing Company. Dubuque, Iowa. 862 pp.
316. Merritt, R. W., K. W. Cummins, and T. M. Burton. 1984. The role of aquatic insects in the processing and cycling of nutrients, pages 134-163. In: V. H. Resh and D. M. Rosenberg (eds.). *The Ecology of Aquatic Insects.* Praeger Publishers. New York, N.Y.
317. Meyer, J. L., and eight co-authors. 1988. Elemental dynamics in streams. *J. N. Am. Benthol. Soc.* 7:410-432.
318. Michael, J. H., Jr. 1995. Enhancement effects of spawning pink salmon on stream rearing juvenile coho salmon: Managing one resource to benefit another. *N.W. Sci.* 69:228-233.
319. Michael, J. H., Jr. 1998. Pacific salmon spawner escapement goals for the Skagit River watershed as determined by nutrient cycling considerations. *N.W. Sci.* 72:239-248.
320. Miegs, R. C., and C. A. Rieck. 1958. Mergansers and trout in Washington. *Proceedings of 47th Annual Conference West Assoc. State Game Fish Comm.* Pages 306-318.
321. Miller, L. 1957. Bird remains from an Oregon Indian Midden. *Condor* 59:59-63.
322. Miller, R. J., and E. L. Brannon. 1982. The origin and development of life history patterns in Pacific Salmon, pages 296 to 309. In E. L. Brannon and E. O. Salo (eds.). *Proceedings of the Salmon and Trout Migratory Behavior Symposium.* June 3-5, 1981, First International Symposium, School of Fisheries, University of Washington, Seattle.
323. Miller, J. A., and C. A. Simenstad. 1997. A comparative assessment of a natural and created estuarine slough as rearing habitat for juvenile chinook and coho salmon. *Estuaries* 20:792-806.
324. Minakawa, N. 1997. The dynamics of aquatic insect communities associated with salmon spawning. Ph. D. Dissertation, University of Washington, School of Fisheries. Seattle.

325. Minshall, G. W., R. C. Petersen, K. W. Cummins, T. L. Bott, J. R. Sedell, C. E. Cushing, and R. L. Vannote. 1983. Interbiome comparison of stream ecosystem dynamics. *Ecol. Mon.* 53:1-25.
326. Minshall, G. W., and J. N. Minshall. 1977. Microdistribution of benthic invertebrates in Rocky Mountain (U.S.A.) Stream. *Hydrobiologia* 55:231-249.
327. Minshall, G. W. 1968. Community dynamics of the benthic fauna in a woodland springbrook. *Hydrobiologia* 32:305-337.
328. Minshall, G. W. 1984. Aquatic insect-substratum relationships. Pages 358-400 in V. H. Resh, and D. M. Rosenberg, editors. *The ecology of aquatic insects*. Praeger Publishers, New York, N.Y.
329. Mitsch, W. J., and J. G. Gosselink. 1986. *Wetlands*. Van Nostrand Reinhold, New York, N.Y. 539 pp.
330. Mizue, K., K. Yoshida, and A. Takemura. 1966. On the ecology of the Dall's porpoise in the Bering Sea and the North Pacific Ocean, [In Japan, English summary]. *Fac. Fish., Nagasaki Univ., Bull.* 21:1-21.
331. Modde, T., A. F. Wasowicz, and D. K. Hepworth. 1996. Cormorant and grebe predation on rainbow trout stocked in a southern Utah reservoir. *North American Journal of Fisheries Management* 16:388-394.
332. Montgomery, D. R., and J. M. Buffington. 1993. Channel classification, prediction of channel response, and assessment of channel condition. Draft report to the Sediment, Hydrology, and Mass Wasting Committee of the Washington State Timber/Fish/Wildlife Agreement. Department of Geological Sciences and Quaternary Research Center, University of Washington. Seattle, WA. 83 pp.
333. Morejohn, G. V., J. T. Harvey and L. T. Krasnow. 1978. The importance of *Loligo opalescens* in the food web of marine vertebrates in Monterey Bay, California. *California Department of Fish and Game Bulletin No. 169*: 67-98.
334. Morrison, M. L., B.G. Marcot, and R.W. Mannan. 1998. *Wildlife-Habitat Relationships*. Second edition. University of Wisconsin Press, Madison, Wisconsin. 435 pp.
335. Mossman, A. S. 1959. Selective predation of glaucous-winged gulls upon adult red salmon. *Ecology* 39:482-486.
336. Moyle, P. 1966. Feeding behavior of the glaucous-winged gull on an Alaskan salmon stream. *Willson Bulletin*. 78:175-190.
337. Muir, W. D., and R. L. Emmett. 1988. The food habits of migrating salmonid smolts passing Bonneville Dam in the Columbia River, 1984. *Re. Riv. Res. Man.* 2:1-10.
338. Mundie, J. H. 1969. Ecological implications of the diet of juvenile coho in streams. Pages 135-152 in T. G. Northcote, (ed.). *Symposium On Salmon and Trout In Streams*, H. R. MacMillan Lectures in Fisheries. Held February 22-24, 1968, at the University of British Columbia, Vancouver.
339. Munn, M. D., R. W. Black, A. L. Haggland, M. A. Hummling, and R. L. Huffman. 1999. An assessment of stream habitat and nutrients in the Elwha River Basin: Implications for restoration. U.S. Geological Survey, Water-Resources Investigations Report 98-4223. Prepared in cooperation with the Lower Elwha Tribe and National Park Service. 38 pp.
340. Munro, J. 1941. Studies of waterfowl in British Columbia. Greater scaup duck, lesser scaup. *Can. J. Res. Section D Zoolog. Sci.* 19:113-138.
341. Munro, J. A. 1932. Food of the American Merganser (*Mergus merganser americanus*) in British Columbia. *Canadian Field-Naturalist* 46:166-168.
342. Munro, J. A., and W. A. Clemens. 1939. The food and feeding habits of the red-breasted merganser in British Columbia. *Journal of Wildlife Management* 3:46-53.

343. Munro, J. A. 1945. Observations of the loon in the Cariboo Parklands, British Columbia. *Auk* 62:38-49.
344. Munro, J. A. 1923. A preliminary report on the relation of various ducks and gulls to the propagation of sockeye salmon at Henderson lake, Vancouver Island. B.C. *Canadian Field-Naturalist* 37:107-116.
345. Munro, J. A. 1938. Studies of waterfowl in British Columbia: Barrow's goldeneye, American goldeneye. *Trans. Royl. Can. Inst.* 22:259-318.
346. Munro, J. A. 1938a. The northern bald eagle in British Columbia. *Wilson Bulletin* 50:38-35.
347. Murie, O. J. 1959. Fauna of the Alutian Islands and Alaska Peninsula. *North American Fauna*. 61 pp.
348. Murota, T. 1998a. Material cycle and sustainable economy. Pages 120-138 *in* D. Bell, L. Fawcett, R. Keil, and P. Penz, editors. *Political Ecology*. Routledge, London-New York.
349. Murota, T. 1998b. Nutrient shadow cast by anadromous fishes: Perspectives in comparison with marine fishery and guano occurrence. Paper presented at the Western Division American Fisheries Society Symposium, held in Anchorage, Alaska (Sept.-Oct., 1998).
350. Murota, T. 1987. The environmental economics of the water planet earth. In: *Environmental Economics - The Analysis of a Major Interface*. Pages 185-199 *in* G. Pilet and T. Murota, editors. R. Leimgruber, Geneva.
351. Murota, T., and Faculty of Environmental Studies. 1994. Material cycle and sustainable economy: A thermodynamical approach to political ecology. A paper read at The Global Political Ecology Conference (Harold Innis Centenary Celebration). Held March 3-6, 1994. York University, Toronto, Canada.
352. Murphy, M. L., and J. D. Hall. 1981. Varied effects of clear-cut logging on predators and their habitat in small streams of the Cascade Mountains, Oregon. *Canadian Journal of Fisheries and Aquatic Sciences* 38:137-145.
353. Myers, G. L., and J. J. Peterka. 1976. Survival and growth of Rainbow trout (*Salmo gairdneri*) in four prairie lakes, North Dakota. *Journal of Fisheries Research Bd. Can.* 33:1192-1195.
354. Myers, K. W. 1980. An investigation of the utilization of four study areas in Yaquina Bay, Oregon, by hatchery and wild juvenile salmonids. M. S. Thesis, Oregon State University. Corvallis, OR.
355. Myers, K. W., and H. F. Horton. 1982. Temporal use of an Oregon estuary by hatchery and wild juvenile salmon. Pages 377-392 *in* V. S. Kennedy, editor. *Estuarine comparisons*. Academic Press, New York, N.Y.
356. Myers, J. M., R. G. Kope, G. J. Bryant, D. Teel, L. J. Lierheimer, T. C. Wainwright, W. S. Grant, F. W. Waknitz, K. Neeley, S. T. Lindley, and R. S. Waples. 1998. Status review of chinook salmon from Washington, Idaho, Oregon, and California. U.S. Dept. of Commerce, NOAA Tech. Memo. NMFS-NWFSC-35. 443pp.
357. Myers, J. P. 1980. Territoriality and flocking by Buff-breasted Sandpipers: variations in non-breeding dispersion. *Condor* 82:241-250.
358. Nagorsen, D. W., K. F. Morrison and J. E. Forsberg 1989. Winter diet of Vancouver Island marten (*Martes americana*). *Canadian Journal of Zoology* 67:1394-1400.
359. Nagorsen, D. W., W. R. Campbell and G. R. Giannico 1989. Winter Food Habits of Marten, *Martes americana*, on the Queen Charlotte Islands. *Canadian Field-Naturalist* 105:55-59.
360. Naiman, R. J., and E. C. Anderson. 1997. Streams and rivers: their physical and biological variability. Pages 131-148 (Chap. 7) *in* P. Schoemaker, B. von Hagen, and E. Wolf, editors. *The Rain Forests of Home: Profile of a North American Bioregion*. Ecotrust/Interain Pacific and Island Press. 480 pp.

361. Naiman, R. J., T. J. Beechie, L. E. Benda, P. A. Bisson, L. H. MacDonald, M. D. O'Connor, C. Oliver, P. Olson, and E. A. Steel. 1992. Fundamental elements of ecologically healthy watersheds in the Pacific Northwest Coastal Ecoregion. Pages 127-188 *in* R. J. Naiman, editor. *Watershed Management - Balancing Sustainability and Environmental Change*. Springer-Verlog. New York, N.Y.
362. Naiman, R. J., and R. E. Bilby. 1998. *River ecology and management: Lessons from the Pacific Coastal Ecoregion*. Springer-Verlag, New York, N.Y. 705 pp.
363. Nakatani, R. E. 1969. Effects of heated discharge on anadromous fish. *In* P. A. Krenkel, and F. L. Parker, editors. *Biological aspects of thermal pollution*. Vanderbilt University Press. Nashville, TN.
364. National Marine Fisheries Service (NMFS). 1997. Investigation of scientific information on the impacts of California sea lions and Pacific harbor seals on salmonids and on the coastal ecosystems of Washington, Oregon, and California. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-NWFSC-28. 172 pp.
365. National Research Council (NRC). 1996. *Upstream - Salmon and society in the Pacific Northwest*. National Academy Press. Washington D. C. 452 pp.
366. National Marine Fisheries Service (NMFS). 1998. *Essential Fish Habitat Advisory Report to the Pacific Fishery Management Council for Pacific salmon species of California, Oregon, and Washington*. Draft report prepared by the Pacific State Marine Fisheries Commission. Gladstone.
- 366a. National Marine Fisheries Service (NMFS). 1991. Final rule for endangered status of Snake River sockeye salmon. November 20, 1991. *Federal Register* 56: 58619.
367. Nehlsen, W., J. E. Williams, and J. A. Lichatowich. 1991. Pacific salmon at the crossroads: stocks at risk from California, Oregon, Idaho, and Washington. *Fisheries* 16 (2):4-21.
368. Netboy, A. 1980. *The Columbia River salmon and steelhead trout: Their fight for survival*. University of Washington Press, Seattle, WA.
369. Newbold, J. D., J. W. Elwood, R. V. O'Neil, and W. V. Winkle. 1981. Measuring nutrient spiralling in streams. *Canadian Journal of Fisheries and Aquatic Sciences* 38:860-863.
370. Newton, M., B. A. El Hassen, and J. Zavitkovski. 1968. Role of red alder in western Oregon forest succession. Pages 73-84 *in* J.M. Trappe, J.F. Franklin, R.F. Tarrant, and G.M. Hansen, editors. *Biology of alder*. USDA Forest Service. Portland, OR.
371. Nichol, L. M., and D. M. Shackleton. 1996. Seasonal movements and foraging behavior of northern resident killer whales (*Orcinus orca*) in relation to the inshore distribution of salmon (*Oncorhynchus* spp.) in British Columbia. *Canadian Journal of Zoology* 74:983-991.
372. Nicholas, J. W., and D. G. Hankin. 1988. Chinook salmon populations in Oregon coastal river basins: Description of life histories and assessment of recent trends in run strengths. *Oreg. Department of Fish and Wildlife, Fisheries Division. Info. Rep., No. 88-1*. 359 pp.
373. Nickelson, T. E. 1986. Influences of upwelling, ocean temperatures, and smolt abundance on marine survival of coho salmon (*Oncorhynchus kisutch*) in the Oregon Production Area. *Canadian Journal of Fisheries and Aquatic Sciences* 43:527-535.
374. Nickelson, T., J. Rodgers, S. Johnson, and M. Solazzi. 1992. Seasonal changes in habitat use by juvenile coho salmon (*Oncorhynchus kisutch*) in Oregon Coastal Streams. *Canadian Journal of Fisheries and Aquatic Sciences* 49:783-789.
375. Nicola, S. J. 1966. The relationship of Alloverla nymphs to incubating pink (*Oncorhynchus gorbuscha*) and chum (*Oncorhynchus keta*) salmon eggs, alevins, and pre-emergent fry in a southeastern Alaska stream. M. S. Thesis, University of Washington, School of Fisheries. Seattle, WA.

376. Nicola, S. J. 1968. Scavenging by *Alloperla* (Plecoptera: Chloroperlidae) nymphs on dead pink (*Oncorhynchus gorbuscha*) and chum (*Oncorhynchus keta*) salmon embryos. *Canadian Journal of Zoology* 46:787-796.
377. Nikol'skii, G. V. 1952. The type of dynamics of stocks and the character of spawning of the chum (*Oncorhynchus keta*, Walb.) and the pink salmon (*Oncorhynchus gorbuscha*, Walb.) in the Amur River. *Doklady Akademii Nauk. SSSR*, 86(4): 873-875
378. Norris, K. S., and J. H. Prescott. 1961. Observations on Pacific cetaceans of California and Mexican waters. University of California. *Publ. Zoo.* 63:291-403.
- 378a. North, M. R. 1994. Yellow-billed loon, *Gavia adamsii*. *The Birds of North America* No.121. In A. Poole, and F. Gill, editors. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologist's Union, Washington, D.C.
379. Northwest Power Planning Council (NPPC). 1986. Compilation of information on salmon and steelhead losses in the Columbia River basin. Appendix D of the 1987 Columbia River Basin Fish and Wildlife Program. Portland, OR. 252 pp.
380. O'Connell, M. A., J. G. Hallett, and S. D. West. 1993. Wildlife use of riparian habitats: A Literature review. Washington Department of Natural Resources, Timber/Fish/Wildlife, Report TFW-WL1-93-001.
381. Oakley, A. L., J. A. Collins, L. B. Everson, D. A. Heller, J. C. Howerton, and R. E. Vincent. 1985. Riparian zones and freshwater wetlands. Pages 58-76 in R. E. Brown, editor. *Management of Wildlife and Fish Habitats in Forests of Western Oregon and Washington*. U. S. For. Serv. Portland, OR.
382. Obermayer, K. E., A. Hodgson, and M. F. Willson. 1999. American Dipper, *Cinclus mexicanus*, foraging on Pacific salmon, *Oncorhynchus* spp., eggs. *Canadian Field-Naturalist* 113:288-290.
383. Ofelt, C. H. 1975. Food habits of nesting bald eagles in southeast Alaska. *Condor* 77:337-338.
384. Olesiuk, P. F. 1993. Annual prey consumption by harbor seal (*Phoca vitulina*) in the Strait of Georgia, British Columbia. *Fish. Bull.* 91:491-515.
385. Olson, R., and W. Hubert. 1994. *Beaver: Water resources and Riparian Habitat Manager*. University of Wyoming. Laramie, WY. 48 pp.
386. Ordal, E., and R. E. Pacha. 1963. The effects of temperatures on disease in fish. In *Water temperature: Influences, effects and control*. Proceedings of the twelfth Pacific Northwest symposium on water pollution research. U.S. Public Health Service, Pacific Northwest Water Laboratory. Corvallis, OR.
387. Oregon Coastal Salmon Restoration Initiative (OCSRI). 1997. OCSRI Conservation Plan. Draft Revision February 24, 1997. Salem, OR.
- 387a. Oregon Department of Fish and Wildlife (ODFW). 1986. Comprehensive plan for production and management of Oregon's anadromous salmon and trout. Part III. Steelhead Plan. Oregon Department of Fish and Wildlife, Division of Anadromous Fish Section. Portland, OR.
- 387b. Oregon Department of Fish and Wildlife. 1981. Oregon's commercial harvest of Coho salmon, *Oncorhynchus kisutch* (Walbaum), 1892-1960. Information Report Series, Fisheries. Number 81-3.
388. Pacific Fishery Management Council (PFMC). 1998. Review of 1997 ocean salmon fisheries. A Report of the Pacific Fishery Management Council pursuant to National Oceanic and Atmospheric Administration, Award No. NA87FC0008. Portland, OR.
- 388a. Pacific Fishery Management Council (PFMC). 1991. Review of 1990 ocean salmon fisheries. Portland, OR.

389. Palmer. 1962. Handbook of North American Birds 1. Yale University Press. New Haven, Connecticut.
390. Palmisano, J. F., R.H. Ellis, and V. W. Kaczynski. 1993. The impact of environmental and management factors on Washington's wild anadromous salmon and trout. Washington Forest Protection Association and State of Washington Department of Natural Resources. Olympia. 371 pp.
391. Parker, M. S. 1994. Feeding ecology of stream-dwelling Pacific giant salamander larvae (*Dicamptodon tenebrosus*). *Copeia* 3:705-718.
392. Parker, M. S. 1991. Relationship between cover availability and larval Pacific giant salamander density. *J. Herp.* 25:355-357.
393. Parker, M. S. 1993. Predation by Pacific giant salamander larvae on juvenile steelhead trout. *Northwest Naturalist* 74:77-81.
394. Percy, W. G. 1997. What have we learned in the last decade? What are research priorities? Pages 271-277 in R.L. Emmett, and M.H. Schiewe, editors. Estuarine and Ocean Survival of Northeastern Pacific Salmon, NOAA Technical Memorandum NMFS-NWFSC-29.
395. Percy, W. G., R. D. Brodeur, and J. P. Fisher. 1990. Distribution and ecology of juvenile cutthroat trout (*Oncorhynchus clarki clarki*) and steelhead (*O. mykiss*) in the ocean off Oregon and Washington. *Fisheries Bulletin* 88:697-711.
396. Percy, W. G., R. D. Brodeur, J. M. Shenker, W. W. Smoker, and Y. Endo. 1988. Food habits of Pacific salmon and steelhead trout, midwater trawl catches and oceanographic conditions in the Gulf of Alaska, 1980-1985. *Bull. Ocean Res. Instit., Univ. of Tokyo* 26(2):29-78.
397. Percy, W. G. 1992. Ocean ecology of north Pacific salmonids. Washington Sea Grant Program, distributed by University of Washington Press. Seattle, WA.
398. Peterson, N. P. 1980. The role of spring ponds in the winter ecology and natural production of coho salmon (*Oncorhynchus kisutch*) on the Olympic Peninsula, Washington. M. S. Thesis, University of Washington, College of Fisheries. Seattle. 96 pp.
399. Peterson, N. P. 1982a. Population characteristics of juvenile coho salmon (*Oncorhynchus kisutch*) overwintering in riverine ponds. *Canadian Journal of Fisheries and Aquatic Sciences* 39:1303-1307.
400. Peterson, N. P. 1982b. Immigration of juvenile coho salmon (*Oncorhynchus kisutch*) into riverine ponds. *Canadian Journal of Fisheries and Aquatic Sciences* 39:1308-1310.
401. Peterson, N. P., and L. M. Reid. 1984. Wall-base channels: their evolution, distribution, and use by juvenile coho salmon in the Clearwater River, Washington. Pages 215-226 in J.M. Walton and D.B. Houston, editors. Proceedings of the Olympic Wild Fish Conference, March 23-25, 1983, Port Angeles, WA.
- 401a. Peterson, N. P., and T. P. Quinn. 1996. Persistence of egg pocket architecture in redds of chum salmon, *Oncorhynchus keta*. *Envir. Biol. Fish.* 46:243-253.
402. Phinney, L. A., and P. Bucknell. 1975. A catalog of Washington streams and salmon utilization. Washington Department of Fisheries, Olympia, WA.
403. Pike, G. C. 1950. Stomach contents of whales caught of the coast of British Columbia. Progress Report of the Pacific Coast Stations. Fisheries Research Board of Canada 83:27-28.

404. Piorkowski, R. J. 1995. Ecological effects of spawning salmon in several southcentral Alaskan streams. Ph. D., University of Alaska, Fairbanks. 177 pp.
405. Poage, N. J., and T. A. Spies. 1996. A tale of two unmanaged riparian forests. COPE Report 9(1):6-9.
406. Poelker, R.J., and H.D. Hartwell. 1973. Black bear of Washington. Washington State Game Department. Biological Bulletin No. 14. Olympia, Washington. 180 pp.
407. Pitcher, K. W. 1981. Prey of the Steller sea lion, *Eumetopias jubatus*, in the Gulf of Alaska. U. S. Fish and Wildlife Service Fishery Bulletin 79:467-472.
410. Pitcher, K. W. 1977. Population productivity and food habits of harbor seals in the Prince William Sound, Copper River Delta Area, Alaska. Pages 1-20 in Report to U.S. Marine Mammal Commission for Contract MM5AC01.
411. Pitcher, K. W., and D. G. Calkins. 1991. Reproductive biology of Steller sea lions in the Gulf of Alaska. Journal of Mammalogy 62:599-605.
412. Pomery, L. R. 1970. The strategy of mineral cycling. Annu. Rev. Ecol. Syst. 1:171-190.
413. Pritchard, D. W. 1967. What is an estuary: Physical viewpoint. Pages 3-5 in G. H. Lauff, editor. Estuaries. Am. Assoc. Adv. Sci. Pub. 83, Washington, D.C.
414. Quinn, T. P., and N. P. Peterson. 1994. The effects of forestry practices on fish populations: Incubation Environment of Chum Salmon (*Oncorhynchus keta*) in Kennedy Creek - Part A. Persistence of Egg Pocket Architecture in Chum Salmon Redds. Washington Department of Natural Resources, Olympia, WA. Timber/Fish/Wildlife Report. TFW-FA-94-001.
415. Rader, R. B. 1997. A functional classification of the drift: traits that influence invertebrate availability to salmonids. Canadian Journal of Fisheries and Aquatic Sciences 54:1211-1234.
416. Ralph, S. C., G. C. Poole, L. L. Conquest, and R. J. Naiman. 1994. Stream channel morphology and woody debris in logged and unlogged basins of western Washington. Canadian Journal of Fisheries and Aquatic Sciences 51:37-51.
417. Reeves, G. H., P. A. Bisson, J. M. Dambacher. 1998. Fish Communities. Pages 200-234 in R. J. Naiman and R. E. Bilby, editors. River Ecology and Management: Lessons From the Pacific Coastal Ecoregion. Springer-Verlag. New York, N.Y.
418. Reeves, G. H., F. H. Everest, and J. D. Hall. 1987. Interactions between the reidside shiner (*Richardsonius balteatus*) and the steelhead trout (*Salmo gairdneri*) in western Oregon: the influence of water temperature. Canadian Journal of Fisheries and Aquatic Sciences 44:1602-1613.
419. Regier, H. A. 1997. Old traditions that led to abuses of salmon and their ecosystems. Pages 17-28 in D. J. Stouder, P. A. Bisson, and R. J. Naiman, editors. Pacific Salmon and Their Ecosystems: Status and Future Options. Chapman & Hall. New York, N.Y.
420. Reice, S. R. 1994. Nonequilibrium determinants of biological community structure. American Scientist 82:424-435.
421. Reid, L. M. 1981. Sediment production from gravel-surfaced forest roads, Clearwater Basin, Washington. M. S. Thesis, University of Washington Department of Geological Sciences. Seattle, WA. 350 pp.
422. Reimchen, T. E. 1994. Further studies of predator and scavenger use of chum salmon in stream and estuarine habitats at Bag Harbour, Gwaii Haanas. Island Ecological Research, Queen Charlotte City, B.C., Prepared for Canadian Parks Service. 58 pp.
423. Reimer, S. D., and R. F. Brown. 1997. Prey of pinnipeds at selected sites in Oregon identified by scat (fecal) analysis, 1983-1996. Oregon Department of Fish and Wildlife, Wildlife Diversity Program, Technical Report No. 97-6-02.

34 pp.

424. Reinhart, D. P., and D. J. Mattson. 1989. Bear use of cutthroat trout spawning streams in Yellowstone National Park. *International Conference of Bear Research and Management* 8:343-350.

424a. Read, L. B. 1997. A Bayesian decision theory approach to harvest management of salmon and steelhead. Report prepared for Washington Trout, P. O. Box 402. Duvall, WA 98019.

425. Richey, J. E., M. A. Perkins, and C. R. Goldman. 1975. Effects of kokanee salmon (*Oncorhynchus nerka*) decomposition on the ecology of a subalpine stream. *J. Fish. Res. Bd. Can.* 32:817-820.

425a. Ricker, W. E. 1954. Stock and recruitment. *J. Fish. Res. Bd. Can.* 11:559-623.

426. Ricker, W. 1980. Causes of the decrease in age and size of chinook salmon (*Oncorhynchus tshawytscha*). *Can. Tech. Rep. Aquat. Sci.* 944

427. Robertson, I. 1973. Predation by fish-eating birds on stocks of Pacific herring, *Clupea pallasii*, in the Gulf Islands of British Columbia. Unpublished Report, Pac. Biol. Stn., Nanaimo, BC. 29 pp.

428. Robertson, I. 1974. The food of nesting double-crested and pelagic cormorants at Mandarte Island, British Columbia, with notes on feeding ecology. *Condor* 76:346-348.

429. Roby, D. D., D. P. Craig, K. Collis, and S. L. Adamany. 1998. Avian predation on juvenile salmonids in the lower Columbia River - 1997 Annual Report. Oregon Coop. Wildl. Res. Unit., Oregon State University, Corvallis, OR. 69 pp.

430. Roby, D. D., D. P. Craig, K. Collis, and S. L. Adamany. 1998. (Unpublished) Avian predators on juvenile salmonids in the lower Columbia River. Annual Report to Bonneville Power Administration and U. S. Army Corps of Engineers. 87 pp.

431. Roche, J., and M. McHutchison. 1998. First fish/first people: Salmon tales of the North Pacific rim. University of Washington Press. Seattle, WA. 199 pp.

432. Roemmich, D., and J. McGowan. 1995. Climatic warming and the decline of zooplankton in the California Current. *Science* 267:1324-1326.

433. Roffe, T. J., and B. R. Mate. 1984. Abundances and feeding habits of pinnipeds in the Rogue River, Oregon. *Journal of Wildlife Management* 48:1262-1274.

434. Rosgen, D. 1996. Applied river morphology. Wildland Hydrology. Pagosa Springs, CO.

435. Ruttner, F. 1971. Fundamentals of limnology. English translation of Third Edition Copyright University of Toronto Press. Toronto, Canada.

436. Rowlett, R. A. 1980. Observations of marine birds and mammals in the northern Chesapeake Bight. U. S. Fish Wildl. Serv., Biol. Serv. Prog., FWS/OBS-80/04.

437. Ruggerone, G. T. 1986. Consumption of migrating juvenile salmonids by gulls foraging below a Columbia River dam. *Transactions of American Fisheries Society* 115:736-742.

438. Ryall, R., and C. D. Levings. 1987. Juvenile salmon utilization of rejuvenated tidal channels in the Squamish estuary, British Columbia. *Can. Manuscript Rep. of Fisheries Aquat. Sci.* 1904.

439. Salo, E. O., and W. H. Bayliff. 1958. Artificial and natural production of silver salmon, *Oncorhynchus kisutch*, at Minter Creek, Washington. State of Washington Department of Fisheries. Res. Bull. No. 4.

440. Salo, E. O. 1991. Life history of chum salmon. Pages 231-307 in C. Groot, and L. Margolis, editors. Pacific salmon life

histories, University of British Columbia Press. Vancouver, British Columbia, Canada.

441. Salyer, J. C., and K. F. Lagler. 1940. The food and habits of the American merganser during winter in Michigan, considered in relation to fish management. *Journal of Wildlife Management* 4:186-219.
442. Sandercock, F. K. 1991. Life history of coho salmon (*Oncorhynchus kisutch*). Pages 397 to 445 in C. Groot and L. Margolis, editors. *Pacific Salmon Life Histories*, University of British Columbia Press. Vancouver.
- 442a. Sanger, G. A. 1983. Diets and Food Web Relationships of Seabirds in the Gulf of Alaska and Adjacent Marine Regions. Pages 631-771 in *Final Report Outer Continental Shelf Environmental Assessment Program*, US Fish and Wildlife Service Denver Wildlife Research Center Migratory Bird Project.
443. Schlorff, R. W. 1978. Predatory ecology of the Great Egret at Humboldt Bay, California. Pages 347-353 in A. J. Sprunt, J. C. Ogden, and S. Winkler, editors. *Wading Birds*. National Audubon Society Research Report No. 7.
444. Scheffer, V. B., and J. W. Slipp. 1944. The harbor seal in Washington State. *American Naturalist* 32:374-416.
445. Scheffer, V. B. 1950. The food of the Alaska fur Seal. United States Dept. of the Interior, Fish and Wildlife Service. *Wildlife Leaflet No. 329*. Washington D.C. 16 pp.
446. Scheffer, T. H., and C. C. Sperry. 1931. Food habits of the Pacific Harbor Seal, *Phoca richardii*. *Journal of Mammalogy* 12:214-226.
447. Schwartz, J. E. II, and G. E. Mitchell. 1945. The Roosevelt elk on the Olympic Peninsula, Washington. *Journal of Wildlife Management* 9:295-319.
448. Scott, J. M. 1973. Resource allocation in four synotic species of marine diving birds. Ph. D. Dissertation, Oregon State University, Corvallis, OR. 107 pp.
449. Scott, W. B., and E. J. Crossman. 1973. *Freshwater fishes of Canada*. Fish. Res. Bd. Can., Bull. 184. 966 pp.
450. Schuldt, J. A., and A. E. Hershey. 1995. Effect of salmon carcass decomposition on Lake Superior tributary streams. *J. N. Amer. Benth. Soc.* 14 (2) 259-268.
451. Sedell, J. R., F. J. Triska, and N. S. Triska. 1975. The processing of conifer and hardwood leaves in two coniferous forest streams. 1. Weight loss and associated invertebrates. *Verh. Int. Ver. Limnol.* 19:1617-1627.
452. Sedell, J. R., R. J. Naiman, K. W. Cummins, G. W. Minshall, and R. L. Vannote. 1979. Transport of particulate organic material in streams as a function of physical processes. *International. Verh. Int. Ver. Limnol.* 20:1366-1375.
453. Sedell, J. R., and K. J. Luchessa. 1981. Using the historical record as an aid to salmonid habitat enhancement. Pages 210-223 in N. B. Armantrout, editor. *Acquisition and Utilization of Aquatic Habitat Information*. Proceedings of a Symposium held 28-30 October, 1981. Portland, OR.
454. Seliskar, D. M., and J. L. Gallagher. 1983. The ecology of tidal marshes of the Pacific Northwest coast: a community profile. USDI - Fish and Wildlife Service, Division of Biological Services. FWS/OBS-82/32. Washington, D.C.
455. Senn, H. G. 1958. Merganser degradation. Unpublished Report to Fisheries Management Division, State of Washington Department of Game. Olympia, WA.
456. Serdar, C. F. 1999. Description, analysis, and impacts of the Grouse Creek Landslide, Jefferson County, Washington, 1997-98. Master of Environmental Studies, The Evergreen State College. Olympia, WA. 171 pp.
457. Servheen, C. W. 1975. Ecology of the wintering bald eagle on the Skagit River, Washington. M. S. Thesis. University of Washington, Seattle, WA.

458. Seufert, F. 1980. *Wheels of fortune*. Oregon Historical Society. 259 pp.
459. Shea, D. S. 1973. White-tailed deer eating salmon. *Murrelet* 54:23.
460. Shreffler, D. K., and R. M. Thom. 1993. *Restoration of Urban Estuaries: New approaches for site location and design*. Prepared for the Washington State Department of Natural Resources by Battelle/Marine Sciences Laboratory. Sequim, WA.
461. Shreffler, D. A., C. A. Simenstad, and R. M. Thom. 1992. Temporary residence of juvenile salmon in a restored estuarine wetland. *Canadian Journal of Fisheries and Aquatic Sciences* 47:2079-2084.
462. Shuldt, J. A., and A. E. Hershey. 1995. Effect of salmon carcass decomposition on Lake Superior tributary streams. *Journal of Wildlife Management* 14:1-9.
463. Sibatani, A. 1992. Naze sake ha kawa wo sozyousuruka (Why do salmon go up rivers). In: Tyuuou Kouron/Chuo Koron (Tokyo) 1992(4): 286-295 (In Japanese).
464. Sibatani, A. 1996. (English translation by: R. Davis). Why do salmon ascend rivers? *Selected Papers on Entropy Studies* 3:3-11.
465. Simenstad, C. A., W. J. Kinney, S. S. Parker, E. O. Salo, J. R. Cordell, and H. Buechner. 1980. Prey community structures and trophic ecology of outmigrating juvenile chum and pink salmon in Hood Canal, Washington: A synthesis of three years' studies, 1977-1979. Final Rep. to Washington Department of Fisheries. Fisheries Research Institute, University of Washington. FRI-UW-8026. Seattle, WA. 113 pp.
466. Simenstad, C. A., K. L. Fresh, and E. O. Salo. 1982. The role of Puget Sound and Washington coastal estuaries in the life history of Pacific salmon: An unappreciated function. Pages 343-364 in V. S. Kennedy, editor. *Estuarine Comparisons*. Academic Press. New York, N.Y. 709 pp.
467. Simenstad, C. A., and E. O. Salo. 1982. Foraging success as a determinant of estuarine and nearshore carrying capacity of juvenile chum salmon (*Oncorhynchus keta*) in Hood Canal, Washington. Pages 21-37 in B. R. Melteff, and R. A. Neve, editors. *Proceedings of the North Pacific Aquaculture Symposium, 18-27 August 1980*. Anchorage, Alaska and Newport, Oregon. Alaska Sea Grant Rep. 82-2. Univ. Alaska, Fairbanks. 372 pp.
468. Simenstad, C. A. B. S. Miller C. F. Nyblade K. Thornburgh and L. J. Bledsoe. 1979. Food web relationships of northern Puget Sound and the Strait of Juan de Fuca. MESA Puget Sound Project, Environ. Res. Labs. DOC/EPA Interagency Energy /Environment R&D Program Rep., EPA-600/7-79-259.
469. Simenstad, C. A., M. Dethier, C. Levings, and D. Hay. 1997b. The Land-Margin Interface of Coastal Temperate Rain Forest Ecosystems: Shaping the Nature of Coastal Interactions. Pages 149-187 (Chap. 7) in P. Schoonmaker, B. von Hagen, and E. Wolf, editors. *The Rain Forests of Home: Profile of a North American Bioregion*. Ecotrust/Interain Pacific and Island Press. 480 pp.
470. Simenstad, C. A., J. R. Cordell, W. G. Hood, B. E. Feist, and R. M. Thom. 1997a. Ecological status of a created estuarine slough in the Chehalis River estuary: Assessment of created and natural estuarine sloughs, January-December 1995., Fisheries Research Institute, School of Fisheries, FRI-UW-9621, University of Washington, Seattle, WA. 47 pp.
471. Simenstad, C. A., J. R. Cordell, J. A. Miller, W. G. Hood, and R. M. Thom. 1993. Ecological status of a created estuarine slough in the Chehalis River estuary: Assessment of created and natural estuarine sloughs, January-December 1992. Fisheries Research Institute, University of Washington. FRI-UW-9305. Seattle, WA. 56 pp.
472. Simenstad, C. A., J. R. Cordell, W. G. Hood, J. A. Miller, and R. M. Thom. 1992. Ecological status of a created estuarine slough in the Chehalis River estuary: Report of monitoring in created and natural estuarine sloughs, January-

- December 1991. Fisheries Research Institute, University of Washington. FRI-UW-9206. Seattle, WA. 49 pp.
473. Simenstad, C. A., W. G. Hood, R. M. Thom, D. A. Levy, and D. L. Bottom. In press. Landscape structure and scale constraints on restoring estuarine wetlands for Pacific Coast juvenile fishes. *In* M. P. Weinstein, and D. A. Kreeger, editors. *Concepts and Controversies in Tidal Marsh Ecology*, Kluwer Academic Publ., Dordrecht.
474. Simons, D. D. 1983. Interactions between California condors and humans in prehistoric far western North America. Pages 470-494 *in* S.R. Wilbur, and J. A. Jackson, editors. *Vulture Biology and Management*. University of California Press, Berkeley, Los Angeles & London.
475. Skagen, S. K., R. L. Knight, and G. H. Orians. 1991. Human disturbance of an avian scavenging guild. *Ecological Applications* 1:215-225.
476. Slaney, P. A., B. O. Rublee, C. J. Perrin, and H. Goldberg. 1994. Debris structure placements and whole-river fertilization for salmonids in a large regulated stream in British Columbia. *Bull. Mar. Sci.* 55:1160-1180.
477. Slaney, P. A., and D. Zaldokas. 1997. Fish habitat rehabilitation procedures. *Watershed Restoration Technical Circular No. 9*. Watershed Restoration Program, Ministry of Environment, Lands and Parks, Vancouver, British Columbia.
478. Smith, C. L. 1979. *Salmon fishers of the Columbia*. Oregon State University Press, Corvallis, OR.
479. Smith, M. R., P. W. Mattocks, Jr., and M. K. Cassidy. 1997. Breeding birds of Washington State. *In* K. M. Cassidy, C. E. Grue, M. R. Smith, and K. M. Dvornich, editors (eds.). Vol. 4. *Washington State GAP Analysis - Final Report*. Seattle Audubon Society, Publications in Zoology, No. 1. Seattle, WA. 538 pp.
480. Smith, J. L., and D. R. Mudd. 1978. Food of the Caspian tern in Grays Harbor, Washington. *Murrelet* 59:105-106.
481. Smith, M. W. 1968. Fertilization and predator control to increase growth rate and yield of trout in a natural lake. *J. Fish. Res. Bd. Can.* 25:2011-2036.
482. Spalding, D. J. 1964. Comparative feeding habits of fur seal, sea lion, and harbor seal on the British Columbia coast. *Fish. Res. Bd. Can. Bull. No. 146*. 52 pp.
483. Spanier, E. 1980. The use of distress calls to repel night herons (*Nycticorax nycticorax*) from fish ponds. *J. Appl. Ecol.* 17:287-294.
484. Speaker, R.W., K. J. Luchessa, J. F. Franklin, and S.V. Gregory. 1988. The use of plastic strips to measure leaf retention by riparian vegetation in a coastal Oregon stream. *American Midland Naturalist* 120:22-31.
485. Spence, B. C., G. A. Lomnicky, R. M. Hughes, and R. P. Novitzki. 1996. An ecosystem approach to salmonid conservation. Report prepared by ManTech. TR-4501-96-6057. Corvallis, OR.
486. Spencer, C. N., B. R. McClelland, and J. A. Stanford. 1989. Shrimp stocking, salmon collapse and eagle displacement: cascading interactions in the food web of a large aquatic ecosystem. *BioScience* 41:14-21.
487. Stalmaster, Mark V. 1980. Salmon carrion as a winter food source for red-tailed hawks. *Murrelet* 61(1):43-44.
488. Stalmaster, M. V., J. R. Newman and A. J. Hansen. 1979. Population dynamics of wintering bald eagles on the Nooksack River, Washington. *Northwest Science* 53:126-131.
489. Stalmaster, M. V. 1976. Winter ecology and effects of human activity on bald eagles in the Nooksack River valley, Washington. M. S. Thesis. Western Washington State College, Bellingham, WA.
490. Stalmaster, M. V., and J. A. Gessaman. 1984. Ecological energetics and foraging behavior of overwintering bald eagles. *Ecol. Monog.* 54:407-428.

491. Stanford, J. A., and J. V. Ward. 1988. The hyporheic habitat of river ecosystems. *Nature* 335:64-66.
492. Stanford, J. A., and A. R. Gaufin. 1974. Hyporheic communities of two Montana rivers. *Science* 185:700-702.
493. Steeger, C., H. Esselink, and R. C. Ydenberg. 1992. Comparative feeding ecology and reproductive performance of ospreys in different habitats of southeastern British Columbia. *Canadian Journal of Zoology* 70:470-475.
494. Stenson, G. B., G. A. Badgero, and H. D. Fisher. 1984. Food habits of the River otter, *Lutra canadensis*, in the marine environment of British Columbia. *Canadian Journal of Zoology* 62:88-91.
495. Stewart, B. S., and S. Leatherwood. 1985. Minke whale. *Balaenoptera acutorostrata Lacepede*, 1804. Pages 91-136 in S. H. Ridgway, and R. Harrison, editors. *Handbook of Marine Mammals. Volume 3. The Sirenians and Baleen Whales.* Academic Press, London, Orlando, FL.
496. Stockner, J. G., and E. A. MacIsaac. 1996. British Columbia lake enrichment programme: two decades of habitat enhancement for sockeye salmon. *Regulated Rivers: Research & Management* 12:547-561.
497. Stockner, J. G., and K. R. S. Shortreed. 1978. Enhancement of autotrophic production by nutrient addition in a coastal rainforest stream on Vancouver Island. *J. Fish. Res. Bd. Can.* 35:28-34.
498. Stolz, J., and J. Schnell. 1991. Trout - The Wildlife Series. Stackpole Books. Cameron and Kelker Streets, Harrisberg, PA. 370 pp.
499. Stouder, D. J., P. A. Bisson, and R. J. Naiman, editors. 1997. Pacific salmon and their ecosystems: Status and future options. I. T. P. Chapman and Hall International Thomson Publishing. New York, N.Y.
500. Stroud, R. K., C. H. Fiscus and H. Kajimura. 1980. Food of the Pacific white-sided dolphin, *Lagenorhynchus obliquidens*, Dall's Porpoise, *Phocoenoides dalli*, and Northern fur seal, *Callorhinus ursinus*, off California and Washington. U. S. Fish and Wildlife Service Fish. Bull. 78(4):951-959.
501. Strub, P. T., C. James, A. C. Thomas, and M. R. Abbott. 1990. Seasonal and nonseasonal variability of satellite-derived surface pigment concentration in the California Current. *J. Geophys. Res.* 95(C7):11503-11530.
502. Suckley, G., and J. G. Cooper. 1860. The natural history of Washington Territory and Oregon. Bailliere Brothers, New York. 399 pp.
503. Sugai, S. F., and D. C. Burrell. 1984. Transport of dissolved organic carbon, nutrients, and trace metals from the Wilson and Blossom rivers to Smeaton Bay, southeast Alaska. *Canadian Journal of Fisheries and Aquatic Sciences* 41:180-190.
504. Sullivan, K., J., Tooley, K. Doughty, J.E. Caldwell, and P. Knudsen. 1990. Evaluation of prediction models and characterization of stream temperature regimes in Washington. Washington Department of Natural Resources. Timber/Fish/Wildlife Rep. No. TFW-WQ3-90-006. Olympia, WA.
505. Sumner, F. H. 1962. Migration and growth of coastal cutthroat trout in Tillamook County, Oregon. *Transactions of American Fisheries Society* 91:77-83.
506. Swanson, F. J., S. V. Gregory, J. R. Sedell, and A.G. Campbell. 1982. Land-water interactions: The riparian zone. Pages 267-291 in R.L. Edmonds, editor. *Analysis of coniferous forest ecosystems in the western United States.* U.S. Int. Biolog. Prog. Synthesis Series 14, Hutchinson Ross. Stroudsburg, PA.
507. Swanson, F. J., L. E. Benda, S. H. Duncan, G. E. Grant, W. F. Megahan, L. M. Reid, and R. R. Ziemer. 1987. Mass failures and other processes of sediment production in Pacific Northwest forest landscapes. Pages 9 to 38 in E. O. Salo, and T. W. Cundy, editors. *Streamside Management: Forestry and Fishery Interactions.* College of Forest Resources, University of Washington. Institute of Forest Resources, Contribution No. 57. Seattle, WA.

508. Sweeney, S. J. 1978. Diet, reproduction and population structure of the bobcat (*Lynx rufus fasciatus*) in western Washington. M. S. Thesis, University of Washington. Seattle, WA.
509. Sydeman, W. J., K. A. Hobson, P. Pyle, and E. B. McLaren. 1997. Trophic relationships among seabirds in central California: combined stable isotope and conventional dietary approach. *Condor* 99:327-336.
510. Tagart, J. V. 1976. The survival from egg deposition to emergence of coho salmon in the Clearwater River, Jefferson County, Washington. M. S. Thesis, University of Washington School of Fisheries, Seattle, WA. 101 pp.
511. Taverner, P. A. 1934. Birds of Canada. Canada Dept. Mines, Natl. Mus. Bull. No. 72 (Biol. Ser. 19). Ottawa. 445 pp.
512. Thedinga, J. F., and K. V. Koski. 1984. A stream ecosystem in an old-growth forest in southeast Alaska. Part VI: The production of coho salmon, *Oncorhynchus kisutch*, smolts and adults from Porcupine Creek. Pages 99-108 in W. R. Meehan, T. R. Merrel, Jr., and T. A. Hanley, editors. Proceedings from a Symposium on Fish and Wildlife Relationships in Old-growth Forests. Am. Inst. Fish. Res. Biol., Juneau, AK.
513. Thom, R. M., and A. B. Borde. 1997. Human intervention in Pacific Northwest coastal ecosystems. Pages 5-37 in G. R. McMurray, and R. J. Bailey, editors. Change in Pacific Northwest Coastal Ecosystems. Science for Solutions, Pacific Northwest Coastal Ecosystems Regional Study (PNCERS). U. S. Department of Commerce. NOAA Coastal Ocean Program Decision Analysis Series No. 11.
514. Thomas, A. C., and P. T. Strub. 1989. Large scale patterns of phytoplankton pigment distribution during the spring transition along the west coast of North America. *J. Geophys. Res.* 94(C12):18-117.
- 514a. Thompson, G. G. 1992. A Bayesian approach to management advice when stock-recruitment parameters are uncertain. *Fishery Bulletin* (U. S.) 90: 561-573.
515. Thomson, R. E. 1981. Oceanography of the British Columbia Coast. *Can. Spec. Publ. Fish. Aquat. Sci.* 56, Dept. Fish. Oceans, Sydney, British Columbia. 291 pp.
516. Thut, R. N. 1970. Feeding habits of the Dipper in southwestern Washington. *Condor* 72:234-235.
- 516a. Tipping, J. M. 1997. Effect of smolt length at release on adult returns of hatchery reared steelhead. *Progressive Fish-Culturist* 59: 310-311.
517. Todd, I., and P. Larkin. 1971. Gillnet selectivity on sockeye (*Oncorhynchus nerka*) and pink salmon (*O. gorbuscha*) of the Skeena River system, British Columbia. *J. Fish. Res. Bd. Can.* 28:821-842.
518. Toweill, D. E. 1974. Winter food habits of River Otters in western Oregon. *Journal of Wildlife Management* 38:107-111.
519. Townsend, C. R., and M. R. Scarsbrook. 1997. The intermediate disturbance hypothesis, refugia, and biodiversity in streams. *Limnol. Oceanogr.* 42(5):938-949.
520. Triska, F. J., J. H. Duff, and R. J. Avanzino. 1990. Influence of exchange flow between channel and hyporheic zone on nitrate production in a small mountain stream. *Canadian Journal of Fisheries and Aquatic Sciences* 47:2099-2111.
521. Trotter, P. C. 1987. Cutthroat - native trout of the West. Colorado Associated University Press. Boulder. 219 pp.
522. Tschaplinski, P. J. 1982. Aspects of the population biology of estuarine-reared and stream-reared juvenile coho salmon in Carnation Creek: A summary of current research. Pages 289-307 in G. F. Hartman, editor. Proceedings of the Carnation Creek Workshop: A Ten-year Review. Malaspina College, Nanaimo, British Columbia.

523. Tschaplinski, P. J. 1987. The use of estuaries as rearing habitats by juvenile coho salmon. Pages 123-141 in T. W. Chamberlin, editor. Proceedings of the Workshop: Applying 15 Years of Carnation Creek Results. Pacific Biol. Station. Nanaimo, British Columbia.
524. Tsuchida, A. 1994. Resource physics and the limitations of nuclear fusion power generation. Selected Papers On Entropy Studies, Vol. 1.
525. Tsuchida, A. 1996. The importance of the "Cycle of Matter": A discussion of nutrient cycling in basins, land, and human society from the viewpoint of entropy, pages 150-170. In Toyohashi, Proceedings of Clean Sea '96. International Workshop and Symposium On Environmental Restoration for Enclosed Seas.
526. United States Army Corps of Engineers (USACOE), U. S. Environmental Protection Agency, U. S. Fish and Wildlife Service, and National Marine Fisheries Service. 1993. Commencement Bay Cumulative Impact Study. U. S. Army Corps of Engineers, Seattle District. Seattle, WA.
527. Van Daele, L. J., and H. A. Van Daele. 1982. Factors affecting the productivity of ospreys nesting in west-central Idaho. Condor 84:292-299.
528. Vander Heyden, M. 1997. Female black bear habitat selection and home range ecology in the central Cascades of Oregon. M.S. Thesis, Oregon State University, Corvallis, OR. 135 pp.
529. Vannote, R. L., G. W. Minshall, K. W. Cummins, J. R. Sedell, and C. E. Cushing. 1980. The river continuum concept. Canadian Journal of Fisheries and Aquatic Sciences 37:130-137.
530. Vermeer, K., and K. Devito. 1986. Size, caloric content, and association of prey fishes in meals of nestling rhinoceros auklets. Murrelet 67:1-9.
531. Vermeer, K., and S. J. Westrheim. 1984. Fish changes in diets of nestling rhinoceros auklets and their implications. Pages 96-105 in D. N. Nettleship, G. A. Sanger, and P. F. Springer, editors. Marine Birds: Their Feeding Ecology and Commercial Fisheries Relationships. Minister of Supply and Services, Canada.
532. Vermeer, K. 1979. Nesting requirements, food and breeding distribution of Rhinoceros Auklets, *Cerorhinca monocerata*, and Tufted puffins, *Lunda cirrhata*. Ardea 67:101-110.
533. Vermeer, K. 1982. Comparison of the diet of the glaucous-winged gull on the east and west coasts of Vancouver Island. Murrelet 63:80-85.
534. Vermeer, K., and K. H. Morgan. 1992. Marine bird populations and habitat use in a fjord on the west coast of Vancouver Island. Canadian Wildlife Service Occasional Paper No. 75:86-96.
535. Verts, B. J., and L. N. Carraway. 1998. Land Mammals of Oregon. University of California Press, Berkeley, CA . 668 pp.
536. Wallace, J. B., S. L. Eggert, J. L. Meyer, and J. R. Webster. 1997. Multiple trophic levels of a forest continuum concept. Canadian Journal of Fisheries and Aquatic Sciences 37:130-104.
537. Watson, J. W., M. G. Garrett and R. G. Anthony. 1991. Foraging ecology of bald eagles in the Columbia River estuary. Journal of Wildlife Management 55:492-499.
- 537a. Ward, B. R., and P. A. Slaney. 1988. Life history and smolt-to-adult survival of Keogh River steelhead trout and the relationship to smolt size. Canadian Journal of Fisheries and Aquatic Sciences 1110-1122.
538. Warren, C. E. 1971. Biology and water pollution control. The Department of Fisheries and Wildlife, Oregon State University. Corvallis, OR.

- 538a. Washington Department of Fisheries. 1992. Draft. Goals and objectives for stocks and fisheries. Olympia, WA.
- 538b. Washington Department of Fisheries. 1983. Strategic plan for management of Washington's salmon resources. Olympia, WA.
539. Washington Department of Fish and Wildlife and Western Washington Treaty Indian Tribes. 1994. 1992 Washington State Salmon And Steelhead Stock Inventory. Appendix One, Puget Sound Stocks. 371 pp.
540. Washington Department of Fisheries (WDF) and Oregon Department of Fish and Wildlife (ODFW). 1992. Status report. Columbia River fish runs & fisheries, 1938-91. Olympia, WA.
541. Washington Department of Fish and Wildlife (WDFW). 1998. 1998 Washington salmonid stock inventory, coastal cutthroat trout appendix. DRAFT. Washington Department of Fish and Wildlife. Olympia, WA.
- 541a. Washington Department of Fish and Wildlife (WDFW). 1998. Forage fish management plan - A plan for managing the forage fisheries in Washington. Adopted by the Washington Fish and Wildlife Commission on January 24, 1998.
542. Washington Department of Fisheries, Washington Department of Wildlife, and Western Washington Treaty Indian Tribes. 1993. 1992 Washington State Salmon And Steelhead Stock Inventory. Washington Department of Fish and Wildlife, Olympia. 212 pp.
543. Washington Department of Ecology. 1994. Inventory of dams in the state of Washington. Washington Department of Ecology, Water Resources Program, Dam Safety Division. Revised Edition, January 1994, Publication #94-16. 150 pp.
544. Washington Department of Fish and Wildlife (WDFW). 1997. Wild Salmonid Policy. Washington Department of Fish and Wildlife. Olympia, WA.
545. Washington Department of Natural Resources (WDNR). 1997. Final Habitat Conservation Plan. Washington Department of Natural Resources, Jennifer Belcher, Commissioner of Public Lands. Olympia, WA.
546. Washington Department of Fish and Wildlife and Oregon Department of Fish and Wildlife. 1994. Status Report. Columbia fish runs and fisheries 1938-93. Olympia, WA. 275 pp.
547. Waters, T. F. 1969. Invertebrate drift - Ecology and significance to stream fishes. Pages 121-134 in T. G. Northcote, editor. H. R. MacMillan Lectures In Fisheries. Held Feb. 22-24, 1968. University of British Columbia. Vancouver.
548. Wehle, D. H. S. 1983. The food, feeding, and development of young tufted and horned puffins in Alaska. Condor 85:427-442.
549. Weitkamp, L. A., T. C. Wainwright, G. J. Bryant, G. B. Milner, D. J. Teel, R. G. Kope, and R. S. Waples. 1995. Status review of coho salmon from Washington, Oregon, and California. U.S. Dept. of Commerce. NMFS-NWFSC-24: 258 pp.
550. Wendler, H. O., and G. Deschamps. 1955. Logging dams on coastal Washington streams. Fish. Res. Pap. 1: 27-38, Washington Department of Fisheries, Olympia, WA.
551. White, H. C. 1936. The food of kingfishers and mergansers on the Margaree River, Nova Scotia. J. Biol. Bd. Can. 2:299-309.
552. White H. C. 1939. Bird control to increase the Margaree River salmon. Bull. Fish. Res. Bd. Can. 58:30 pp.
553. White, H. C. 1957. Food and natural history of mergansers on salmon waters in the Maritime provinces of Canada. Fish. Res. Bd. Can., Bulletin No. 116. 63 pp.

554. White, R. 1992. Land use, environment, and social change: The shaping of Island County, Washington. University of Washington Press. Seattle, WA.
555. Whitman, J. S. 1981. Ecology of the mink (*Mustela vison*) in west-central Idaho. M.S. Thesis, University of Idaho. 101 pp.
556. Williams, W. R., M. Laramie, and J. J. Ames. 1975. Catalog of Washington Streams and Salmon Utilization - Volume 1 Puget Sound. Washington Department of Fisheries. Olympia.
557. Williams, D. D., and H. B. N. Hynes. 1974. The occurrence of benthos deep in the substratum of a stream. *Freshwat. Biol.* 4:233-256.
558. Williams, D. D., and B. W. Feltmate. 1992. Aquatic insects. CAB International. United Kingdom. 358 pp.
559. Willson, M. F., and K. C. Halupka. 1995. Anadromous fish as keystone species in vertebrate communities. *Conser. Biol.* 9:489-497.
560. Winter, B. D. 1992. Determinate migratory behavior of steelhead (*Oncorhynchus mykiss*) parr. Ph.D. Dissertation. University of Washington, School of Fisheries. Seattle, WA. 255 pp.
561. Willson, M. F., S. M. Gende, and B. H. Marston. 1998. Fishes and the Forest: Expanding perspectives on fish-wildlife interactions. *BioScience* 48:455-462.
562. Willson, M. F., and K.C. Halupka 1995. Anadromous fish as keystone species in vertebrate communities. *Conservation Biology* 9:489-497.
563. Wilson, U. W., and D. A. Manuwal. 1986. Breeding biology of the Rhinoceros Auklet in Washington. *Condor* 88:143-155.
564. Wipfli, M. S., J. Hudson, and J. Caouette. 1998. Influence of salmon carcasses on stream productivity: response of biofilm and benthic macroinvertebrates in southeastern Alaska, U.S.A. *Canadian Journal of Fisheries and Aquatic Sciences* 55:1503-1511.
565. Wissmar, R. C., and C. A. Simenstad. 1988. Energetic constraints of juvenile chum salmon (*Oncorhynchus keta*) migrating in estuaries. *Canadian Journal of Fisheries and Aquatic Sciences* 45:1555-1560.
566. Wissmar, R. C., and C. A. Simenstad. 1998. Variability of estuarine and riverine ecosystem productivity for supporting Pacific salmon. Pages 253-301 in G. R. McMurray, and R. J. Bailey, editors. *Change in Pacific Northwest Coastal Ecosystems*. NOAA Coastal Ocean Prog., Decision Analysis Series No. 11, NOAA Coastal Ocean Office. Silver Spring, MD. 342 pp.
567. Wood, C. C. 1986. Dispersion of Common Merganser (*Mergus merganser*) breeding pairs in relation to availability of juvenile Pacific salmon in Vancouver Island streams. *Canadian Journal of Zoology* 64:756-765.
568. Wood, C. C. 1985. Aggregative response of Common Mergansers (*Mergus merganser*): predicting flock size and abundance on Vancouver Island salmon streams. *Canadian Journal of Fisheries and Aquatic Sciences* 42:1259-1271.
569. Wood, C. C., and C. M. Hand. 1985. Food searching behaviour of the common merganser (*Mergus merganser*) 1: functional responses to prey and predator density. *Canadian Journal of Zoology* 63:1260-1270.
570. Wood, C.C. 1987a. Predation by the common merganser (*Mergus merganser*) on Eastern Vancouver Island. I: Predation during the seaward migration. *Canadian Journal of Fisheries and Aquatic Sciences* 44:941-949.
571. Wood, C.C. 1987b. Predation by the common merganser (*Mergus merganser*) on Eastern Vancouver Island. II: Predation of stream-resident juvenile salmon by merganser broods. *Canadian Journal of Fisheries and Aquatic Sciences* 44:950-959.
572. Wurtsbaugh, W. A., and G. E. Davis. 1977. Effects of temperature and ration level on the growth and food conversion efficiency of *Salmo gairdneri* Richardson. *J. Fish Biol.* 11:87-98.

573. Wydoski, R.S., and R.R. Whitney. 1979. Inland fishes of Washington. University of Washington Press. Seattle, WA.
574. Yoakum, J. 1964. Observations on bobcat-water relationships. *Journal of Mammalogy* 45:477-479.
575. Young, S. P. 1944. The Wolves of North America. Their history, life habits, economic status, and control. The American Wildlife Institute, Washington, D.C. 635 pp.
576. Zarnowitz, J. E., and K. J. Raedeke. 1984. Winter predation on coho fingerlings by birds and mammals in relation to pond characteristics. College of Forest Resources, University of Washington. Service Contract No. 1480. Seattle, WA. 34 pp.

Appendix I. The 9 wildlife species identified as having (or historically had) a *strong, consistent* relationship with salmon in Oregon and Washington. An "x" identifies the life stage(s) of salmon applicable to the species.

	<i>Incubtion</i>	<i>Freshwater Rearing</i>	<i>Saltwater</i>	<i>Spawning</i>	<i>Carcass</i>	<i>Comments</i>
Common Merganser	x	x	x			
Harlequin Duck	x		x			Strong relationship w/ drift eggs and alevin; indirect relationship w/ carcass-derived insects
Osprey		x	x	x		
Bald Eagle			x	x	x	Strong relationship w/ salmon; also indirect relationship -- feeds on gulls, terns, and waterfowl that eat salmon; occasionally have been seen catching and consuming smolts.
Caspian Tern		x	x			
Black Bear				x	x	
Grizzly Bear				x	x	
Northern River Otter		x		x	x	
Killer Whale			x			

Appendix II. The 58 wildlife species identified as having (or historically had) a *recurrent* relationship with salmon in Oregon and Washington. An "x" identifies the life stage(s) of salmon applicable to the species.

	<i>Incubtion</i>	<i>Freshwater Rearing</i>	<i>Saltwater</i>	<i>Spawning</i>	<i>Carcass</i>	<i>Comments</i>
Cope's Giant Salamander	x	x			?	Also potential occasional relationship w/ carcasses
Pacific Giant Salamander	x	x				
Pacific Coast Aquatic Garter Snake	x	x				
Red-throated Loon		x	x			
Pacific Loon			x			Also potential relationship with spawning salmon or carcasses
Common Loon		x	x			Also potential relationship with spawning salmon or carcasses
Pied-billed Grebe		x				
Western Grebe		x	x			
Clark's Grebe			x			
American White Pelican		x				
Brandt's Cormorant		x	x			
Double-crested Cormorant		x	x			
Pelagic Cormorant		x	x			
Great Blue Heron		x	x			
Black-crowned Night-heron		x	x			
Turkey Vulture					x	
California Condor					x	A historic relationship based on 1800's literature and archaeological evidence
Common Goldeneye	x	x			x	
Barrow's Goldeneye	x	x			x	
Common Merganser					x	
Red-breasted Merganser	x	x	x			
Golden Eagle				x	x	

Bonaparte's Gull	x		x		x	
Heermann's Gull			x			
Ring-billed Gull		x	x		x	
California Gull			x		x	
Herring Gull		x	x		x	
Thayer's Gull			x			
Western Gull			x		x	
Glaucous-winged Gull	x		x	x	x	
Glaucous Gull			x		x	
Common Tern		x	x			
Arctic Tern		x	x			
Forster's Tern		x	x			
Elegant Tern			x			
Common Murre			x			
Marbled Murrelet		x	x			
Rhinoceros Auklet			x			
Tufted Puffin			x			
Belted Kingfisher		x	x	x		
American Dipper	x	x			x	Direct relationship w/ drift eggs and fry; indirect relationship with carcass-derived insects
Steller's Jay					x	
Black-billed Magpie		x			x	
American Crow		x			x	
Northwestern Crow		x	x		x	
Common Raven		x		x	x	
Virginia Opossum					x	
Water Shrew	x	x			x	May eat drift eggs, fry; indirect relationship with carcass-derived insects
Coyote					x	
Gray Wolf				x	x	
Raccoon		x			x	
Mink		x		x	x	

Bobcat				x	x	
Northern Fur Seal			x			
Northern (Steller) Sea Lion			x	x	x	
California Sea Lion			x	x		
Harbor Seal			x	x	x	
Pacific White-sided Dolphin			x			

Appendix III. The 25 wildlife species identified as having an *indirect* relationship with salmon in Oregon and Washington. An "x" identifies the life stage(s) of salmon applicable to the species.

	<i>Incubation</i>	<i>Freshwater Rearing</i>	<i>Saltwater</i>	<i>Spawning</i>	<i>Carcass</i>	<i>Comments</i>
Harlequin Duck					x	Indirect relationship w/ carcass-derived insects; direct relationship w/ eggs
Bald Eagle	x	x	x		x	Indirect relationship -- feeds on gulls, terns, and waterfowl that eat salmon; also strong, direct relationship w/ salmon
Gyrfalcon		x	x		x	Feeds on waterfowl and gulls that eat fish
Peregrine Falcon		x	x		x	Feeds on waterfowl and gulls that eat fish
Killdeer					x	Carcass nutrients support insect supply
Spotted Sandpiper					x	Carcass nutrients support insect supply
Snowy Owl		x				Feeds on waterfowl that eat fish
Willow Flycatcher					x	Carcass nutrients likely to support insect supply
Tree Swallow					x	Carcass nutrients likely to support insect supply
Violet-green Swallow					x	Carcass nutrients likely to support insect supply
Northern Rough-winged Swallow					x	Carcass nutrients likely to support insect supply
Bank Swallow					x	Carcass nutrients likely to support insect supply
Cliff Swallow					x	Carcass nutrients likely to support insect supply
Barn Swallow					x	Carcass nutrients likely to support insect supply
American Dipper					x	Indirect relationship with carcass-derived insects; direct relationship w/ drift eggs and fry
Masked Shrew					x	Likely to eat both carcass meat and insects associated w/ carcasses
Vagrant Shrew					x	Likely to eat both carcass meat and insects associated w/ carcasses
Montane Shrew					x	Likely to eat both carcass meat and insects associated w/ carcasses
Fog Shrew					x	Likely to eat both carcass meat and insects associated w/ carcasses

Pacific Shrew					x	Likely to eat both carcass meat and insects associated w/ carcasses
Water Shrew					x	May eat drift eggs, fry; indirect relationship with carcass-derived insects
Pacific Water Shrew					x	Likely to eat both carcass meat and insects associated w/ carcasses
Trowbridge's Shrew					x	Likely to eat both carcass meat and insects associated w/ carcasses
Harbor Porpoise			x			Feeds on species that feed on smolts
Dall's Porpoise			x			Feeds on species that feed on smolts

Appendix IV. The 65 wildlife species identified as having (or historically had) a *rare* relationship with salmon in Oregon and Washington. An "x" identifies the life stage(s) of salmon applicable to the species.

	<i>Incubation</i>	<i>Freshwater Rearing</i>	<i>Saltwater</i>	<i>Spawning</i>	<i>Carcass</i>	<i>Comments</i>
Snapping Turtle		x				
Western Pond Turtle		x				
Western Terrestrial Garter Snake		x				
Common Garter Snake		x				
Pacific Loon					x	
Common Loon					x	
Yellow-billed Loon			x			
Horned Grebe	x		x			
Red-necked Grebe			x		x	
Western Grebe					x	
Sooty Shearwater			x			
Brown Pelican			x			
Great Egret		x	x			
Snowy Egret		x				
Green Heron		x	x			
Trumpeter Swan	x	x			x	
Mallard	x				x	
Green-winged Teal	x					
Canvasback					x	
Greater Scaup	x				x	
Surf Scoter			x		x	
White-winged Scoter			x		x	
Common Goldeneye			x		x	
Barrow's Goldeneye			x			

	<i>Incubation</i>	<i>Freshwater Rearing</i>	<i>Saltwater</i>	<i>Spawning</i>	<i>Carcass</i>	<i>Comments</i>
Hooded Merganser	x	x			x	
Red-tailed Hawk					x	
Greater Yellowlegs	x					
Franklin's Gull		x				
Mew Gull	x					
Black-legged Kittiwake			x		x	
Pigeon Guillemot			x			
Ancient Murrelet			x			
Gray Jay					x	
Winter Wren					x	
American Robin	x					
Varied Thrush	x				x	
Spotted Towhee					x	
Song Sparrow					x	
Masked Shrew					x	Likely to eat both carcass meat and insects associated w/ carcasses
Vagrant Shrew					x	Likely to eat both carcass meat and insects associated w/ carcasses
Montane Shrew					x	Likely to eat both carcass meat and insects associated w/ carcasses
Fog Shrew					x	Likely to eat both carcass meat and insects associated w/ carcasses
Pacific Shrew					x	Likely to eat both carcass meat and insects associated w/ carcasses
Pacific Water Shrew					x	
Trowbridge's Shrew					x	

	<i>Incubation</i>	<i>Freshwater Rearing</i>	<i>Saltwater</i>	<i>Spawning</i>	<i>Carcass</i>	<i>Comments</i>
Douglas' Squirrel					x	
Northern Flying Squirrel					x	
Deer Mouse					x	
Red Fox					x	
Gray Fox					x	
Ringtail					x	
American Marten					x	
Fisher					x	
Long-tailed Weasel					x	
Wolverine					x	
Striped Skunk					x	
Mountain Lion				x	x	
White-tailed Deer					x	
Black-tailed Deer					x	
Minke Whale			x			
Sperm Whale			x			
Humpback Whale			x			
Northern Right-whale Dolphin			x			
Dall's Porpoise			x			
Harbor Porpoise			x			

Appendix V. The 60 wildlife species identified as having an *unknown* relationship with salmon in Oregon and Washington.

Baird's Shrew	Hoary Bat	Rough-skinned Newt
Big Brown Bat	Keen's Myotis	Short-billed Dowitcher
Black Phoebe	Least Flycatcher	Shrew-mole
Brazilian Free-tailed Bat	Little Brown Myotis	Silver-haired Bat
Brewer's Blackbird	Long-eared Myotis	Southern Torrent Salamander
Bullfrog	Long-legged Myotis	Spotted Bat
California Myotis	Long-toed Salamander	Tailed Frog
Cascade Torrent Salamander	Merriam's Shrew	Townsend's Big-eared Bat
Columbia Torrent Salamander	Northern Leopard Frog	Townsend's Chipmunk
Columbian Mouse	Northern Waterthrush	Townsend's Vole
Cordilleran Flycatcher	Northwestern Salamander	Van Dyke's Salamander
Dunlin	Olympic Torrent Salamander	Warbling Vireo
Dunn's Salamander	Oregon Spotted Frog	Water Vole
Dusky Flycatcher	Pacific-slope Flycatcher	Western Pipistrelle
Ermine	Painted Turtle	Western Sandpiper
European Starling	Pallid Bat	Western Small-footed Myotis
Foothill Yellow-legged Frog	Purple Martin	Western Spotted Skunk
Fringed Myotis	Pygmy Shrew	Western Toad
Gray Catbird	Red-eared Slider Turtle	Woodhouse's Toad
Hammond's Flycatcher	Red-legged Frog	Yuma Myotis

Appendix VI. The 407 wildlife species identified as having (or historically had) *no* relationship with salmon in Oregon and Washington.

Acorn Woodpecker	Boreal Chickadee	Common Nighthawk
Aleutian Canada Goose	Boreal Owl	Common Poorwill
Allen's Chipmunk	Botta's (Pistol River) Pocket Gopher	Common Porcupine
Allen's Hummingbird	Brant	Common Redpoll
American Avocet	Brewer's Sparrow	Common Snipe
American Badger	Broad-footed Mole	Common Yellowthroat
American Beaver	Broad-tailed Hummingbird	Cooper's Hawk
American Bittern	Brown Creeper	Creeping Vole
American Black Duck	Brown-headed Cowbird	Dark Kangaroo Mouse
American Coot	Brush Rabbit	Dark-eyed Junco
American Golden-Plover	Buff-breasted Sandpiper	Del Norte Salamander
American Goldfinch	Bufflehead	Desert Horned Lizard
American Kestrel	Buller's Shearwater	Desert Woodrat
American Pika	Bullock's Oriole	Downy Woodpecker
American Pipit	Burrowing Owl	Dusky Canada Goose
American Redstart	Bushtit	Dusky-footed Woodrat
American Tree Sparrow	Bushy-tailed Woodrat	Eared Grebe
American Wigeon	Cackling Canada Goose	Eastern Cottontail
Anna's Hummingbird	California Bighorn Sheep	Eastern Fox Squirrel
Ash-throated Flycatcher	California Ground Squirrel	Eastern Gray Squirrel
Baird's Sandpiper	California Kangaroo Rat	Eastern Kingbird
Band-tailed Pigeon	California Mountain Kingsnake	Ensatina
Barn Owl	California Quail	Eurasian Wigeon
Barred Owl	California Slender Salamander	European Rabbit
Belding's Ground Squirrel	California Towhee	Evening Grosbeak
Bewick's Wren	California Vole	Feral Horse
Bison	Calliope Hummingbird	Feral Pig
Black Oystercatcher	Camas Pocket Gopher	Ferruginous Hawk
Black Rat	Canyon Mouse	Flammulated Owl
Black Rosy-finch	Canyon Wren	Flesh-footed Shearwater
Black Salamander	Cascade Golden-mantled Ground Squirrel	Fork-tailed Storm-petrel
Black Scoter	Cascades Frog	Fox Sparrow
Black Swift	Cassin's Auklet	Gadwall
Black Tern	Cassin's Finch	Giant Canada Goose
Black Turnstone	Cassin's Vireo	Golden-crowned Kinglet
Black-backed Woodpecker	Cattle Egret	Golden-crowned Sparrow
Black-bellied Plover	Cedar Waxwing	Golden-mantled Ground Squirrel
Black-capped Chickadee	Chestnut-backed Chickadee	Gopher Snake
Black-chinned Hummingbird	Chipping Sparrow	Grasshopper Sparrow
Black-footed Albatross	Chisel-toothed Kangaroo Rat	Gray Flycatcher
Black-headed Grosbeak	Chukar	Gray Partridge
Black-necked Stilt	Cinnamon Teal	Gray Whale
Black-tailed Jackrabbit	Clark's Nutcracker	Gray-crowned Rosy-Finch
Black-throated Gray Warbler	Clay-colored Sparrow	Gray-tailed Vole
Black-throated Sparrow	Clouded Salamander	Great Basin Pocket Mouse
Blue Grouse	Coast Mole	Great Basin Spadefoot
Blue Whale	Columbia Spotted Frog	Great Gray Owl
Blue-gray Gnatcatcher	Columbian Ground Squirrel	Great Horned Owl
Blue-winged Teal	Columbian White-tailed Deer	Greater White-fronted Goose
Bobolink	Common Kingsnake	Green Frog
Bohemian Waxwing		Green-tailed Towhee
		Hairy Woodpecker

Harris's Sparrow	Mute Swan	Red Knot
Heather Vole	Nashville Warbler	Red Phalarope
Hermit Thrush	Night Snake	Red Squirrel
Hermit Warbler	North Pacific Bottle-nosed Whale	Red Tree Vole
Hoary Marmot	Northern Alligator Lizard	Red-breasted Nuthatch
Horned Lark	Northern Bobwhite	Red-breasted Sapsucker
House Finch	Northern Bog Lemming	Red-eyed Vireo
House Mouse	Northern Elephant Seal	Redhead
House Sparrow	Northern Flicker	Red-naped Sapsucker
House Wren	Northern Fulmar	Red-necked Phalarope
Hutton's Vireo	Northern Goshawk	Red-shouldered Hawk
Juniper Titmouse	Northern Grasshopper Mouse	Red-tailed Chipmunk
Kit Fox	Northern Harrier	Red-winged Blackbird
Lapland Longspur	Northern Mockingbird	Ringneck Snake
Larch Mountain Salamander	Northern Pintail	Ring-necked Duck
Lark Sparrow	Northern Pocket Gopher	Ring-necked Pheasant
Laysan Albatross	Northern Pygmy-owl	Risso's Dolphin
Lazuli Bunting	Northern Saw-whet Owl	Rock Dove
Leach's Storm-petrel	Northern Shoveler	Rock Sandpiper
Least Bittern	Northern Shrike	Rock Wren
Least Chipmunk	Northwestern Garter Snake	Rocky Mountain Bighorn Sheep
Least Sandpiper	Norway Rat	Rocky Mountain Elk
Leatherback Turtle	Nutria	Roosevelt Elk
Lesser Canada Goose	Nuttall's (Mountain) Cottontail	Ross's Goose
Lesser Goldfinch	Oak Titmouse	Rough-legged Hawk
Lesser Scaup	Oldsquaw	Rubber Boa
Lesser Yellowlegs	Olive-sided Flycatcher	Ruby-crowned Kinglet
Lewis's Woodpecker	Olympic Marmot	Ruddy Duck
Lincoln's Sparrow	Orange-crowned Warbler	Ruddy Turnstone
Little Pocket Mouse	Ord's Kangaroo Rat	Ruff
Loggerhead Shrike	Oregon Slender Salamander	Ruffed Grouse
Long-billed Curlew	Pacific Golden-Plover	Rufous Hummingbird
Long-billed Dowitcher	Pacific Jumping Mouse	Sabine's Gull
Long-eared Owl	Pacific Treefrog	Sage Grouse
Long-nosed Leopard Lizard	Palm Warbler	Sage Sparrow
Long-tailed Jaeger	Parasitic Jaeger	Sage Thrasher
Long-tailed Vole	Pectoral Sandpiper	Sagebrush Lizard
Lynx	Pileated Woodpecker	Sagebrush Vole
Macgillivray's Warbler	Pine Grosbeak	Sanderling
Marbled Godwit	Pine Siskin	Sandhill Crane
Marsh Wren	Pink-footed Shearwater	Savannah Sparrow
Meadow Vole	Pinon Mouse	Say's Phoebe
Merlin	Pinyon Jay	Scaled Quail
Merriam's Ground Squirrel	Piute Ground Squirrel	Sea Otter
Mojave Black-collared Lizard	Plateau Striped Whiptail	Semipalmated Plover
Montane Vole	Plumbeous Vireo	Semipalmated Sandpiper
Moose	Pomarine Jaeger	Sharp-shinned Hawk
Mountain Beaver	Prairie Falcon	Sharptail Snake
Mountain Bluebird	Preble's Shrew	Sharp-tailed Grouse
Mountain Caribou	Pronghorn Antelope	Sharp-tailed Sandpiper
Mountain Chickadee	Purple Finch	Short-eared Owl
Mountain Goat	Pygmy Nuthatch	Short-finned Pilot Whale
Mountain Quail	Pygmy Rabbit	Short-horned Lizard
Mourning Dove	Racer	Short-tailed Albatross
Mule Deer	ed Crossbill	Short-tailed Shearwater
Muskrat		Side-blotched Lizard

Siskiyou Chipmunk
Siskiyou Mtns. Salamander
Sky Lark
Snow Bunting
Snow Goose
Snowshoe Hare
Snowy Plover
Solitary Sandpiper
Sora
South Polar Skua
Southern Alligator Lizard
Southern Red-backed Vole
Spotted Owl
Spruce Grouse
Stilt Sandpiper
Striped Whipsnake
Surfbird
Swainson's Hawk
Swainson's Thrush
Swamp Sparrow
Taverner's Canada Goose
Three-toed Woodpecker
Tiger Salamander
Townsend's Ground Squirrel
Townsend's Mole
Townsend's Pocket Gopher
Townsend's Solitaire
Townsend's Warbler
Tricolored Blackbird

Tundra Swan
Upland Sandpiper
Vancouver Canada Goose
Vaux's Swift
Veery
Vesper Sparrow
Virginia Rail
Wandering Tattler
Washington Ground Squirrel
Western Bluebird
Western Canada Goose
Western Fence Lizard
Western Gray Squirrel
Western Ground Snake
Western Harvest Mouse
Western Jumping Mouse
Western Kingbird
Western Meadowlark
Western Pocket Gopher
Western Rattlesnake
Western Red-backed Salamander
Western Red-backed Vole
Western Screech-owl
Western Scrub-Jay
Western Skink
Western Tanager
Western Whiptail
Western Wood-pewee
Whimbrel

White-breasted Nuthatch
White-crowned Sparrow
White-faced Ibis
White-footed Vole
White-headed Woodpecker
White-tailed Antelope Squirrel
White-tailed Jackrabbit
White-tailed Kite
White-tailed Ptarmigan
White-throated Sparrow
White-throated Swift
White-winged Crossbill
Wild Burro
Wild Turkey
Willet
Williamson's Sapsucker
Wilson's Phalarope
Wilson's Warbler
Wood Duck
Wrentit
Wyoming Ground Squirrel
Yellow Rail
Yellow Warbler
Yellow-bellied Marmot
Yellow-billed Cuckoo
Yellow-breasted Chat
Yellow-headed Blackbird
Yellow-pine Chipmunk
Yellow-rumped Warbler

Appendix VII. List of published and unpublished observations of wildlife predation and scavenging on salmon.

Salmon life stage	Species	Relationship to salmon	References	Location of study/report
INCUBATION – EGGS AND ALEVIN	Cope's Giant Salamander	Recurrent	Johnson et al. 2000	
	Pacific Giant Salamander	Recurrent	Graf 1949	California
	Pacific Coast Aquatic Garter Snake	Recurrent	Brown et al. 1995	Oregon
	Common Merganser	Strong	Munro and Clemens 1932, 1937, 1939	British Columbia
	Hooded Merganser	Rare	Reimchen 1994	British Columbia
	Red-breasted Merganser	Recurrent	Munro and Clemens 1939	Canada
	Barrow's Goldeneye	Recurrent	Willson and Halupka 1995	Alaska
			Munro 1938, 1939	British Columbia
	Common Goldeneye	Recurrent	Willson and Halupka 1995	Alaska
			Munro 1938, 1939	British Columbia
	American Dipper	Recurrent	Ehinger 1930	Washington
			Munro 1923, Obermayer et al. 1999, Piorkowski 1995, Willson and Halupka 1995	Alaska
			Reimchen 1994, Burcham 1904	British Columbia
	Glaucous-winged Gull	Recurrent	Baird 1990	Washington
			Reimchen 1994	British Columbia
			Mossman 1958, Moyle 1966	Alaska
	Bonaparte's Gull	Recurrent	Moyle 1966	Alaska
	Harlequin Duck	Strong	Dzinbal and Jarvis 1982, Obermayer et al. 1999	Alaska
	Horned Grebe	Rare	Palmer 1962	Washington
			Munro 1941	British Columbia
Trumpeter Swan	Rare	Farley 1980	Alaska	
Mallard	Rare	Willson and Halupka 1995	Alaska	
Green-winged Teal	Rare	Ned Pittman, pers. comm.	Washington	
Greater Scaup	Rare	Munro 1941	British Columbia	
Greater Yellowlegs	Rare	Elphick and Tibbitts 1998	Alaska	
Mew Gull	Rare	Moyle 1966	Alaska	
American Robin	Rare	Willson and Halupka 1995	Alaska	

FRESHWATER REARING – FRY AND PUPP	Varied Thrush	Rare	Ned Pittman pers. comm.	Washington
	Water Shrew	Recurrent	Banfield 1974	Canada
	Cope's Giant Salamander	Recurrent	Antonelli et al. 1972	Washington
	Pacific Giant Salamander	Recurrent	Antonelli et al. 1972	Washington
			Parker 1993, Parker 1994	California
	Snapping Turtle	Rare	Johnson and Hasler 1954, Lagler 1943	Michigan
	Western Pond Turtle	Rare	Johnson et al. 2000	
	Pacific Coast Aquatic Garter Snake	Recurrent	Brown et al. 1995, Fitch 1941, 1984, Drummond 1983 Hansen 1980	Oregon, California
	Western Terrestrial Garter Snake	Rare	Anderson 1977, Tanner 1949	Montana, Utah
	Common Garter Snake	Rare	Lagler and Salyer 1945	Michigan
	Red-throated Loon	Recurrent	Eriksson et al. 1990	Sweden
			Palmer 1962	Labrador
	Common Loon	Recurrent	Palmer 1962, Johnson and Hasler 1954, Alexander 1977	Michigan
			Fraser 1972, 1974, Matkowski 1989, Matkowski 1984, Barr 1973, Smith 1968, Munro 1945	Canada, British Columbia
			Willson and Halupka 1995	Alaska
	Pied-billed Grebe	Recurrent	Zarnowitz and Raedeke 1984	Washington
	Western Grebe	Recurrent	Modde and Wasowicz 1996	Utah
	American White Pelican	Recurrent	Myers and Peterka 1976, Lingle 1977	North Dakota
			Palmer 1962	Wyoming
	Brandt's Cormorant	Recurrent	Johnson et al. 2000	
Double-crested	Recurrent	Modde and Wasowicz 1996	Utah	
		Mayers and Peterka 1976	North Dakota	
Pelagic Cormorant	Recurrent	Johnson et al. 2000		
Great Blue Heron	Recurrent	Alexander 1977, 1979, Bent 1926, Johnson and Hasler 1954	Michigan	
		Fraser 1972, Matkowski 1989, Smith 1968	Canada	
		Dolloff 1993, Willson and Halupka 1995	Alaska	
		Zarnowitz and Raedeke 1984	Washington	
		Henney and Bethers 1971	Oregon	

**FRESHWATER
REARING –
FRY AND
PARR**

Great Egret	Rare	Johnson et al. 2000	
Snowy Egret	Recurrent	Johnson et al. 2000	
Green Heron	Rare	Jurek 1974	California
Black-crowned Night	Recurrent	Spanier 1980	Israel
		Myers and Peterka 1976	North Dakota
Trumpeter Swan	Rare	Farley 1980	Alaska
		Hampton 1981	Montana
Common Goldeneye	Recurrent	Beach 1937	Michigan
		White 1939	Nova Scotia
Barrow's Goldeneye	Recurrent	Munro and Clemens 1938, 1939	British Columbia
Hooded Merganser	Rare	Zarnowitz and Raedeke 1984	Washington
Common Merganser	Strong	Alexander 1977, 1979, Beach 1937, Johnson and Halser 1954, Salyer and Lagler 1940, Shetter 1970	Michigan
		Miegs and Rieck 1967, Senn 1958	Washington
		Fraser 1972, Huntsman 1941, Munro and Clemens 1932, 1937, 1939, Smith 1968	Canada
		Willson and Halupka 1995	Alaska
		White 1936, 1957	Nova Scotia
Red-breasted merganser	Recurrent	Munro and Clemens 1939	Canada
		Marquiss and Duncan 1993	Scotland
Osprey	Strong	Swenson 1978	Wyoming
		Steeger et al. 1992	British Columbia
		MacCarter 1972	Montana
		Johnson and Hasler 1954	Michigan
		Van Daele and Van Daele 1982	Idaho
		French and Koplín 1977	California
		Hughes 1983	Alaska
		Lind 1976	Oregon
Franklin's Gull	Rare	Myers and Peterka 1976	North Dakota
Ring-billed Gull	Recurrent	Nui Tateyama, pers. comm.	Washington
Caspian Tern	Strong	Johnson et al. 2000	

**FRESHWATER
REARING –
FRY AND
PARR**

Herring Gull	Recurrent	Mendall 1939	Maine
Forster's Tern	Recurrent	Ayles et al. 1976	Canada
Common Tern	Recurrent	Ayles et al. 1976	Canada
Arctic Tern	Recurrent	Mossman 1959, Willson and Halupka 1995	Alaska
Marbled Murrelet	Recurrent	Brooks 1928	British Columbia
		Carter and Sealy 1984	Alaska
Belted Kingfisher	Recurrent	Alexander 1977, 1979	Michigan
		Willson and Halupka 1995	Alaska
		Gould 1934, Eipper 1956	New York
		White 1936	Nova Scotia
Black-billed Magpie	Recurrent	Elson 1962, Huntsman 1941	Canada
		Willson and Halupka 1995	Alaska
American Crow	Recurrent	Willson and Halupka 1995	Alaska
Northwestern Crow	Recurrent	Willson and Halupka 1995	Alaska
Common Raven	Recurrent	Johnson et al. 2000	
American Dipper	Recurrent	Dolloff 1993	Alaska
		Loegering 1997	Oregon
		Thut 1970	Washington
		Kingery 1996, Cottam and Uhler 1937	British Columbia
Water Shrew	Recurrent	Lampman 1947	Oregon
		Banfield 1974	Canada
		Conaway 1952	Montana
Raccoon	Recurrent	Alexander 1977	Michigan
Mink	Recurrent	Whitman 1981	Idaho
		Dunstone 1993	New York
		Banfield 1974, Burgess and Bider 1980, Fraser 1972	Canada
		Ben-David et al. 1997	Alaska
		Grinnell et al. 1937	California
		Alexander 1977, 1979	Michigan
Akande 1972	Scotland		
Northern River Otter	Strong	Zarnowitz and Raedeke 1984	Washington

FRESHWATER – FRY AND PARR		Strong	Banfield 1974, Stenson et al. 1984	British Columbia
			Dolloff 1993	Alaska
			Alexander 1979	Michigan
SALTWATER – SMOLT, IMMATURE ADULTS AND ADULTS	Pacific Loon	Recurrent	Mace 1983	British Columbia
	Red-throated Loon	Recurrent	Johnson et al. 2000	
	Common loon	Recurrent	Palmer 1962	Alaska
			Mace 1983	British Columbia
	Yellow-billed Loon	Rare	North 1994	Russia
	Horned Grebe	Rare	Mace 1983, Vermeer 1992	British Columbia
	Red-necked Grebe	Rare	Mace 1983	British Columbia
	Western Grebe	Recurrent	Vermeer et al. 1992, Mace 1983	British Columbia
	Clark's Grebe	Recurrent	Johnson et al. 2000	
	Sooty Shearwater	Rare	Emmett 1997, Bayer 1989	Oregon
	Brown Pelican	Rare	Bayer 1986a, Emmett 1997, McNeil et al. 1991	Oregon
	Brandt's Cormorant	Recurrent	Aniley and Sanger 1979	California
			Bayer 1986a, Scott 1973	Oregon
	Double-crested Cormorant	Recurrent	Bayer 1986a, Bayer 1989, Erickson 1988, Hoffman and Hall 1988, Roby et al. 1998	Oregon
			Ainley and Anderson 1981, Mace 1993, Robertson 1974	British Columbia
	Pelagic Cormorant	Recurrent	Bayer 1986a, Scott 1973	Oregon
			Jewett et al. 1953	Washington
			Mace 1983	British Columbia
	Great Blue Heron	Recurrent	Forbes and Simpson 1982, Mace 1983, Myers 1980	British Columbia
	Green Heron	Rare	Johnson et al. 2000	
Black-crowned Night Heron	Recurrent	Johnson et al. 2000		
Great Egret	Recurrent	Johnson et al. 2000, Schlorff 1978	California	
Harlequin Duck	Strong	Cottam 1939, Mace 1983	British Columbia	
Surf Scoter	Rare	Mace 1983	British Columbia	
White-winged Scoter	Rare	Mace 1983	British Columbia	
Common Goldeneye	Rare	Mace 1983	British Columbia	
Barrow's Goldeneye	Rare	Mace 1983	British Columbia	

**SALTWATER –
SMOLT,
IMMATURE
ADULTS AND
ADULTS**

Common Merganser	Strong	Elson 1962	Canada
		Mace 1983, Macdonald et al. 1988, Wood and Hand 1985, Wood 1985, 1986, 1987	British Columbia
Red-breasted Merganser	Recurrent	Mace 1983	British Columbia
Osprey	Strong	Bayer 1986a, Emmett 1997, Roby et al. 1998	Oregon
Bald Eagle	Recurrent	Knight et al. 1990, Watson et al. 1991	Washington
Bonaparte's Gull	Recurrent	Macdonald et al. 1988, Mace 1983	British Columbia
Heermann's Gull	Recurrent	Bayer 1986a, 1989	Oregon
Ring-billed Gull	Recurrent	Bayer 1989, Roby et al. 1998	Oregon
		Ruggerone 1986	Washington
California Gull	Recurrent	Roby et al. 1998	Oregon
		Mace 1983	British Columbia
Herring Gull	Recurrent	Bent 1921	Washington
		Mace 1983, Macdonald et al. 1988	British Columbia
Thayer's Gull	Recurrent	Mace 1983	British Columbia
Western Gull	Recurrent	Bayer 1986a, 1986b, Roby et al. 1998	Oregon
Glaucous-winged Gull	Recurrent	Baird 1990	Washington
		Roby et al. 1998, Bayer 1986a	Oregon
		Mace 1983, Vermeer 1982	British Columbia
Glaucous Gull	Recurrent	Sanger 1983	Alaska
Black-legged Kittiwake	Rare	Rowlett 1980, Sanger 1983	Alaska
		Simenstad et al. 1979	Oregon
Caspian Tern	Strong	Smith and Mudd 1978	Washington
		Roby et al. 1998	Oregon
Elegant Tern	Recurrent	Johnson et al. 2000	
Common Tern	Recurrent	Johnson et al. 2000, Simenstad et al. 1979	Oregon
Forster's Tern	Recurrent	Johnson et al. 2000	
Arctic Tern	Recurrent	Simenstad et al. 1979	Oregon
		Willson and Halupka 1995	Alaska
Common Murre	Recurrent	Sydeman et al. 1996	California
		Bayer 1986a, Bayer 1986b, Matthews 1983	Oregon

**SALTWATER –
SMOLT,
IMMATURE
ADULTS AND
ADULTS**

		The OR Inst. Of Marine Biol. 1982	Oregon
		Ainley et al. 1990	Alaska
Pigeon Guillemot	Rare	Bayer 1986a	Oregon
Marbled Murrelet	Recurrent	DeGange 1996	Alaska
		Mace 1983	British Columbia
Ancient Murrelet	Rare	Mace 1983	British Columbia
Rhinoceros Auklet	Recurrent	Wilson and Manuwal 1986	Washington
		Burger et al. 1993, Vermeer and DeVito 1986, Vermeer and Westrheim 1984, Vermeer 1979	British Columbia
		Gaston and Dechesne 1996, Sydeman et al. 1997	California
		Sanger 1983	Alaska
Tufted Puffin	Recurrent	Baird 1990, 1991, Wehle 1983	Alaska
Belted Kingfisher	Recurrent	Bayer 1989	Oregon
		Mace 1983	British Columbia
Northwestern Crow	Recurrent	Mace 1983	British Columbia
Northern Fur Seal	Recurrent	Antonelis an Perez 1984, Baker et al. 1970, Kajimura 1980	Washington,
		Clemens and Wilby 1933, Spalding 1964	British Columbia
		Scheffer 1950, Wilke and Kenyon 1957	Bering Sea
		Banfield 1974	Canada
		Fiscus and Baines 1966, Imler and Sarber 1947, Pitcher 1981	Alaska
		Dalquest 1948	Washington
		Riemer and Brown 1997, Roffe and Mate 1983	Oregon
		Banfield 1974, Spalding 1964	British Columbia
California Sea Lion	Recurrent	Gearin et al. 1988, Jeffries 1984	Washington
		Jameson and Kenyon 1977, Riemer and Brown 1997, Roffe and Mate 1983	Oregon
		Baltz and Morejohn 1978, Harvey and Weise 1997, Jones 1981, NMFS 1997	California
Harbor Seal	Recurrent	Dalquest 1948, Everitt et al. 1981, Sheffer and Sperry 1931, Scheffer and Slipp 1944	Washington

SALTWATER – SMOLT, IMMATURE ADULTS AND ADULTS			Beach et al. 1985, Brown and Mate 1983, Brown et al. 1995, Browne et al. 1997, Graybill 1981, Jeffries 1985, Riemer and Brown 1997, Roffe and Mate 1984	Oregon
			Olesiuk 1993, Olesiuk et al. 1990, Spalding 1964	British Columbia
			Imler and Sarber 1947, Pitcher and Calkins 1981, Pitcher 1977, 1981	Alaska
			Briggs and Davis 1972, Jones 1981, Herder 1983, Hanson 1993	California
	Minke Whale	Rare	Banfield 1974	Canada
			Stewart and Leatherwood 1985	Atlantic Ocean
	Humpback Whale	Rare	Johnson and Wolman 1984	Northern Hemisphere
	Pacific White-sided Dolphin	Recurrent	Kajimura et al. 1980, Stroud et al. 1980	Washington
	Northern Right-Whale Dolphin	Rare	Johnson et al. 2000	
	Killer Whale	Strong	Hall 1986	Alaska
			Scheffer and Slipp 1948	Washington
			Nichol and Shackleton 1996, Banfield 1974	British Columbia
			Balcomb et al. 1982, Heimlich-Boran 1986, 1987, Felleman et al. 1991	Pacific Northwest
	Harbor Porpoise	Rare	Gearin et al. 1994	Washington
			Fontaine et al. 1994	Canada
	Dall's Porpoise	Rare	Norris and Prescott 1961	Oregon
		Mizue et al. 1996	Washington	
Sperm Whale	Rare	Leatherwood and Reeves 1983, Pike 1950	British Columbia	
Osprey	Strong	Johnson et al. 2000		
Bald Eagle	Strong	Hunt et al. 1992, Servheen 1975, Spencer et al. 1989, Stalmaster 1976	Washington	
		Simons 1983	Oregon	

SPAWNING			Munro 1938a	British Columbia
			Ofelt 1975	Alaska
	Golden Eagle	Recurrent	McClelland 1973	Montana
	Glaucous-winged Gull	Recurrent	Mossman 1958	Alaska
	Belted Kingfisher	Recurrent	Reimchen 1994	British Columbia
	Common Raven	Recurrent	Reimchen 1994	British Columbia
	Gray Wolf	Recurrent	Young 1944	Alaska
	Black Bear	Strong	Kellyhouse 1975	California
			Reimchen 1994	British Columbia
			Mattson 1989	Idaho
			Chi 1999, Moyle 1966, Piorkowski 1995, Wilson et al. 1998, Frame 1974	Alaska
			Banfield 1974	Canada
	Grizzly Bear	Strong	Banfield 1974	Canada
	Mink	Recurrent	Eagle and Whitman 1987	Canada
			Melquist et al. 1981	Idaho
			Hatler 1976	British Columbia
	Northern River Otter	Strong	Melquist et al. 1981	Idaho
			Toweill 1974	Oregon
	Mountain Lion	Rare	Ned Pittman pers. comm.	Washington
	Bobcat	Recurrent	Yoakum 1964	Washington
	Habor Seal	Recurrent	Reimchen 1994	British Columbia
Everitt et al. 1981			Washington	
Riemer and Brown 1997			Oregon	
California Sea Lion	Recurrent	Jeffries 1984	Oregon	
		Riemer and Brown 1997, Roffe and Mate 1983	Oregon	
Northern (Steller) Sea	Recurrent	Roffe and Mate 1983, Riemer and Brown 1997	Oregon	
		Reimchen 1994	British Columbia	
CARCASSES	Western Pond Turtle	Rare	Holland 1985	California
	Common Loon	Rare	Reimchen 1994	British Columbia
	Western Grebe	Rare	Reimcher 1994	British Columbia

CARCASSES

Pacific Loon	Rare	Reimchen 1994	British Columbia
Red-necked Grebe	Rare	Reimchen 1994	British Columbia
Turkey Vulture	Recurrent	Jewett et al. 1953	Washington
California Condor	Recurrent	Gabrielson and Jewett 1940, Simons 1983	Oregon
		Suckley and Cooper 1860	Washington
Trumpeter Swan	Rare	Butler 1973	British Columbia
Mallard	Rare	Jewett et al. 1953	Washington
		Reimchen 1994	British Columbia
Canvasback	Rare	Jewett et al. 1953	Washington
Greater Scaup	Rare	Munro 1941, Reimchen 1994	British Columbia
Surf Scoter	Rare	Reimchen 1994	British Columbia
White-winged Scoter	Rare	Reimchen 1994	British Columbia
Common Goldeneye	Recurrent	Dawson 1909, Jewett et al. 1953, Servheen 1975	Washington
		Taverner 1934	Canada
Barrow's Goldeneye	Recurrent	Jewett et al. 1953	Washington
Hooded Merganser	Rare	Reimchen 1994	British Columbia
Common Merganser	Recurrent	Stalmaster and Gessaman 1984	Washington
		Munro and Clemens 1932, 1937, 1939	British Columbia
Bald Eagle	Strong	Cederholm et al. 1989, Jewett et al. 1953, Knight and Knight 1983, Skagen et al. 1991, Stalmaster and Gessaman 1984, Owen et al. 1990	Washington
		Simons 1983	Oregon
		Munro 1938a	British Columbia
		Frame 1974	Alaska
		Shea 1970	Montana
Red-tailed Hawk	Rare	Willson and Halupka 1995	Alaska
		Cederholm et al. 1989, Ned Pittman pers. comm., Stalmaster and Gessaman 1984, Stalmaster 1980, Nui Tateyama, pers. comm.	Washington
Golden Eagle	Recurrent	Stalmaster and Gessaman 1984	Washington
Ring-billed Gull	Recurrent	Johnson et al. 2000	
California Gull	Recurrent	Johnson et al. 2000, Reimchen 1994	British Columbia

CARCASSES

Bonaparte's Gull	Recurrent	Moyle 1966, Willson and Halupka 1995, Frame 1974	Alaska
Western Gull	Recurrent	Johnson et al. 2000	
Herring Gull	Recurrent	Reimchen 1994	British Columbia
Glaucous-winged Gull	Recurrent	Reimchen 1994	British Columbia
		Simons 1983	Oregon
		Moyle 1966, Bent 1921, Frame 1974	Alaska
		Servheen 1975, Skagen et al. 1991, Stalmaster and Gessaman 1984	Washington
Glaucous Gull	Recurrent	Johnson et al. 2000	
Black-legged Kittiwake	Rare	Reimchen 1994	British Columbia
Gray Jay	Rare	Cederholm et al. 1989	Washington
Steller's Jay	Recurrent	Cederholm et al. 1989	Washington
		Willson and Halupka 1995, Frame 1974	Alaska
Black-billed Magpie	Recurrent	Willson and Halupka 1995	Alaska
		McClelland 1973	Montana
American Crow	Recurrent	Cederholm et al. 1989, Skagen et al. 1991, Stalmaster and Gessaman 1984	Washington
Northwestern Crow	Recurrent	Frame 1974	Alaska
		Campbell et al. 1990, Reimchen 1994	British Columbia
Common Raven	Recurrent	Murie 1959, Willson and Halupka 1995, Frame 1974	Alaska
		Knight and Anderson 1990, Stalmaster and Gessaman 1984, Cederholm et al. 1989, Suckley and Cooper 1860	Washington
Winter Wren	Rare	Cederholm et al. 1989	Washington
		Reimchen 1994	British Columbia
American Dipper	Recurrent	Willson and Halupka 1995	Alaska
		Cederholm et al. 1989	Washington
Varied Thrush	Rare	Reimchen 1994	British Columbia
Spotted Towhee	Rare	Ned Pittman pers. comm.	Washington
Song Sparrow	Rare	Ned Pittman pers. comm.	Washington
Virginia Opossum	Recurrent	Johnson et al. 2000	
Deer Mouse	Rare	Cederholm et al. 1989	Washington

Douglas' Squirrel	Rare	Cederholm et al. 1989	Washington
Northern Flying Squirrel	Rare	Cederholm et al. 1989	Washington
Water Shrew	Recurrent	Cederholm et al. 1989	Washington
		Conaway 1952	Montana
Vagrant Shrew	Rare	Cederholm et al. 1989	Washington
Masked Shrew	Rare	Cederholm et al. 1989	Washington
Trowbridge's Shrew	Rare	Johnson et al. 2000	
Pacific Water Shrew	Rare	Johnson et al. 2000	
Pacific Shrew	Rare	Johnson et al. 2000	
Montane Shrew	Rare	Johnson et al. 2000	
Fog Shrew	Rare	Johnson et al. 2000	
Coyote	Recurrent	Young 1944	Oregon
		Cederholm et al. 1989, Stalmaster and Gessaman 1984	Washington
Gray Wolf	Recurrent	Young 1944	Oregon, British
Red Fox	Rare	Young 1944	Oregon
		Willson and Halupka 1995	Alaska
		Hewson 1995	Scotland
Gray Fox	Rare	Young 1944	Oregon
Black Bear	Strong	Young 1944	British Columbia
		Piorkowski 1995, Chi 1999, Frame 1974	Alaska
		Cederholm et al. 1989, Hilderbrand et al. 1996	Washington
Grizzly Bear	Strong	Mattson et al. 1991	Idaho
		Hamilton and Archibald 1985, Hamilton and Bunnell 1987, Young 1944	British Columbia
		Banfield 1974	Canada
		Verts and Carraway 1998	Oregon
Ringtail	Rare	Johnson et al. 2000	
Raccoon	Recurrent	Cederholm et al. 1989	Washington
American Marten	Rare	Reimchen 1994, Nagorsen et al. 1989, 1991, Hatler 1976	British Columbia
Striped Skunk	Rare	Cederholm et al. 1989	Washington
Long-tailed Weasel	Rare	Cederholm et al. 1989	Washington