Sunnyside-Snake River Wildlife Area Management Plan



Acknowledgements

Planning team members

Scott McCorquodale, Region 3 Wildlife Program Manager Ross Huffman, Region 3 Lands Operations Manager Phillip Buser, Former Wildlife Area Manager Pat Kaelber, Wildlife Area Manager Noel Ferguson, Wildlife Area Assistant Manager Jorge Garcia, Water Access Manager Mike Ritter, District Habitat Biologist Jason Fidorra, District Wildlife Biologist Paul Hoffarth, District Fish Biologist Sergeant Brian Fulton, Enforcement Program Mark Teske, Diversity Species Biologist

Plan leadership and content development

Scott McCorquodale, Region 3 Wildlife Program Manager Ross Huffman, Region 3 Lands Operations Manager Phillip Buser, Former Wildlife Area Manager Pat Kaelber, Wildlife Area Manager Patricia Jatczak, Lands Planner Cynthia Wilkerson, Lands Division Manager

Other WDFW Support

John Talmadge, GIS Shelly Snyder, GIS Brian Cosentino, GIS Rachel Blomker, Public Affairs Madonna Luers, Public Affairs*, retired* Matthew Wilson, Statewide Waterfowl Specialist Lynn Helbrecht, Climate Change Coordinator

Cover photo: Sumac and goldenrod on Sunnyside-Snake River Wildlife Area Photo by Justin Haug



Flicker and robin Photo by Justin Haug, WDFW

Wildlife Area Advisory Committee

Sunnyside-Snake River Wildlife Area Advisory Committee Roster Ross Huffman, WDFW Staff Lead

Name	Representing
Mike Estes	Richland Rod and Gun Club
Suzanne Sullivan	Ducks Unlimited
Randy Cline	Pheasants Forever
Kathy Criddle	Lower Columbia Basin Audubon / Native Plant Society
Betsy Crysel	Franklin County Noxious Weed Control Board
Jeff Knutson	Yakima County Noxious Weed Control Board
Jed Pauley	Agricultural Producer
Rachel Voss	Mule Deer Foundation
Craig Adams	Rocky Mountain Elk Foundation
Dave Solem	South Columbia Basin Irrigation District
Danny Chappel	Backcountry Horsemen's Association
Dave Blodgett	Yakama Nation
Dennis Dauble	Columbia Basin Fly Casters Association
Vic Reeve	Benton County Noxious Weed Control Board



Snow geese Photo by Alan L. Bauer





Great horned owl Photo by Justin Haug, WDFW

Sunnyside-Snake River Wildlife Area Management Plan

unni line

Kelly Susewind, Director, Washington Department of Fish and Wildlife



Table of contents

Acknowledgements	1
Wildlife Area Advisory Committee	2
Tables	6
Figures	6
Acronyms	7
Wildlife area management planning overview	8
Introduction and agency mission	8
Wildlife area management planning framework	8
Purpose and organization of the plan	8
Public outreach and stakeholder involvement process	9
Statewide WDFW lands vision	9
Statewide planning goals	10
Welcome to the Sunnyside-Snake River Wildlife Area	11
Introduction	11
Wildlife Area Description	13
Sunnyside-Snake River Wildlife Area Overview	13
Wildlife area vision	14
Success Stories	17
Sunnyside-Snake River Wildlife Area Unit Descriptions	20
Land Ownership and Management, Funding and Agreements	73
Acquisition history, funding, and purpose	73
Management setting	76
Administration	76
Recreation and stewardship	
Wildlife area goals, objectives, and monitoring	
Goals, objectives and performance measures	
Monitoring and adaptive management	
Physical characteristics	
Geology and soils	
Hydrology and watersheds	
Climate	



Ecological values	
Species management	113
Diversity Species	
Fish species overview and management	126
Habitat management	
Climate change approach	132
References	138
Appendices	139
Appendix A. Species and habitat information	140
Appendix B. Weed management plan	142
Appendix C. Fire response information	146
Appendix D. Sunnyside-Snake River Wildlife Area Wetland Management Plan	149
Appendix E. Cultural History Summary	
Appendix F. Public Response Summary (SEPA)	164



Tables

- Table 1Statewide wildlife area planning goals
- Table 2Wildlife area land instruments
- Table 3 Agricultural leases
- Table 4 Land Use designation by wildlife area unit
- Table 5Research activities
- Table 6 Recreational use
- Table 7Water access areas
- Table 8On-going volunteer opportunities
- Table 9
 Goals, objectives, and performance measures
- Table 10 Ecological systems of concern
- Table 11 Species conservation status
- Table 12 Fish species
- Table 13Focal fish species in Yakima River
- Table 14 Fire history



Shrike at Sunnyside Unit Photo by Justin Haug, WDFW

- Table 15
 Species of greatest conservation need, vulnerability, and confidence
- Table 16
 Species associated with shrubsteppe and with a sensitivity to climate change
- Table 17Plan objectives with climate nexus

Figures

- Figure 1 Sunnyside-Snake River Wildlife Area Vicinity West
- Figure 2 Sunnyside-Snake River Wildlife Area Vicinity East
- Figure 3 I-82 Ponds (Union Gap)
- Figure 4 I-82 Ponds (Wapato)
- Figure 5 I-82 Ponds (Zillah)
- Figure 6 Glover Unit
- Figure 7 Sunnyside Unit
- Figure 8 Vance-Ferry Road Unit
- Figure 9 Byron Unit
- Figure 10 Whitstran Unit
- Figure 11 Thornton Unit
- Figure 12 Rattlesnake Slope Unit
- Figure 13 Benton City Unit
- Figure 14 Hope Valley Unit
- Figure 15 Mesa Lake Unit
- Figure 16 Esquatzel Coulee Unit
- Figure 17 Windmill Ranch Unit
- Figure 18 Bailie Unit
- Figure 19 Thompson Seeps
- Figure 20 Fire District Boundaries West
- Figure 21 Fire District Boundaries East



Acronyms

ADA	Americans with Disabilities Act
BLM	Bureau of Land Management
BMP	Best Management Practices
BOR	Bureau of Reclamation
BPA	Bonneville Power Administration
CRP	Conservation Reserve Program
DAHP	Washington State Department of Archaeology & Historic Preservation
DNR	Washington State Department of Natural Resources
EIA	Ecological Integrity Assessment
EIM	Ecological Integrity Monitoring
ESA	Endangered Species Act
IPM	Integrated Pest Management
NPS	National Park Service
NRCS	National Resources Conservation Service
PHS	Priority Habitats and Species
RCW	Revised Code of Washington
RCO	Washington State Recreation and Conservation Office
SEPA	State Environmental Policy Act
SGCN	Species of Greatest Conservation Need
SRFB	Salmon Recovery Funding Board
SSSRMU	Sunnyside-Snake River Management Unit
SWAP	State Wildlife Action Plan
USACE	United States Army Corps of Engineers
USFS	United States Forest Service
USFWS	United States Fish and Wildlife Service
WAC	Washington Administrative Code
WAAC	Wildlife Area Advisory Committee
WDFW	Washington State Department of Fish and Wildlife
WHCWG	Washington Wildlife Habitat Connectivity Working Group
WLA	Wildlife Area
WWRP	Washington Wildlife and Recreation Program



Wildlife area management planning overview

Introduction and agency mission

Welcome to your fish and wildlife lands!

The Washington Department of Fish and Wildlife (WDFW) provides active management for more than 1 million acres of publicly owned land, most of which falls within 33 wildlife areas across the state (https://wdfw.wa.gov/places-to-go/wildlife-areas). These diverse lands contain nearly all species and habitats present in Washington. With the loss of natural habitat posing the single greatest threat to native fish and wildlife, these wildlife areas play a critical conservation role. The wildlife area management plan addresses all aspects of resource management, highlights areas for public access, education, and stewardship, and aligns with statewide conservation goals.

In addition to protecting lands and water for habitat and people, WDFW manages land to preserve Washington's natural and cultural heritage, provide access for hunting, fishing, and wildlife-related recreation, and to foster outdoor experiences and exploration throughout the state. We do this to support the species and habitats of Washington and ensure they prosper for our collective enjoyment well into the future.

An interdisciplinary team of WDFW staff members, including fish, habitat, and wildlife biologists, as well as enforcement, real estate, and management, developed the Sunnyside-Snake River Wildlife Area Management Plan along with significant public involvement. This included input from the local stakeholder-based Sunnyside-Snake River Wildlife Area advisory committee, tribes, public agencies, and interested residents.

Wildlife area management planning framework

Management of wildlife areas is guided by WDFW's mission and strategic plan, as well as by state and federal laws. Each new plan is guided by the Wildlife Area Management Planning Framework (framework), which summarizes the agency's mission, laws, policies, and approaches to management of fish and wildlife, as well as public use and recreation. The framework summarizes priorities and guidance developed in each of the agency's program—Fish, Wildlife, Habitat, and Enforcement. Readers are encouraged to review the framework in advance, or as a companion document to this wildlife area plan. The framework (<u>https://wdfw.wa.gov/publications/01810</u>) provides context for the organization and content of wildlife area plans across the state. The framework is a living document, and is updated periodically to reflect new agency initiatives, guidance, and directives.

Purpose and organization of the plan

The purpose of this management plan is to guide all management activities, including conservation and recreation, occurring on the Sunnyside-Snake River Wildlife Area for the next 10 years. Management goals, objectives, and performance measures are defined in the plan and provide a clear road map of projects and management actions to support statewide conservation and



recreation goals. Actions in the plan are dependent on available budget. Budget reductions made during the life of this plan may delay implementation of some actions.

The plan is designed to be a resource for internal and external audiences, and is organized into four parts: 1) overview of the wildlife area and associated units and success stories; 2) goals and objectives, and performance measures for the planning area; 3) environment information, wildlife species, and habitat management; and 4) appendices which support different areas of the plan.

Public outreach and stakeholder involvement process

The agency is committed to a transparent and inclusive public outreach process for all wildlife area management plans. Under the umbrella of the statewide planning goals (Table 1), a customized outreach strategy was developed for this area, tailored to local and regional stakeholders, as well as



Fishing at I-82 Ponds Photo by Alan L. Bauer local and out-of-the-area visitors and user groups. For this plan, the public process included three elements: 1) tribal, public, and advisory committee meetings; 2) development and distribution of fact sheets, meeting announcements, and news releases; and 3) solicitation of public comments through meetings, phone calls, email, social media, and the WDFW website. Comments on the Final Draft Plan were solicited through the State Environmental Policy Act (SEPA) process. The Public Response Summary for this is included in Appendix F.

Statewide WDFW lands vision

The statewide vision sets the agency expectations for the future state of all Washington Department of Fish and Wildlife lands.

A Washington where fish and wildlife thrive in healthy habitats, and where people experience and enjoy our state's natural gifts for generations to come.

By actively managing lands, restoring habitats, and preserving wild places, we serve as stewards and guardians for Washington's natural places by protecting lands and water for wildlife and people.



Statewide planning goals

A complete list of goals, objectives, and performance measures specific to this wildlife area starts on page 90.

Goal 1	Restore and protect the integrity of priority ecological systems and sites. This goal originates from the 2017-2019 WDFW Strategic Plan, Goal #1: "Conserve and protect native fish and wildlife". Ecological integrity monitoring on priority systems and sites will be developed as part of implementation of the management plan for each individual wildlife area plan.
Goal 2	Sustain individual species through habitat and population management actions, where consistent with site purpose and funding. This goal also relates to 2017-19 WDFW Strategic Plan, Goal #1. Each individual wildlife area plan will provide a summary of species associated with the wildlife area and will focus on target species for habitat management actions.
Goal 3	Provide fishing, hunting, and wildlife-related recreational opportunities where consistent with Goals 1 and 2. This goal is consistent with the WDFW Strategic Plan, Goal #2: "Provide sustainable fishing, hunting, and other wildlife-related recreational and commercial experiences". Each plan will provide a summary of recreation activities associated with the wildlife area, aiming to balance recreational activities with species and habitat protection.
Goal 4	Engage stakeholders in consistent, timely, and transparent communication regarding wildlife area management activities. This goal relates to Strategic Plan Goal #3: "Promote a healthy economy, protect community character, maintain an overall high quality of life, and deliver high-quality customer service". As described under the public outreach section of this document, public input and involvement are key components in the development of the management plan through the advisory committee efforts and public meetings. After the plan is adopted, the management plan updates will be reviewed by the wildlife area advisory committee on a biannual basis.
Goal 5	Maintain productive and positive working relationships with local community neighbors, lessee partners, and permittees. As part of day-to-day business, wildlife area staff strives to maintain positive working relationships with grazing and agricultural lessees and the local community.
Goal 6	Hire, train, equip, and license, as necessary, wildlife area staff to meet the operation and management needs of wildlife areas. This goal is consistent with Goal #4 of the Strategic Plan: "Build an effective and efficient organization by supporting the workforce, improving business processes, and investing in technology". Specific activities on wildlife areas include attending training and hiring qualified staff.
Goal 7	Maintain safe, highly functional, and cost-effective administration and operational facilities and equipment. This goal is consistent with WDFW Strategic Plan Goal #4. Maintenance of facilities and equipment is a key activity on wildlife areas. Annual reporting is required by WDFW and agencies that provide operations and maintenance funding, such as U.S. Fish and Wildlife Service, Pittman-Robertson.

Table 1: Statewide planning goals



Welcome to the Sunnyside-Snake River Wildlife Area



Thornton Unit Photo by Justin Haug, WDFW

Introduction

The Sunnyside-Snake River Wildlife Area is managed in 15 units in southcentral Washington on 21,200 acres in Yakima, Benton, and Franklin counties. The wildlife area units span roughly 100 miles from west to east, starting at the Sunnyside units (south of Union Gap along I-82), through the Benton City Unit along the Yakima River, and ending at the Snake River units (north of Pasco on SR 395).

Acquisitions or land management agreements began in 1944 and continued through 2017 to protect and enhance habitat and provide public recreation. The Snake River Units in Franklin County were purchased to provide partial mitigation for the construction of four dams on the Lower Snake River. The most recent acquisitions were three small parcels totaling 79 acres along the Yakima River, which were deeded to WDFW in 2017. WDFW owns 12,598 acres; 4,693 acres are either under agreement, leased, or are Bureau of Reclamation property managed by WDFW; and 3,909 acres are in easement, almost all of which is in one hunting easement.



Some of the Sunnyside units along the Yakima River upstream of the Mabton bridge (SR 241 crossing), border the Yakama Indian Reservation. The Confederated Bands and Tribes of the Yakama Nation partners with WDFW and other state and federal agencies and non-governmental organizations on activities that affect fish and wildlife and habitat. The 1855 Treaty with the Yakama says "The exclusive right of taking fish in all the streams, where running through or

bordering said reservation, is further secured to said confederated tribes and bands of Indians, as also the right of taking fish at all usual and accustomed places, in common with the citizens of the Territory, and of erecting temporary buildings for curing them; together with the privilege of hunting, gathering roots and berries, and pasturing their horses and cattle upon open and unclaimed land". Tribal members also harvest culturally important plants, such as wapato, camas, cattails, and taxus (dogbane). Treaty tribes have been recognized as managers of their treatyreserved resources, and have interest in the management decisions in the



Yakima River Photo by Alan L. Bauer

Sunnyside-Snake River Wildlife Area (US. v Washington 1974). WDFW honors and respects tribal treaty reserved rights, and will take into consideration traditional hunting and gathering sites in any actions in this plan that may affect them. Communicating and coordinating with tribes is an objective of this plan.

Wildlife area lands have been affected by past agricultural practices, fire, road construction, and general development. Some of these factors, such as fire, continue to impact some wildlife areas. Irrigation water provides both managed wetland habitats and a water source for limited agricultural operations on some units. These wetlands provide important habitat and riparian areas for a variety of resident and migratory wildlife. Non-irrigated land on the units provide habitat more characteristic of the semi-arid Columbia Basin and supports native bunch grasses and shrubsteppe vegetation. Overall, the wildlife area units support more native habitats than the surrounding landscapes that are characterized as mostly irrigated croplands.

The Sunnyside-Snake River units provide important native and managed (wetland) habitats in the Columbia Basin for both resident and migratory wildlife. Some units are managed under prior agreements with the Bonneville Power Administration (BPA) and US Army Corps of Engineers for upland wildlife, waterfowl, big game, and a variety of non-game species that are consistent with WDFW goals. The units that are owned by the Bureau of Reclamation (BOR) are managed under a separate agreement that is also consistent with WDFW goals. In this arid environment, these wetland and riparian areas provide habitat, browse, and cover for wildlife.

Close to one-half of the acreage is shrubsteppe. Open water, cultivated cropland, and riparian are the other main land covers. The area is popular for fishing, hunting, and bird watching.



Wildlife Area Description

Sunnyside-Snake River Wildlife Area Overview

Size	21,199 acres
Acquisition, Agreement Dates	1944 - 2017
Acquisition Funding and Agreement	National Park Service: Land and Water Conservation Fund US Fish and Wildlife Service: Pittman-Robertson Wildlife Restoration Program (PR) US Army Corps of Engineers: Snake River Mitigation Account US Congress Programs: Transfer of Land WA Recreation and Conservation Office: WA Wildlife and Recreation Program; Boating Facilities Program; State Bond Account Private Grantor: Private Donations WA Dept. of Fish and Wildlife: Wildlife Fund; State Migratory Waterfowl Fund Utility Districts: Mitigation Funds
Elevation Range	448 - 2,446 feet
Recreational Opportunities	Hunting, fishing, wildlife viewing, birding, walking, horseback riding
Units	West side: I-82 Ponds, Glover, Sunnyside, Vance-Ferry Road, Byron, Whitstran, Thornton, Benton City, and Rattlesnake Slope. East side: Hope Valley (<i>includes Clark Pond and Nipper</i>), Mesa Lake, Esquatzel Coulee, Windmill, Bailie, and Thompson Seeps.
Counties	Yakima, Benton, and Franklin

Just under 60% of the wildlife area is owned by WDFW, with the remainder managed by WDFW under a variety of agreements. Ownership does not play a large role in how lands are managed, unless there are restrictions or conditions within the agreements.

During the planning process for this management plan, a couple unit names were changed, and three small contiguous units were combined as one. The number of wildlife area properties organized by units went from 17 to 15. The Nipper, Clark Pond, and Hope Valley units were combined under the name of Hope Valley.

Previous Name	New Name
Headquarters (Sunnyside) Unit	Sunnyside Unit
WB-10 Wasteway	Thompson Seeps Unit
Nipper	Hope Valley
Clark Pond	Hope Valley
Hope Valley	Hope Valley

Figures 1 and 2 show the west and east locations of the wildlife area units.



Wildlife area vision

The overall vision for the Sunnyside-Snake River Wildlife Area is to restore, enhance, and maintain a variety of native and managed habitats to support a diversity of wildlife, which support fishing, quality waterfowl hunting and other hunting, and birding, hiking, and a variety of educational and nature-based activities.

The focus for recreational and educational activities will be on those units that support high visitor use or have the potential to support more activities through partnerships. These priority units are: Sunnyside, Rattlesnake Slope, Thornton, Mesa Lake, Windmill Ranch, and the Esquatzel Coulee.



Blue-winged teal Photo by Phillip Buser





Figure 1. Sunnyside-Snake River Wildlife Area Vicinity - West



Figure 2. Sunnyside-Snake River Wildlife Area Vicinity - East

Success Stories

Enhancing wetlands to improve habitat on the wildlife area

The Sunnyside-Snake River Wildlife Area contains of about 4,000 acres of freshwater wetland habitats. These wetland habitats are primarily supported by direct surface flow of irrigation water, but seeps and springs support wetlands that are not directly connected to the irrigation canals and waterways.

Since 2011, approximately 400 acres of wetland habitat have been restored and enhanced at the Mesa Lake, Windmill Ranch, Bailie, and Sunnyside units. The enhancement and restoration work included the replacement of water control structures, pumps, culverts, and piping, and beaver management. Beavers play an important role in the ecosystem, and their dam-building activities provide habitat for a diversity of species, but they can also cause problems. Beavers instinctively build dams where they hear the water channeled through a culvert. This can lead to flooding and road and other damage. To reduce the damage caused by beavers, sturdy wire mesh was used to prevent damming up the pipe. These "beaver deceivers" allow the water to keep flowing,



Worth Lake, Windmill Ranch Unit Photo by Alan L. Bauer

Restoration and enhancement projects targeted shallower wetlands where management actions promote open water and emergent wetland vegetation. This vegetation provides nesting and escape cover for water birds and waterfowl. These wetlands also provide year round habitat for a wide variety of wetland-dependent plant and animal species, such as muskrats, mallards, Canada geese, and river otters. Additionally, wetland restoration and enhancement provides more opportunities for hiking, bird watching, and quality waterfowl hunting. Wetlands throughout the

Sunnyside-Snake River Wildlife Area provide waterfowl and upland bird hunting opportunities for the public. Hunters come from around the state to these areas for quality hunting on public land. To maintain these wetlands and continue to provide for public recreation, all wetlands continue to be actively managed through seasonal mowing, seeding, and weed treatments.



Engaging youth and volunteers on the wildlife area

The Sunnyside-Snake River Wildlife Area has a long tradition of providing outdoor learning

experiences for local youth and school groups. Teachers value the opportunity for students to learn life skills outside of the classroom, and students learn about the environment and gain skills from real life examples. This collaboration is also good for wildlife. In 2010, students from Mabton Junior High School built bluebird nest boxes and helped place them throughout the Snipes Reserve of the Sunnyside Unit. For four years, students from Artz Fox Elementary School have helped plant 200 shrubs at the Sunnyside Unit each year. Species of shrubs planted include big sagebrush, woods rose, hawthorn, and wild current.

Another avenue of youth engagement on the wildlife area has been through the Boy Scouts of America, specifically their Eagle Scout Merit Award program. Since 2000, six scouts have conducted projects on the wildlife area, and three just in 2017-2018. One scout organized a work party to build five large bat boxes that were placed on the Windmill Ranch Unit. Another scout organized a work party to build three osprey nest platforms, which have been placed at the



School group learning wetland ecology Photo by Phillip Buser



Eagle Scout project Photo by Phillip Buser

Windmill Ranch and Mesa Lake units. The third scout organized a crew to replace safety zone signs that burned up during a recent fire on the Rattlesnake Slope Unit. The scouts also replaced signs on two and a half miles of fence line. The wildlife and the visitors appreciate and benefit from the efforts of the scouts!

You do not have to be a student, a scout, or a member of an organization volunteer. There are many opportunities for individuals to volunteer on the wildlife area (see page 85 for some ongoing opportunities). Individual volunteers have donated their time and talents to clean up parking lots, repair fences, paint over graffiti, and help fix roads. Volunteers have even helped build ADA-accessible hunting and viewing blinds.



Clinics teach youth about hunting

Youth can learn to be safe and successful hunters at the Sunnyside-Snake River Wildlife Area. In 2014, a fire swept across the Sunnyside Unit. Along with burning nearly 250 acres of wildlife habitat, the fire also destroyed one of the shop buildings. The shop was rebuilt and now offers an

indoor space for hosting youth hunting clinics, mentored hunts, and hunter education classes.

Since 2015, there have been six combined pheasant clinics and hunts (two each year) at the Sunnyside Unit. These events have had an average of 20 participants and 30 volunteers per clinic. WDFW also held a dove clinic and hunt in 2017 with 15 participants and 11 volunteers.

There have been two hunter education classes, and more will be held based on interest and availability. Students learn about wildlife conservation, how to handle firearms safely, and the rules and regulations of hunting and outdoor recreation.



Youth hunt clinic at Sunnyside Photo by Phillip Buser

Diversity of hunting opportunities

Upland bird hunting in native habitats and waterfowl hunting in managed wetlands are just two of the hunting opportunities found on the Sunnyside-Snake River Wildlife Area. This wildlife area is comprised of 15 units that stretch across three counties and span almost 100 miles. There are a variety of habitat types, including wetlands, upland grasslands, shrubsteppe, and riparian corridors



Waterfowl Photo by Phillip Buser

that provide suitable habitats for a variety of wildlife species, including deer, elk, upland game birds, waterfowl, and small game.

Big game hunting opportunities include elk, which are routinely found on four different units in Benton and Franklin counties, and mule deer that occur on nearly every unit of the wildlife area. Wild turkeys are found in the units along the Yakima River corridor. Waterfowl hunting opportunities include ducks (except sea ducks) and geese (Canada, snow, and white-fronted) that can number in the thousands during the fall and

winter. The wildlife area provides significant habitat for wintering waterfowl. Hunting opportunities for mourning doves, coots, and snipe occur at many of the units. Upland hunting opportunities include California quail, ring-necked pheasant, coyotes, crows, and cottontail rabbits. Hunting for ring-necked pheasants is popular for both pen-raised birds released at the Hope Valley and Sunnyside units, and wild birds at the Bailie, Esquatzel Coulee, Mesa Lake, Thompson Seeps, and Windmill Ranch units. The Richland Rod and Gun Club has hosted pheasant clinics at Esquatzel Coulee in the past.

Sunnyside-Snake River Wildlife Area Unit Descriptions

I-82 Ponds

Size	804 acres
Acquisition Date	1963 - 2017
Acquisition Funding	US Army Corps of Engineers: Snake River Mitigation Account National Park Service: Land and Water Conservation Fund WA Recreation and Conservation Office: Boating Facilities Program; State Bond Account WA Dept. of Fish and Wildlife: Wildlife Fund; State Migratory Waterfowl Fund Utility Districts: Mitigation Funds
Purpose of Funding	Fishing, waterfowl habitat, small game habitat
Elevation Range	646 - 989 feet
Recreational Opportunities	Fishing, waterfowl and upland bird hunting, bird watching
County	Yakima
Site Access	Accessed from Donald Road (exit 44 from I-82) and Buena Loop (exit 50 from I-82)



Fishing at I-82 Ponds Photo by Alan L. Bauer



Overview

The I-82 Ponds Unit consists of 804 acres of wetland/riparian and upland habitats located along the Yakima River in Yakima County, in the lower Yakima River watershed. The entire unit is made up of 17 separate land parcels stretching from Union Gap to Zillah. The I-82 Unit is displayed in three sections: Union Gap, Wapato, and Zillah (Maps 3, 4, and 5). The land is owned by WDFW except for less than 10 acres in easement. Two small inholdings, the McDonald and Kroll properties, were added to the unit in 2017. These properties support the goal of the Draft Wapato Reach Action Plan (2017) to acquire land for conservation and to restore riparian and floodplain functions. Surrounding lands are primarily agriculture.

Overall, ponds and riverine wetlands comprise about 25% of this unit. The healthy riparian habitat, all of which transitions into upland shrubsteppe habitat, is dominated by cottonwood, various species of willow, Wood's rose, Nootka rose, blue elderberry, as well as introduced cultivars like mulberry trees. The backwater sloughs and dense riparian growth offer excellent nesting and food resources for waterfowl, upland birds, neotropical migrants, and birds of prey. The majority of the open water wetland habitat is associated with the seven ponds that range in size from 13 to 28 acres. Several of the ponds were formed when gravel borrow pits were created during the construction of the interstate and later filled with groundwater.

A primary focus of this unit is to provide facilities and public access to the Yakima River and ponds. There are 10 river access sites and seven ponds. The recreation section includes a summary of fishing and boating access for details on what is available at each site.

The ponds are popular fishing areas and offer a variety of fishing opportunities that include both bank fishing and small boat fishing. The ponds support populations of sunfish, black crappie, yellow perch, and largemouth bass, and periodically other fish species such as channel catfish are stocked. Legal sized rainbow trout are also stocked in most of the ponds each year.

Maintenance and restoration of some of the ponds is being addressed through a larger stakeholder process associated with the Yakima River Basin Integrated Water Resource Management Plan. Specifically, Pond 4, is separated by a failing levee to the downstream Pond 5. The location of ponds 4 and 5 adjacent to the Yakima River lies in the Wapato Reach, a section of the Yakima River that is a critical migration zone for salmon and steelhead, and provides spawning and rearing habitat for fall and summer Chinook salmon. An analysis was conducted by Ducks Unlimited on the feasibility of making Pond 5 open to the river and protecting Pond 4. Funding has been sought to do this work.

Columbia clubtail (a state species of concern), and white-belted ringtail dragonfly may occur here. Bull trout and steelhead, both federally threatened species in this area, and Pacific lamprey and mountain sucker utilize the Yakima River. This unit also sustains important cultural values to the Yakama Nation. A portion of the Yakima River is within the borders of the Yakama Indian Reservation, and salmon fishing in this reach is a valuable cultural resource to the Yakama Nation. Public salmon fishing is rarely open along this stretch of river, but other fish species can be pursued. A Yakama Nation fishing license is required, as well as a WDFW fishing license. Anglers must follow the rules and regulations of the Yakama Nation. To hunt the Yakima River, hunters need a Yakama Nation hunting permit as well as a WDFW hunting license.

The I-82 Unit offers waterfowl as well as deer, Eurasian collared dove, mourning dove, quail, pheasant, and small game. Again, waterfowl hunters in or on the river must be licensed by and



follow the rules and regulations of the Yakama Nation. There are multiple parking areas and restrooms on the unit.

Primary management objectives on the I-82 Ponds Unit

- Coordinate with Yakima Basin Integrated Plan (YBIP) work on Wapato Reach, which includes the I-82 ponds, as well as other areas of the Yakima River system (1.E).
- Maintain and restore floodplain functions throughout the Yakima River Valley (1.I).
- Develop regularly scheduled waterfowl surveys for use in assessing management actions (2.B).
- Assess potential to improve lakes and stock with fish (trout and/or warm water) by 2024 (4.A).
- Research areas where possible access or access agreements need to be updated by 2024 (5.B).

Figure 3: I-82 Ponds Unit (Union Gap)



Figure 4: I-82 Ponds Unit (Wapato)



Figure 5: I-82 Ponds Unit (Zillah)



Glover Unit

Size	51 acres
Acquisition Date	1992
Acquisition Funding	WA Recreation and Conservation Office: WA Wildlife and Recreation Program
Purpose of Funding	Small game and upland birds
Elevation Range	669 – 697 feet
Recreational Opportunities	Upland bird and waterfowl hunting
County	Yakima
Site Access	From Highway I-82, exit 67 to Wendell-Phillips Road

Overview

The Glover Unit is composed of 51 WDFW-deeded acres surrounded by private agricultural land, located southwest of Sunnyside in Yakima County. The property was acquired for upland bird and small game habitat. About 10 acres were historically used in an agricultural lease. The unit is a mosaic of hardwood trees, shrubs, and grasslands with a slough running east-west through the center. The mixture of trees and edge habitat with the added water component make it a surprisingly diverse patch of land within a farmland monoculture of corn and dairies. Mule deer, dabbling ducks, songbirds, birds of prey, doves, quail, and monarch butterflies can all be found in this small unit.

Recreational opportunities are some upland bird and waterfowl hunting, birding, and nature observation. There is a gravel parking area and information kiosk. Visitors must register on site when arriving, and report their activity before leaving. Registration provides usage and harvest data for future management of the unit.



Tree snags at Glover Unit Photo by Alan L. Bauer

Primary management objectives on the Glover Unit

- Manage and improve upland bird habitat and nesting cover (4.B).
- Ongoing, as agricultural leases come up for renewal or for new ones developed, evaluate agricultural leases per agency policy for multiple benefits, especially to improve waterfowl forage for hunting opportunities (7.B).

Figure 6: Glover Unit





Sunnyside Unit

Size	2,741 acres
Acquisition Date	1944 - 2017
Acquisition Funding	National Park Service: Land and Water Conservation Fund US Fish and Wildlife Service: Pittman-Robertson Wildlife Restoration Program (PR) US Army Corps of Engineers: Snake River Mitigation Account US Congress Programs: Transfer of Land WA Recreation and Conservation Office: WA Wildlife and Recreation Program; Boating Facilities Program; State Bond Account WA Dept. of Fish and Wildlife: Wildlife Fund; State Migratory Waterfowl Fund Private Grantor: Donation
Purpose of Funding	Waterfowl and small game habitat, riparian and floodplain habitat conservation
Elevation Range	632 - 681 feet
Recreational Opportunities	Waterfowl and upland bird hunting, small game and deer hunting, and bird watching
County	Yakima
Site Access	From Highway I-82, exit 67 – six parking areas



Giffin Lake, Sunnyside Unit Photo by Alan Bauer



Overview

This unit was formerly called the Headquarters – Sunnyside Unit, but was recently changed to the Sunnyside Unit. It covers 2,741 of mostly deeded acres, one mile north of Mabton and five miles south of Sunnyside. This unit is a collection of small agricultural fields, interspersed with diverse habitats. About 90% of the unit is in the floodplain of the Yakima River, with 13 miles of riverfront. Evidence of old river oxbows can be found throughout the unit. Vegetation ranges from wetland species to upland perennial grasses and forbs to mature riparian woodlands. Non-native Russian olive trees have been invading for several years, and staff manage them to reduce the spread. In 2017, a small inholding, the Grow property, was purchased and became part of the unit. This property was named in the Yakima River Basin Integrated Water Resource Management Plan (Sandisen et al 2012), and also supports the goal of the Wapato Reach Assessment Report (Yakama Nation 2012) to acquire land for riparian and floodplain habitat conservation.

In 2014, two fires, two weeks apart, burned into the unit. The fires burned large swaths of bunchgrasses and greasewood, and burned down the Unit's maintenance shop building. Funds were provided for replacing the office and shop and for the restoration of 250 acres of habitat. Restoration included weed treatments, seeding of native grasses, and shrub plantings. In 2015, a new office and shop facility were built, and habitat restoration projects continue.

The flora of the Sunnyside Unit varies widely, ranging from dense riparian thickets bordering the wetlands and Yakima River, to upland expanses of bunchgrasses and greasewood, to isolated stands of aspen, cottonwoods, and willows. Invasive, exotic plant species are common throughout the unit with new loads of seed being delivered almost annually with spring flooding from the river. Weed control is a continual effort on the unit as well as encouraging growth of native and wildlife-friendly plant species. In recent years, small sections of the historical agricultural fields that were planted in the floodplain were taken out of production and planted with native grasses and shrubs. The remaining agricultural fields are dryland agriculture.

Irrigation water supplies virtually all of the water to wetland habitats at this unit. The water level at the wetlands are managed through water control structures to provide wildlife habitat and forage, bird watching, and waterfowl hunting. Irrigation water is diverted to fill nearly all of the wetlands at the unit, with natural springs and sub-irrigation from the nearby Yakima River supplementing the remainder. The Sunnyside Unit is at the downstream end of three sizable irrigation ditches just before they spill back into the Yakima River. The wildlife area is allowed to withdraw as much water as necessary to maintain wetland and lake levels. In addition to wildlife habitat, the wetlands serve an additional purpose in filtering contaminants such as fertilizers and pesticides from the open-air irrigation water supply before it returns to the river. The abundant cattails in the wetlands are well-known water cleaners, efficiently up-taking heavy metals and other contaminants.

The Sunnyside Unit experiences the highest amount of recreation use of any unit of the wildlife area. Waterfowl and upland game bird hunting are popular activities on the unit, and pheasant hunting is particularly popular. Visitors must register on site when arriving, and report their activity before leaving. Registration provides usage and harvest data for future unit management.

WDFW has released pheasants on the site for many years, and it's become a family tradition for locals and non-locals alike to pheasant hunt on the unit every fall. Birding is another popular activity as many thousands of shorebirds and waterfowl stopover in the wetlands during spring



migration. American white pelicans, sandhill cranes, great egrets, white-faced ibis, black-necked stilts and American avocets are all regularly seen in the spring months.

The Snipes Game Reserve is an area established by <u>Chapter 220-411 WAC</u> in the center of the unit. Game reserves are areas where hunting and trapping for all wild animals and wild birds is prohibited, and no guns or dogs are allowed. Within the game reserve is Horseshoe Lake, an old river oxbow, an important resting place for wintering waterfowl. While some game reserves allow passive recreation, Snipes is closed to all public entry.

Six ponds and lakes that vary in size from 15 to 100 surface acres are spread throughout the unit. The 100-acre Giffin Lake, the largest one, was once popular with anglers but is no longer stocked with fish. There is not much access as the area has grown in with cattails and lily pads. Giffin is still a popular destination for birders, hikers, and some waterfowl hunters. The gate for the driveway to Giffin Lake is closed in the summer and open in the fall for hunting.

There are a couple of special circumstances on this unit, besides the Snipes closure. It is unlawful to possess shot (either in shotshells or as loose shot for muzzleloading) other than nontoxic shot for any purpose (WAC 220-414-040) on this unit. The Yakama Indian Reservation borders the south side of the Yakima River the entire length of the Sunnyside Unit. Yakama Nation fishing or hunting licenses are required to fish or duck hunt on the river, and sportsmen must follow the Yakama Nation regulations. WDFW licenses are also required.

Primary management objectives on the Sunnyside Unit

- Implement the 10-year plan for wetland management and restoration (Appendix D), which is focused on enhancements and also includes development of new wetlands on the wildlife area (1.B).
- Increase wetland function and value through vegetative restoration (1.C).
- Develop a protocol for management of nuisance beavers on the wildlife area by 2020 (1.G).
- Maintain and restore floodplain functions throughout the Yakima River Valley (1.I).
- Develop a prescribed burn plan for wetlands by 2020 (1.K).
- Develop regularly scheduled waterfowl surveys for use in assessing management actions (2.B).
- Identify and implement opportunities to enhance monarch butterfly, bumble bee, and other pollinator habitat by 2022 (2.C).
- Continue to improve recreational experience, user expectations, and support of the wildlife area by providing information on the web, at kiosks, in maps, and directional signage (3.B).
- Improve waterfowl hunting and recreational bird watching by restructuring and maintaining berms that form rice paddies by 2020 (3.F).
- Improve waterfowl hunting by working by increasing food resources and water supply in the Johnson Wetland by 2020 (3.G).
- Evaluate the function of the refuges and reserves for hunting and recreation opportunities by 2021 (3.H).
- Assess potential to improve lakes and stock with fish (trout and/or warm water) by 2024 (4.A).
- Manage and improve upland bird habitat and nesting cover (4.B).
- Maintain Disabled Hunter Access Program, including the Thornton ADA Special Permit for Elk hunting (limited entry special draw) (5.C).
- Ongoing, as agricultural leases come up for renewal or for new ones developed, evaluate agricultural leases per agency policy for multiple benefits, especially to improve waterfowl forage for hunting opportunities (7.B).



- Continue to work with BPA on operations and maintenance budget, securing adequate, sustainable funding, and reporting, and ensuring mitigation obligations are met (8.A).
- Remove derelict structures (10.A).
- Maintain and upgrade water control structures as necessary (10.B).



Sunnyside Unit Photo by Alan Bauer





Figure 7: Sunnyside Unit

Vance-Ferry Road Unit

Size	176 acres
Acquisition Date	1944 and 1993
Acquisition Funding	US Fish and Wildlife Service: <i>Pittman-Robertson Wildlife Restoration</i> <i>Program (PR)</i> US Army Corps of Engineers: <i>Snake River Mitigation</i> <i>Account</i> WA Dept. of Fish and Wildlife: <i>Wildlife Fund</i>
Purpose of Funding	Upland bird and small game habitat
Elevation Range	645 – 675 feet
Recreational Opportunities	Upland bird hunting and limited fishing
County	Yakima
Site Access	From Highway 22 to Ferry Road

Overview

The Vance-Ferry Road unit is made of two parcels totaling 176 acres, located one mile northeast of Mabton. It was purchased in 1944 for small game habitat. There is about three quarters of a mile of Yakima River frontage along the south shore of the river, but with a steep cut-bank there is little to offer for fishing access.

About 82 acres of the unit are leased agricultural fields in small grain and alfalfa hay rotation. Cattle were removed from this site and a buffer was established along the river that exists as a mix of annual and perennial grasses and broadleaf weeds. A small island of cottonwoods and shrubs separates the Vance acquisition from the old Ferry Road habitat plot. A single row of mature elm trees lies along the boundary. Two shrub plots were planted on the Vance Unit to break up the agricultural fields and provide more woody cover for wildlife.

This unit offers opportunities to hunt deer, waterfowl, Eurasian collared dove, mourning dove, quail, and small game. About two-thirds of the unit is within the Yakima River Waterfowl Closure area. In this special migratory bird closure it is unlawful to hunt migratory waterfowl, coot, and snipe within a quarter of a mile of the Yakima River. The unit has resident trout and anadromous spring Chinook salmon in the Yakima River.

Primary management objectives on the Vance-Ferry Road Unit

- Ongoing, as agricultural leases come up for renewal or for new ones developed, evaluate agricultural leases per agency policy for multiple benefits, especially to improve waterfowl forage for hunting opportunities (7.B).
- Explore the best option for long-term management of the Vance-Ferry Unit (10.D).







Byron Unit

Size	1,031 acres
Acquisition Date	1947 - 1994
Acquisition Funding	WA Dept. of Fish and Wildlife: Wildlife Fund
Purpose of Funding	Waterfowl habitat
Elevation Range	656-723 feet
Recreational Opportunities	Waterfowl and upland bird hunting, deer hunting, warm water fishing (limited), and birding
County	Yakima
Site Access	From SR-22 East from Mabton and Bus Road off of SR-22



Byron Unit Photo by Alan L. Bauer


The Byron unit encompasses 1,031 acres, and is located five miles east of Mabton and five miles south of Grandview in Yakima County. This unit is within the Yakima River Corridor and includes shrubsteppe habitat plus a wetland system that runs the entire length of the property.

The unit was established to provide waterfowl wintering and production habitat. The topography consists of rolling hills and potholes, and is very rocky with a large number of protruding quaternary flood basalt boulders and slabs. A large irrigation water ditch runs east to west through the unit, terminating at the Yakima River just past the east end of the unit. The ditch meanders through several large pothole ponds for approximately 2.5 miles, creating wetland habitat totaling over 400 surface acres. The wetlands often hold a variety of dabbling ducks during the winter months, giving way to nesting resident Canada geese and migrating shorebirds and ducks in spring.

The unit is also bisected by a large concrete pipe, the Mabton Siphon. The eastern side of the pipe is the Byron Game Reserve, which was established by <u>Chapter 220-411 WAC</u>. Game reserves are areas where hunting and trapping for all wild animals and wild birds is prohibited, and no guns or dogs are allowed. However, the game reserve on this unit is open year round to the public for walking, birding, photography, and other low-impact recreation. The reserve extends approximately one mile east of the pipe. It contains numerous pothole ponds, a 10-foot wide meandering irrigation ditch flowing west to east, and intact shrubsteppe flora with expanses of large, healthy big sagebrush and native forbs. A parking area to access the reserve is being developed at Byron Road at the eastern end of the unit.

The area west of the pipe is open to public access and hunting <u>only</u> during the general hunting season (go to <u>https://wdfw.wa.gov/hunting/regulations/summary-of-seasons</u> for a summary of hunting seasons). Under an agreement with the City of Grandview, it is closed to all use the rest of the year to provide refuge for nesting waterfowl and protect public health.

Part of the water supply for the Byron ponds comes from treated wastewater discharged from the adjacent City of Grandview Waste Water Treatment Plant (WWTP). Historically, non-hunting season shutdown of the Byron Unit was partially due to concerns about public exposure to treated wastewater (although deemed safe, there were still inherent risks). The WWTP has revamped their treatment processes and now have a purer standard of treated water exiting their system. The wildlife area management team is working with the City of Grandview to increase the amount of purified wastewater entering the Byron ponds throughout the year, maintaining uniform pond levels, and to possibly open up the west section of the unit for public access year-round once again.

Mosquitoes are very prevalent throughout the Byron area during the warmer months. The first detection of West Nile Virus in the area occurred at the Byron ponds in 2008. The Benton County Mosquito Control District conducts mosquito collection and testing around the Byron ponds. Ground spraying is routinely carried out and aerial spraying is initiated when a positive test for West Nile Virus occurs, as has happened every year since 2008 (except 2011). Maintaining a higher water level in the ponds year round, with the additional input from the City of Grandview WWTP, will enable the ponds to be stocked with fish and facilitate natural mosquito larvae control.

The unit is home to Columbia Basin shrubsteppe obligate plant species, including a healthy population of mature big sagebrush (*Artemisia tridentata*). Due to habitat degradation from historical cattle grazing in the area, cheat grass, Russian knapweed, and other invasive species have



a solid hold in the area. Canada geese thrive on wetlands due to abundant forage on newly sprouted cheat grass shoots in the spring. The sandy, rocky, alkaline soils make native plant restoration challenging on this unit. Alkaline tolerant species such as saltgrass *(Distichlis spp.)* have successfully been used to recolonize disturbed areas at the Byron Unit.

The Byron Unit supports a variety of upland and shrubsteppe wildlife species including abundant California quail and mourning doves, northern and loggerhead shrikes, sagebrush sparrows, mule deer, yellow-bellied marmots, and numerous reptiles. Byron is particularly well-known for its dense population of western rattlesnakes, which, according to anecdotal evidence, are of a unique greenish coloration. While rattlesnakes have been nearly extirpated locally due to intensive agriculture and housing development, Byron provides a protective refugium for this unique rattlesnake population.

In addition to hunting, birding is a popular activity at Byron. During the winter months, many bird species utilize the wetlands and islands of trees in the unit for cover and food sources. Red-winged blackbirds gather in large flocks in the cattails around the ponds and wintering flocks of American robins can be found throughout the unit feeding on Russian olive fruits. Waterfowl including mallards, northern pintails, green-winged teal, common goldeneye, snow geese, Canada geese, and tundra swans can all be found on the ponds during colder months. Lesser scaup arrive in large flocks during spring migration along with blue-winged teal, shorebirds, and occasional sandhill cranes. Mallards, green-winged teal and Canada geese stay in the area to breed in the springtime. Many birds of prey are found in the Byron Unit as well, including northern harriers, red-tailed hawks, sharp-shinned hawks, Cooper's hawks, American kestrels, and the occasional prairie falcon.

On this unit, it is unlawful to possess shot (either in shotshells or as loose shot for muzzleloading) other than nontoxic shot for any purpose (<u>WAC 220-414-040</u>).

Primary management objectives on the Byron Unit

- Update and implement the 10-year plan for wetland management and restoration (Appendix D), which is focused on enhancements and also includes development of new wetlands on the wildlife area (1.B).
- Increase wetland function and value through vegetative restoration (1.C).
- Coordinate with the City of Grandview to ensure consistent water supply to Byron ponds for wildlife and recreation (1.F).
- Develop a prescribed burn plan for wetlands by 2020 (1.K).
- Develop regularly scheduled waterfowl surveys for use in assessing management actions (2.B).
- Identify and implement opportunities to enhance monarch butterfly, bumble bee, and other pollinator habitat by 2022.
- Evaluate the function of the refuges and reserves for hunting and recreation opportunities by 2021 (3.H).
- Assess potential to improve lakes and stock with fish (trout and/or warm water) by 2024 (4.A).
- Open the west side of the Byron Unit year round (5.D).
- Ongoing, as agricultural leases come up for renewal or for new ones developed, evaluate agricultural leases per agency policy for multiple benefits, especially to improve waterfowl forage for hunting opportunities (7.B).
- Continue to work with BPA on operations and maintenance budget, securing adequate, sustainable funding, and reporting, and ensuring mitigation obligations are met (8.B).





Washington Department of Fish and Wildlife

Whitstran Unit

Size	31 acres
Acquisition Date	1993
Acquisition Funding	US Army Corps of Engineers: <i>Snake River Mitigation Account</i> Private Grantor: <i>Private Donations</i>
Purpose of Funding	Water access
Elevation Range	573 – 629 feet
Recreational Opportunities	Warm water fishing, and limited waterfowl and upland bird hunting, deer and elk hunting
County	Benton
Site Access	From I-82, exit 80 to Bunn Road



Fishing the Yakima River at the Whitstran Unit Photo by Alan L. Bauer



The Whitstran Unit is made of two parcels comprising 31 acres, located two miles east of Prosser along the Yakima River in the lower Yakima watershed. The northern parcel is an irrigation canal. It is primarily shrubsteppe and riparian habitat.

Whitstran is home to a healthy population of well-established big sagebrush (*Artemisia tridentata*), a plant that is in decline regionally due to fires and development. The unit also hosts a healthy riparian zone of cottonwood, willow species, Wood's rose, and other riparian obligates. The unit offers birders, quail hunters, duck hunters, and anglers a unique place to recreate in a small, but relatively intact shrubsteppe ecosystem within close reach of Interstate 82 and the town of Prosser. Fishing for fall Chinook salmon and waterfowl hunting are the primary activities here.

The only access to the Whitstran Unit is via an irrigation canal access road. The access road is narrow, and visitors used to park on the road, providing challenges for neighbors and larger vehicles. Due to the large volume of salmon anglers in the fall, a second parking lot was recently added to reduce road parking.

Primary management objectives on the Whitstran Unit

- Maintain and restore floodplain functions throughout the Yakima River Valley (1.I).
- Research records on this and other units to determine legal boundaries and access by 2024 (5.B).



Figure 10: Whitstran Unit



Thornton Unit

Size	2,080 acres
Acquisition Date	1994
Acquisition Funding	WA Dept. of Fish and Wildlife: Wildlife Fund
Purpose of Funding	Small game habitat
Elevation Range	1,471 – 2,446 feet
Recreational Opportunities	Upland bird and deer and elk hunting, birding, horseback riding and hiking
County	Benton
Site Access	From I-82, exit 90 to Rothrock Road

Overview

The Thornton Unit comprises 2,080 acres along the southern slopes of Rattlesnake Mountain, about eight miles northeast of Prosser in Benton County. The unit provides habitat for many species including mule deer, upland birds, winter concentrations of Rocky Mountain elk (the Rattlesnake Hills herd), and whitetailed jackrabbit. Ferruginous hawks have been known to nest in proximity to this unit. A mitigation project is being planned which will install one or two platforms on raised poles on the Thornton and Rattlesnake Slope units. Continued monitoring of historic nest sites is also needed, and protection of occupied sites from March-July from disturbance would be needed.

The unit supports ridges of excellent shrubsteppe, intermixed with former wheat fields. Restoration of these fields was completed in 1999, and they now support sage brush sparrows, sage thrashers, and Gray



Thornton Unit Photo by Justin Haug, WDFW



(Hungarian) partridge and quail. Other shrubsteppe-obligate species may also benefit from shrubsteppe habitat.

Snipes Creek, an intermittent spring fed stream flowing until mid-summer, bisects the unit. Snipes spring in the center of this unit provides water nearly year round and supports important riparian habitat and cover for wildlife. Other smaller seeps exist in the canyons creating a diversity of habitat and wildflower blooms.

With limited huntable public land on Rattlesnake Mountain, the Thornton Unit provides one of the best places to hunt the Rattlesnake Mountain herd. More elk are harvested here than any other unit of the Sunnyside Wildlife Area. The unit is also a popular hunting area for mule deer, partridge, and quail. Lucky hunters might also find chukar, but populations have declined in recent years.

Abundant wildflowers and quality shrubsteppe attract hikers and naturalists in the spring. The unit is a reliable place to see the uncommon and local sagebrush sparrow singing from sagebrush tips from March through May. While small and not particularly flashy, this species requires large patches of undisturbed sagebrush and its presence indicates the quality of the habitat here. Limited disabled hunter drive-in access is available on this site by permit only. Otherwise the unit is walk-in only. It has two parking lots, one along Rothrock Road on the west side and one along Case Road on the east side.

Primary management objectives on the Thornton Unit

- Install nest platforms for ferruginous hawks to increase nesting opportunities by 2021 (2.D).
- Maintain Disabled Hunter Access Program, including the Thornton ADA Special Permit for Elk hunting (limited entry special draw) (5.C).
- Ongoing, as agricultural leases come up for renewal or for new ones developed, evaluate agricultural leases per agency policy for multiple benefits, especially to improve waterfowl forage for hunting opportunities (7.B).
- Continue to work with BPA on operations and maintenance budget, securing adequate, sustainable funding, and reporting, and ensuring mitigation obligations are met (8.B).



Figure 11: Thornton Unit



Rattlesnake Slope Unit

Size	3,661 acres
Acquisition Date	1973
Acquisition Funding	US Congress Programs: Transfer of Land
Purpose of Funding	Small game habitat
Elevation Range	448 – 2,069 feet
Recreational Opportunities	Chukar, deer and elk hunting, hiking, horseback riding, target shooting in designated range, mountain biking
County	Benton
Site Access	From Highway 225, north of Benton City



Balsamroot on Rattlesnake Slope Photo by Alan L. Bauer



The Rattlesnake Slope Unit comprises 3,661 acres of ridge and canyon landscape, located just five miles north of Benton City. High points along the eastern ridge of Rattlesnake Mountain at 2000 feet provide for commanding views of the lower Yakima River valley, vineyards, and a bird's eye perspective of prominent landmarks including Red, Candy, and Badger Mountains in linear formation. While Rattlesnake Ridge is the prominent feature along the south boundary of the property, gently sloping grassland terrain flows down to the east and north, leveling out to 400 feet elevation toward Horn Rapids.

The unit provides habitat for many species including white-tailed jackrabbit, winter concentrations of Rocky Mountain elk, mule deer, chukar, and black-tailed jackrabbits. Ferruginous hawks have been known to nest in proximity to this unit. A mitigation project is being planned which will install one or two platforms on raised poles on the Thornton and Rattlesnake Slope units. Continued monitoring of historic nest sites is also needed, and protection of occupied sites from March-July from disturbance would be needed.

The east side of the property, near to Hwy 225, is highly disturbed and dominated by invasive cheat grass and knapweed. The interior portion of the site remains quality bunchgrass with abundant and diverse wildflowers that are noticeably visible in the early spring. However, a large fire in 2018 burned 90% of the unit, and restoration and natural regrowth have yet to occur on the site.

Access to the site is from the parking lot on the west side of Hwy 225 about 7.5 miles north of the I-82 Benton City Exit roundabout, or 4 miles south of Hwy 240. Visitors must register on site when arriving, and report their activity before leaving. Registration provides usage and harvest data for future management of the unit.

No motorized transport is authorized on the unit. The area is popular for recreation, including horseback riding, hiking, and mountain biking. Several unofficial and unmaintained trails have been made by users throughout the property. Because the trails were not designed, some of them cause damage to the habitat because of improper placement and lack of erosion and drainage control. WDFW will work with the advisory committee and user groups to plan for sustainable trails. In addition to trail use, hunting is also popular, with chukar, deer, and elk the most likely species to be hunted.

A shooting range was established on land leased to Benton County and run by the Tri Cities Shooting Association. The shooting range is staffed and well-maintained as a multiple use range open to the public with a small fee and set hours. This is the only place where target shooting is allowed on the wildlife area.

A notable adjacent property is the Arid Lands Ecology Reserve (ALE). It is managed for the US Department of Energy by the Pacific Northwest National Laboratory. No general public access is currently allowed on the ALE, and use is limited to approved research and education. The nearby Saddle Mountain National Wildlife Reserve is part of the Hanford Reach National Monument with some use restrictions. For more details, go to: <u>https://www.fws.gov/refuge/Hanford Reach/</u>.



Primary management objectives on the Rattlesnake Slope Unit

- Install nest platforms for ferruginous hawks to increase nesting opportunities by 2021 (2.D).
- Evaluate habitat loss from recent fires on Rattlesnake Slope and develop restoration strategy by 2020 (1.J).
- Continue to improve recreational experience, user expectations, and support of the wildlife area by providing information online, at kiosks, in maps, and directional signage (3.B).
- Work with the advisory committee and user groups to assess user-built trails and plan for sustainable trail system 2022 (3.E).
- Assess potential boundary issues on Rattlesnake Slope by 2020 and others as necessary (7.A).
- Continue to work with BPA on operations and maintenance budget, securing adequate, sustainable funding, and reporting, and ensuring mitigation obligations are met (8.B).







Benton City Unit

Size	25 acres
Acquisition Date	1968 - 1994
Acquisition Funding	National Park Service: <i>Land and Water Conservation Fund</i> US Army Corps of Engineers: <i>Snake River Mitigation Account</i> WA Recreation and Conservation Office: <i>State Bond Account</i>
Purpose of Funding	Yakima River access
Elevation Range	452 – 486 feet
Recreational Opportunities	Fishing, birding, and walking
County	Benton
Site Access	From Highway 225 to 14 th Street



Benton City Unit Photo by Alan L. Bauer



This unit consists of two parcels along the Yakima River near Benton City. They provide fishing and water access. One parcel contains about 16 acres of Yakima riverfront property directly west of the Benton City limits. A second nine-acre parcel, named River Park, is in town off of 1st Street. The properties are owned by WDFW and managed by Benton City under a long-term agreement.

Columbia clubtail (a state species of concern), and white-belted ringtail dragonfly may occur here. Bull trout, steelhead, both federally-threatened species in this area, and Pacific lamprey and mountain sucker utilize the Yakima River.

This unit offers a shoreline fishing access site on the Yakima River and good riparian habitat. The city manages a developed boat ramp and picnic area at River Park, and hiking trails on the west parcel. Anglers can shoreline fish for salmon, bass, carp, suckerfish, and walleye, or launch boats for fisheries in the Yakima River. While waterfowl hunting is not allowed in city limits, hunters can launch craft from the ramp at River Park and hunt non-incorporated waters up and down stream.

The trails on the west parcel provide for short walks and bird watching opportunities with neotropical migrants passing through or breeding during the spring and summer in the wooded riparian areas.

Primary management objectives on the Benton City Unit

Since the Benton City Unit is managed by the city, the only specific objective that WDFW has, along with the Whitstran, I-82 Ponds, and Sunnyside Unit is to:

- Maintain and restore floodplain functions throughout the Yakima River Valley (1.I).
- Maintain agreement with Benton City to manage the unit for recreation (7.D).



Benton City Unit trail Photo by Alan L. Bauer





Hope Valley Unit

Size	549 acres (deeded, leased, and managed)
Acquisition Date	1965 - 1995
Acquisition Funding	WA Recreation and Conservation Office: WA Wildlife and Recreation Program
Purpose of Funding	Small game habitat and water access
Elevation Range	743 – 824 feet
Recreational Opportunities	Waterfowl, pheasant, and upland bird hunting, fishing, and bird watching
County	Franklin
Site Access	For southern entrance: From Hwy 395, take Eltopia West Road and head west 4.5 miles to Hooper Road. Turn right and head north for 2.5 miles to the parking lot. For Clark Pond: From I-82, take the Glade Road exit. Head north on Glade for 17 miles. Turn right and go east on Ironwood 3 miles.



Clark Pond, Hope Valley Unit Photo by Alan L. Bauer



Three small units, Hope Valley, Clark Pond, and Nipper have been combined to be called the Hope Valley Unit. About 213 acres are owned by WDFW, primarily in the south parcel, 60 acres are leased from the DNR, and 274 acres are BOR land managed by WDFW. The part of the unit that contains Clark Pond consists of acreage leased from DNR and owned by BOR. The unit is located between the Glade North Rd and I-395, between Mesa and Eltopia in Franklin County.

The Hope Valley Unit combines quality shrubsteppe uplands with riparian and wetland water features for a diversity of wildlife habitat and opportunities packed into a small unit. Irrigation water flows from the north into Clark Pond and south through a chain of wetland pools leading into the Eltopia Branch Canal that bisects the southern parcel. Waterfowl hunting, pheasant hunting at the release site, and some fishing comprise most of the current recreation on the unit.

Clark Pond off Ironwood Road contains various warmwater fish species for year-round fishing; however, water quality is poor in the summer months. A primitive boat launch at the parking area provides access for hunters and anglers with small craft. During winter months, puddle ducks and a few divers can cover Clark Pond or be completely absent, providing erratic hunting opportunities. Hunters and bird watchers can scope the entire pond from the roadside and parking area to quickly assess waterfowl numbers. The cattail-fringed wetlands to the south usually hold small numbers of puddle duck as does the irrigation canal on the south parcel. The south parking lot is also part of the Eastside Pheasant Enhancement Program's Release sites. Staff release farm-raised roosters for the youth pheasant hunt and into the general season.

Most of the soil in this unit is sandy and alkaline, and generally undesirable for farming, though there are about 30 acres in an agriculture lease. Dryland grasses and shrubs have been planted to stabilize the site and provide habitat for wildlife. The once-common and now imperiled monarch butterfly benefits from the presence of milkweed on the unit, a main food source for its caterpillar. Beavers have caused some concern in the irrigation ditches, and in the past have been trapped and relocated. As part of this management plan, a protocol for management of nuisance beavers will be developed.

The southern part of the unit (the original Hope Valley) is the other commonly used area. The Eltopia Branch Canal, a main irrigation canal bisects this part of the unit. The parcel west of the canal is dryland, with no irrigation allotment. Three shallow draws receive some sub-irrigation water from the canal, which supports emergent vegetation and Russian olive trees. The parcel on the north and east of the canal is a mix of shrubsteppe habitat and cropland. There is some good upland habitat for a variety of species such as California quail and pheasant. Like many units, controlling weeds is an on-going effort here. Visitors must register on site when arriving, and report their activity before leaving. Registration provides usage and harvest data for future management of the unit.

Access to the unit is via a north parking lot along Ironwood Road and a south lot at the bend in Hooper Road. Access between Clark Pond parking area and the south parcels is possible along the 2-track west of the small irrigation ditch heading south, thanks to a public access easement. The land on both sides of this track is private and users must stay on the track and respect private property. No off-road driving is permitted on the unit. Currently, the irrigation canal prevents



access to and from the south parking lot to the rest of the unit. Funds have been sought to build a footbridge over the canal in order to increase accessibility to the site.

Primary management objectives on the Hope Valley Unit

- Increase wetland function and value through vegetative restoration (1.C).
- Develop a protocol for management of nuisance beavers on the wildlife area by 2020 (1.G).
- Develop regularly scheduled waterfowl surveys for use in assessing management actions (2.B).
- Identify and implement opportunities to enhance monarch butterfly, bumble bee, and other pollinator habitat by 2022 (2.C).
- Install footbridge across irrigation canal at Hope Valley Unit by 2022 (3.D).
- Assess potential to improve lakes and stock with fish (trout and/or warmwater) by 2024 (4.A).
- Manage and improve upland bird habitat and nesting cover (4.B).
- Ongoing, as agricultural leases come up for renewal or for new ones developed, evaluate agricultural leases per agency policy for multiple benefits, especially to improve waterfowl forage for hunting opportunities (7.B).





Mesa Lake Unit

Size	695 acres
Acquisition Date	1965-2011
Acquisition Funding	National Park Service: Land and Water Conservation Fund US Fish and Wildlife Service: Pittman-Robertson Wildlife Restoration Program (PR) WA Recreation and Conservation Office: State Bond Account WA Dept. of Fish and Wildlife: Wildlife Fund; State Migratory Waterfowl Fund
Purpose of Funding	Waterfowl habitat
Elevation Range	647 – 867 feet
Recreational Opportunities	Waterfowl and upland bird hunting, fishing, birdwatching
County	Franklin
Site Access	From Highway 395 to Langford Road

Overview

The Mesa Lake Unit contains the 50-acre namesake lake, part of a much larger irrigation system and network of connected "lakes", canals, and wetlands found through central and north Franklin County. The unit is located in directly adjacent to (and partially in) the city limits of Mesa. While mostly owned by WDFW, the Mesa Lake Unit contains 93 acres of Bureau of Reclamation (BOR) land. Though WDFW started managing some of the BOR lands in 1965, the majority of the unit was purchased by WDFW in



Mesa Lake Unit Photo by Alan L. Bauer

2010 from a private hunting club. It consists of shrubsteppe, upland grasslands, wetlands, and a lake. About 30 acres are in an agriculture lease.

There are approximately 15-20 acres of managed wetlands on the plateau above and to the north of the lake. These wetlands range in size from two to seven acres and are managed to provide shallow open water and emergent wetland vegetation. Water levels are regulated through flashboard risers to raise and lower water levels and promote vegetation growth and cover. Periodically wetlands are drained and mowed or sprayed, or both, to control both beneficial and noxious wetland vegetation.



These efforts provide valuable vegetation cover and food sources for primarily waterfowl, but other wetland dependent bird species also benefit.

In addition to several managed wetlands and upland habitats, the unit provides for a diversity of wildlife and recreation. Uplands consists of shrubsteppe, grasslands, and a 30-acre agriculture lease. The unit is a great site for waterfowl, quail and dove hunting, provides a warmwater fishery, and wildlife viewing year round.

Historically, the lake was treated and stocked with trout, but has not been stocked for many years. It is now managed primarily as a warmwater fishery, relying on naturally reproducing species such as largemouth bass, bluegill, and yellow perch. Additionally, species such as walleye and lake whitefish are found in the lake as they move in via the irrigation canals. The ponds don't hold many fish as they are drained and filled seasonally for waterfowl management. The smaller wetlands are managed for waterfowl with several blinds established on site. The west side of the lake has a parking lot and boat launch for fishers and hunters.

In late winter during migration, several thousand geese of mixed species may roost in the area. The managed shallow wetlands attract migrant shorebirds during spring and fall, and support breeding stilts, waterfowl, and a variety of songbirds during the summer. Over 110 bird species have been recorded on the unit in the past couple years, with many more likely still to be found. The once-common and now imperiled monarch butterfly benefits from the presence of milkweed on the unit, a main food source for its caterpillar. Ferruginous hawks have been known to nest close to this unit.

Visitors must register on site when arriving and report their activity before leaving. Registration provides usage and harvest data for future management of the unit. All users must sign in at the kiosk at one of two parking lots. One lot is located at the southwest end of the lake where there is also a boat ramp, and the other parking lot is just the south of the managed wetlands.

An issue not unique to Mesa, beavers are found throughout west Franklin County's irrigated landscape, and on many of the other WDFW units such as Bailie and Windmill. They have proved challenging at times regarding water management as they frequently dam up pipes and water control structures. As a result, a nuisance beaver management protocol is being developed that will help guide beaver management activities in the future.

Primary management objectives on the Mesa Lake Unit

- Update and implement the 10-year plan for wetland management and restoration (Appendix D), which is focused on enhancements and also includes development of new wetlands on the wildlife area (1.B).
- Develop a protocol for management of nuisance beavers on the wildlife area by 2020 (1.G).
- Develop a prescribed burn plan for wetlands by 2020 (1.K).
- Develop regularly scheduled waterfowl surveys for use in assessing management actions (2.B).
- Identify and implement opportunities to enhance monarch butterfly, bumble bee, and other pollinator habitat by 2022 (2.C).
- Continue to improve recreational experience, user expectations, and support of the wildlife area by providing information online, at kiosks, in maps, and directional signage (3.B).
- Assess potential to improve lakes and stock with fish (trout and/or warmwater) by 2024 (4.A).
- Manage and improve upland bird habitat and nesting cover (4.B).



- Ongoing, as agricultural leases come up for renewal or for new ones developed, evaluate agricultural leases per agency policy for multiple benefits, especially to improve waterfowl forage for hunting opportunities (7.B).
- Remove derelict structures (10.A).
- Maintain and upgrade water control structures as necessary (10.B).



Figure 15: Mesa Lake Unit



Esquatzel Coulee Unit

Size	1,847 acres
Transaction Date	1969 - 1984
Acquisition Funding	No deeded land
Purpose of Funding	Waterfowl and small game habitat
Elevation Range	681 – 984 feet
Recreational Opportunities	Waterfowl and upland bird hunting, deer hunting, fishing, bird watching
County	Franklin
Site Access	From Highway 395, to Highway 17

Overview

The Esquatzel Coulee Unit is 1,785 acres of upland canyons, grassland, and wetland, including a narrow lake, located less than half of a mile north of the town of Mesa. This unit is owned by the Bureau of Reclamation (BOR) and managed by WDFW in cooperation with BOR and the South Columbia Basin Irrigation District. It is roughly shaped like the letter "V" that has tipped to the right with its open end tilted to the northeast. Approximately 1,200 acres lie in the bottom of the Esquatzel Coulee along a northeast axis bisected by



Esquatzel Coulee Unit Photo by Alan L. Bauer

the Burlington Northern Railroad and the remaining contiguous acreage is situated on a northerly axis that is the terminus of a long and narrow linear valley through mostly arid rangelands. The lake is a flooded canyon bottom composed of irrigation and seepage water year round and attracts waterfowl.

Columnar basalt and basalt outcrops occur through the unit, especially on the south (north facing) wall of the coulee. This rugged south wall also supports native bunch grasses and shrubsteppe habitats. Pockets of shrubsteppe occur through the remainder of the unit giving way to mostly grass dominated rangelands. Pheasants Forever, through a permit with the BOR, maintains approximately 20-40 acres of irrigated habitat and crops in small plots for wildlife.



The portion along the railroad easement supports approximately three miles of the Esquatzel Wasteway, which is operated by the South Columbia Basin Irrigation District through an agreement



Northern harrier Photo by Alan L. Bauer

with BOR. The wasteway occurs in topographically low areas and may follow a historic stream channel, but it frequently follows channels that have been engineered for water conveyance. In some adjacent areas, wetland habitats have formed. The waters and flow of the wasteway are irrigation driven and not dependent on precipitation or natural flow.

The unit is managed for public fishing and hunting. There is potential for birdwatching and hiking here as well. There is a public parking area located at the southwest end, where only walk-in access is permitted. Wildlife that utilize the unit include mule deer and a variety of waterfowl and birds. SGCN species including ferruginous hawk, a (state threatened species), and Washington ground squirrels (state candidate species) may also occur here, but there have not been any recent observations.

Primary management objectives on the Esquatzel Coulee Unit

- Identify and implement opportunities to enhance monarch butterfly, bumble bee, and other pollinator habitat by 2022 (2.C).
- Continue to improve recreational experience, user expectations, and support of the wildlife area by providing information online, at kiosks, in maps, and directional signage (3.B).
- Manage and improve upland bird habitat and nesting cover (2.B).
- Research areas where possible access or access agreements need to be updated by 2024 (5.B).



Figure 16: Esquatzel Coulee Unit



Windmill Ranch Unit

Size	2,139 acres
Acquisition Date	1960 - 2001
Acquisition Funding	US Army Corps of Engineers: Snake River Mitigation Account
Purpose of Funding	Small game habitat and lake access
Elevation Range	773 – 924 feet
Recreational Opportunities	Upland bird and waterfowl hunting, deer and elk hunting, fishing and ice fishing, bird watching
County	Franklin
Site Access	From Highway 17 to Colonial Road



Windmill Ranch Unit Photo by Alan L. Bauer

Overview

The Windmill Ranch Unit is about half deeded land, and the other half a combination of land leased from DNR and owned by the BOR. The unit consists of four separate parcels totaling over 2,000 acres located about three miles northwest of Mesa in Franklin County. This former irrigated ranch provides a variety of diverse habitats at the southern end of the channeled scablands. A large lake runs across the northern boundary. Sandhill cranes are a common site in the fields in the spring, and heavy concentrations of waterfowl are common in the hunting season.

Approximately 432 acres are in an agriculture lease and row crops are grown for the benefit of wintering waterfowl and migrating sandhill cranes. The rest of the unit is a mosaic of several habitat types, including shrubsteppe, grasslands, wetlands, and riparian. A history of grazing and fire has removed much of the shrub component from the shrubsteppe areas, but replanting efforts are contributing to successful restoration in some areas.



The Columbia Basin Irrigation Project, which is the source of all water on this site, has created a variety of wetland habitats. Wetlands on this unit are primarily formed by irrigation seepage and wastewater, and return flows, and are scattered throughout the unit. Some are intensively managed for moist soil conditions, waterfowl production, and winter food while others are left to function naturally.

Noxious wetland vegetation, including purple loosestrife and Russian olive, are prevalent in many wetlands, and some upland areas are plagued with Canada thistle and knapweed. Cattails are dense in many wetlands and beavers are common. The north part of the unit has some of the best milkweed plants for monarch butterfly.

The 50-acre Powerline Lake is located on this unit. Visitors must park and walk in about 1.5 miles to access to the lake. This is a deep (>60 feet) lake that is spring fed and maintains a fairly stable water level but occasionally discharges excess water into wetland habitat to the west. There is minimal irrigated agriculture to the southeast of the lake, with grassland and shrubsteppe as the majority of the surrounding habitat.

Visitors must register on site when arriving, and report their activity before leaving. Registration provides usage and harvest data for future management of the unit.

The lots are strictly limited to eight vehicles to control overuse during popular hunting and fishing seasons. Registration at this site ensures a



Sign access at Powerline Lake, Windmill Ranch Photo by Alan L. Bauer

quality experience for users and provides usage and harvest data for future management of the unit. This unit provides opportunities for bird watching, upland bird and waterfowl hunting, and deer and elk hunting. Fishing for warmwater species includes bass, bluegill, crappie, and occasionally triploid (sterile and non-reproductive) trout are stocked. Waterfowl hunting is common throughout the unit at suitable open water wetlands and, if the winter is cold enough, ice fishing is done at Powerline Lake.

Primary management objectives on the Windmill Ranch Unit

- Update and implement the 10-year plan for wetland management and restoration (Appendix D), which is focused on enhancements and also includes development of new wetlands on the wildlife area (1.B).
- Increase wetland function and value through vegetative restoration (1.C).
- Develop a protocol for management of nuisance beavers on the wildlife area by 2020 (1.G).
- Develop a prescribed burn plan for wetlands by 2020 (1.K).
- Develop regularly scheduled waterfowl surveys for use in assessing management actions (2.B).



- Identify and implement opportunities to enhance monarch butterfly, bumble bee, and other pollinator habitat by 2022 (2.C).
- Continue to improve recreational experience, user expectations, and support of the wildlife area by providing information online, at kiosks, in maps, and directional signage (4.B).
- Assess potential to improve lakes and stock with fish (trout and/or warmwater) by 2024 (4.A).
- Improve fishing opportunities by managing aquatic weeds in stocked lakes (4.D).
- Ongoing, as agricultural leases come up for renewal or for new ones developed, evaluate agricultural leases per agency policy for multiple benefits, especially to improve waterfowl forage for hunting opportunities (7.B).
- Maintain and upgrade water control structures as necessary (10.B).



Figure 17: Windmill Ranch Unit

Bailie Unit

Size	3,897 acres (hunting easement)
Transaction Date	1986
Acquisition Funding	No deeded land
Purpose of Funding	Small game habitat, hunting easement
Elevation Range	632 – 968 feet
Recreational Opportunities	Waterfowl and upland bird hunting, deer and elk hunting,
County	Franklin
Site Access	From Highway 17 to Sagehill Road

Overview

The Bailie Unit is 3,897 acres of the slightly larger (4,200 acre) Bailie Memorial Youth Ranch (Ranch). The Ranch provides a structured setting for pre-adoption youth 8-14 years old. The Ranch is privately owned and managed by a foundation. In 1986, WDFW purchased a perpetual hunting easement to provide limited public access.

As a result of the Columbia Basin Project, wetlands on this unit, just like those on the Windmill and Mesa, are part of a much larger irrigation system and network of connected "lakes", canals, and wetlands. Uplands are a mix of grasslands and degraded shrubsteppe. Approximately 300-400 acres are managed by the Ranch as irrigated croplands.

The Woodhouse's toad, an SGCN species, breeds on this unit. Other species that could utilize this unit include great blue heron, cinnamon teal, ducks, and waterfowl.



Bailie Unit Photo by Alan L. Bauer





Bailie Youth Ranch Sign Photo by Alan Bauer

Over the last decade, the Ranch has used the unit for cattle grazing that has impacted native habitats. WDFW works closely with the Ranch board of trustees to maintain and restore grasslands and shrubsteppe and provides direct management of wetlands through water level management and wetland vegetation mowing and spraying.

At this unit, hunting is the only activity allowed under the hunting easement, and you must register to hunt. Hunting is allowed only on Saturdays, Sundays, Wednesdays, and legal holidays. Waterfowl, deer, elk, small game, and upland bird hunting are available. Other recreational activities such as fishing may be available by contacting the Bailie Youth Ranch.

Primary management objectives on the Bailie Unit

- Update and implement the 10-year plan for wetland management and restoration (Appendix D), which is focused on enhancements and also includes development of new wetlands on the wildlife area (1.B).
- Increase wetland function and value through vegetative restoration (1.C).
- Develop a protocol for management of nuisance beavers on the wildlife area by 2020 (1.G).
- Develop regularly scheduled waterfowl surveys for use in assessing management actions (2.B).
- Identify and implement opportunities to enhance monarch butterfly, bumble bee, and other pollinator habitat by 2022 (2.C).
- Manage and improve upland bird habitat and nesting cover (4.B).





Thompson Seeps Unit

1,459 acres
1969 - 1984
No deeded land
Small game and waterfowl habitat
749 – 1,000 feet
Pheasant and quail hunting, bird watching
Franklin
From SR 17 to Mt. Vista Road



Thompson Seeps Unit Photo by Alan L. Bauer



The Thompson Seeps Unit, previously known as WB-10 Wasteway Unit, is a 1,459-acre unit, located about eight miles northwest of Basin City in Franklin County. The unit is narrow, about five to six miles long and about a half mile wide located at the bottom of a natural coulee. It is owned by the BOR and is managed for public fishing and hunting access by WDFW.

The large coulee carries irrigation return water to the Columbia River through a series of engineered canals and naturally occurring topographic low areas. The south side is flanked by high quality shrubsteppe while the north is adjacent to irrigated crop lands. In the past, more irrigation return water flowed through the unit, providing a variety of shallow and deeper water habitats for fish and waterfowl, but upgrades to irrigation pumps and improved water diversions have decreased the amount of water in this unit.

This unit is the focus of a wetland restoration project with Ducks Unlimited as well as the site of proposed project to improve native habitat connectivity through the restoration of riparian and shrub habitat for mule deer. WDFW is working with Ducks Unlimited, the Bureau of Reclamation, and other partners to evaluate the potential site improvements, and are in the preliminary stages of planning. It will take coordination, time, and funding to develop a plan and begin implementing these projects.

WDFW management of this unit consists mostly of weed control and some road maintenance where flooding occurs because of beaver activity.

Public recreation includes pheasant and quail hunting, as well as bird watching.

Primary management objectives on the Thompson Seeps Unit

- Update and implement the 10-year plan for wetland management and restoration, which is focused on enhancements and also includes development of new wetlands on the wildlife area (1.B).
- Develop a prescribed burn plan for wetlands by 2020 (1.K).
- Develop wetland habitat in Thompson Seeps Unit by 2022 (1.L).
- Develop regularly scheduled waterfowl surveys for use in assessing management actions (2.B).
- Assess potential to improve lakes and stock with fish (trout and/or warm water) by 2024 (4.A).




Figure 19: Thompson Seeps Unit

Land Ownership and Management, Funding, and Agreements

Acquisition history, funding, and purpose

WDFW and The Sunnyside-Snake River Wildlife Area staff work closely with many organizations such as conservation groups, local governments, irrigation districts, and user groups to identify priorities for purchase, management, and restoration of lands. Many of the units of the wildlife area are near other public and private conservation lands and holdings and coordination about planning and restoration and recovery projects provides efficient and effective habitat and species management over a larger landscape.

The fundamental reason for acquiring lands within the Sunnyside-Snake River Wildlife Area was to protect and enhance native habitats and to provide public recreation. Some of the lands also serve to help meet mitigation goals for the Bonneville Power Administration (BPA) and U.S. Army Corps of Engineers (Corps) construction projects. Although the mitigation details differ between these two entities, they both address habitat losses for upland wildlife, waterfowl, big game, and a variety of non-game species.

In the fall of 2001, and as a result of mitigation for the construction of four dams on the Snake River, the Lower Snake River Fish & Wildlife Compensation Program was designed to compensate for habitat and wildlife losses that were incurred due to the construction of those dams. The Corps was the project proponent for Snake River dam construction and funded mitigation efforts to replace habitat losses. Part of that program included the acquisition of 24,000 acres within southeast Washington. Some of those lands, specifically those in Franklin, Benton, and Yakima counties, were combined with existing wildlife areas.

The 17 units of the wildlife area are managed by WDFW under a variety of instruments, including deeds, agreements, leases, and easements.

Acres	Instrument
12,598	Deeded acres (owned by WDFW)
4,198	Owned by BOR and managed by WDFW under an agreement
320	Owned by USFWS and managed by WDFW under agreement
175	Leased (primarily from DNR)
2 000	Easements – primarily one large hunting easement and some
3,303	fishing easements
21,120	TOTAL

Table 2: Wildlife area land instruments



Agreements

Deeded land

Almost 60% of the wildlife area is owned by WDFW. The primary funders for the deeded land include: WA Recreation and Conservation Office (*State Bond Account*); WA Dept. of Fish and Wildlife (*Wildlife Fund*); National Park Service (*Land and Water Conservation Fund*); US Army Corps of Engineers (*Snake River Mitigation Account*); and US Fish and Wildlife Service (*Pittman-Robertson Wildlife Restoration Program*).

Agreements

Under agreement with the Bureau of Reclamation, WDFW manages about 4,198 acres in multiple units, primarily on the eastern side of the wildlife area. Additionally, 320 acres of the Byron Unit are managed under an agreement with the US Fish and Wildlife Service for habitat protection. Under a long-term agreement, Benton City manages the Benton City Unit, owned by WDFW, for recreation, and takes care of maintenance and operations.

Easements

Easements are a right, held by an entity other than the underlying fee title landowner, to cross or otherwise use a portion of the land for a specified purpose. WDFW holds easements for public recreational access, conservation, and property management throughout the state. On the Bailie Ranch, WDFW holds a 3,897-acre hunting easement. A number of other small easement are held for river and property access.

Leases

WDFW leases land from other entities and manages it under the wildlife area. About 180 acres are leased from the Department of Natural Resources on the Hope Valley and Windmill Ranch units, and 35 acres are leased from the Northern Pacific Railroad on the Byron Unit. Leasing land allows WDFW to fulfil management objectives of protecting waterfowl and small game habitat and providing access.

Agricultural leases

Agriculture is an integral part of the management practices on the wildlife area and provides multiple benefits for wildlife, habitat, and the local economy. It is an effective way to enhance forage and cover for wildlife, and provide weed and erosion control.



Canola fields at Windmill Ranch Unit



Wildlife area and regional staff negotiate leases, develop farm plans in collaboration with lessees, and oversee farming activities on leased sites. Leases are negotiated and designed to meet needs of the agency, wildlife, the farmer, and the community. Specific wildlife goals may be included, such as increased food and cover for upland birds, increased diversity on the landscape, and forage for deer. Practices to maintain pollinator habitat on WDFW owned and managed agricultural and grazing lands should be implemented where practicable. Funds generated from these leases supplement local budgets for the management activities on the wildlife area. As of 2019, 1,918.2 acres are in agriculture.

Unit	Agriculture acres	Expiration
Hope Valley	29.8	2024
Thornton	1,240.5	2019
Windmill Ranch	432.0	2022
Byron/Ferry Rd	101.2	2022
Sunnyside	339.0	2023
TOTAL	2,142.5	

Table 3: Agricultural leases

Grazing permits

Grazing is allowed under certain conditions on the wildlife areas, subject to specific grazing management plans, and in accordance with <u>WAC 220-500-200</u> and the Fish and Wildlife Commission <u>Policy C-6003</u>, Domestic Livestock Grazing on Department Lands. There are permits for cattle to graze on agricultural fields on the Windmill Ranch and Mesa Lake units. Cattle graze fields after the harvest to consume cornstalks and other crop residue which makes it easier for waterfowl and sandhill cranes to access the remaining spilled grain.

Temporary use permits

WDFW allows commercial and group activities on wildlife areas with a permit from the regional office. Non-commercial group activities of 30 participants or more must have a permit. For commercial operators such as rafting companies or hunting and fishing guides, a fee-based permit is required to operate on the wildlife area. The wildlife area issues a couple of permits per year.

Water use

WDFW holds seasonal water rights that are used to flood habitat for waterfowl, shorebirds, and other wildlife found in the wetlands. WDFW's agriculture leases are irrigated with water supplied by the South Columbia Basin and Sunnyside Valley irrigation districts. This provides winter forage for waterfowl, upland birds, turkeys, and deer. Not all agriculture is irrigated; some agriculture close to the river is grown by using the moist soil method.

Ownership and use of adjacent lands

The Sunnyside-Snake River Wildlife Area units are scattered throughout Yakima, Benton, and Franklin counties, where much of the adjacent property is in private agricultural ownership. This



juxtaposition within the converted landscape matrix increases the importance of these units as islands of habitats, stepping stones of connectivity, and refugia, breeding sites, or stopover areas for many wildlife species. Hops, grapes, orchards, corn, potatoes, onions, alfalfa, and wheat are the primary agriculture near the wildlife area units. The Sunnyside and I-82 units share the Yakima River as a common boundary with the Yakama Indian Nation. The Byron Unit is bordered on the north by the City of Grandview's water treatment facility. The Thornton Unit shares a common two-mile boundary with DNR on the west side of the property. About one-third of the DNR section is enrolled in the CRP program; the remainder is high quality shrubsteppe, which is grazed periodically under a private lease. The Rattlesnake Slope Unit borders the Arid Lands Ecology Reserve (ALE), which is owned by the Department of Energy as part of the Hanford Reach National Monument and is managed by the US Fish and Wildlife Service. The Saddle Mountain National Wildlife Refuge is also part of the national monument.

Rattlesnake Slope contains in-holdings owned by the Bureau of Land Management (BLM) and shares a common boundary on the north with ALE. The Rattlesnake Slope Unit is separated from the Thornton Unit by two separate, private ownerships. The Windmill Ranch was purchased as two



Crop from Sunnyside Unit Photo by Alan L. Bauer

separate parcels but a small parcel that is owned by DNR connects them. A small parcel in the SW corner, owned by the Bureau of Reclamation (BOR) makes up part of this management unit.

The Bailie Memorial Youth Foundation shares one mile of common boundary with DNR, which is held under a private lease for grazing.

The Hope Valley Unit (formerly Nipper Unit) borders DNR property on the east and BOR property on the south (Clark Pond). All other lands within the wildlife area complex are bordered by private lands, which are generally managed for some type of agriculture.

Interagency agreements

WDFW cooperates across jurisdictions as this is crucial for sharing resources to get things done across boundaries and at a landscape scale, and to coordinate activities. WDFW manages the Esquatzal Coulee and Thompson Seeps units which are owned by BOR for wildlife habitat and public recreation. WDFW manages these like WDFWowned units. Benton City conversely manages the Benton

City Unit, owned by WDFW, for recreation, and takes care of maintenance and operations.

Management setting

Administration

Day-to-day management of the Sunnyside-Snake River Wildlife Area is the responsibility of staff based out of the offices and shop located in Sunnyside and Pasco. Personnel consists of one full-



time Wildlife Area Manager, two Assistant Wildlife Area Managers, two to three Natural Resource Technicians, one full-time Access Manager, and one career seasonal Natural Resource Worker.

Other activities, such as wildlife surveys and development of species, habitat, wetlands and floodplain recovery are often undertaken by other agency staff and experts in coordination with wildlife area staff.

Operating funds

Operating funds to manage the Sunnyside-Snake River Wildlife Area come from two main sources: locally generated revenue from agriculture leases, and mitigation funds from Bonneville Power Administration. WDFW issues Agriculture Permits to local farmers and receives payments for the right to farm cropland on the wildlife area. This revenue is used to support staff salaries and habitat projects on the wildlife area. In 1996, as part of a mitigation agreement for habitat losses due to dam construction on the Columbia River, the Bonneville Power Administration (BPA) agreed to fund reasonable operation and maintenance activities on the wildlife area

Facilities and maintenance

Activities on WDFW lands include maintaining fences, roads, trails, signs, campgrounds, facilities, and performing weed control. The goal is to ensure wildlife area facilities and infrastructure remain in good working order over time. There are 12 structures located on the units that include two barns, two shops, six storage sheds, and two historic buildings (former residence and a wooden building), and other miscellaneous farm/ranch structures.



Crop harvest Photo by Phillip Buser

Fences are used to delineate

property boundaries where applicable. Some interior fences still remain from previous landowners. These fences are being removed as time and staff capacity allow.

Road management

All of the roads within the wildlife area are administrative roads, and are closed to public motorized access for the protection of wildlife and habitat. Each unit of the wildlife area is relatively small and the need to use motorized vehicles to access the interior portions of the unit is not necessary. Parking lots and access points are mostly located on the periphery of each unit. There is some ADA access for hunting allowed for public access on the Sunnyside and Thornton by arrangement.



Local land use designation

The 15 units of the Sunnyside-Snake River Wildlife Area are located in three counties of Yakima, Benton, and Franklin. Zoning of the units is not consistent across all three counties, but all reflect the local landscape where each unit is located. For example, even though the Whitstran Unit is located along the Yakima River, it is Zone RL (Rural lands)-5, which allows for residential development on up to 5 acres. Likewise, the Benton City Unit is also along the Yakima River, but is within the city limits. Land use must be consistent with each county's Comprehensive Plan, Natural Resource Ordinance, Critical Areas Ordinance, and Shoreline Management Plan. Table 4 describes the relationship of these land use regulations to the wildlife area land.

Wildlife area unit	Comprehensive Plan land use designation and zoning*	Shoreline Management Plan designation
Yakima Co.		
Byron	Remote/Extremely limited	Conservancy
Vance-Ferry	Agriculture	Rural
Sunnyside	Remote/Extremely limited	Conservancy/Rural
Glover	Agriculture	
I-82 Ponds	Remote/Extremely limited	Conservancy
Benton Co.		
Rattlesnake Slope	Park District	
Thornton	Agriculture	
Benton City	Industrial	Conservancy
Whitstran	RL-5	Conservancy
Franklin Co.		
Hope Valley (<i>Clark Pond</i> and Nipper)	Agriculture	Rural Conservancy
Mesa	Agriculture	Rural Conservancy
Esquatzel	Agriculture	
Windmill	Agriculture	
Bailie	Agriculture	Rural Conservancy
Thompson Seeps (WB-10)	Agriculture	

Table 4. La	and use de	esignations	by wildlife	area unit
I UDIC II D	unu use u	Signations	by whathe	ai ca unit

Yakima County

Lakes:<u>https://www.codepublishing.com/WA/YakimaCounty/html/YakimaCounty16D/YakimaCounty16DAppxB.html</u> Rivers: <u>https://www.codepublishing.com/WA/YakimaCounty/html/YakimaCounty16D/YakimaCounty16DAppxC.html</u> Zoning:

http://yakimacounty.maps.arcgis.com/apps/webappviewer/index.html?id=1bf3a144659f4a72a140973904f17612 Benton County

Shoreline: http://benton.municipalcms.com/files/documents/document174120123051013.pdf

Zoning: <u>https://bentonco.maps.arcgis.com/apps/MapSeries/index.html?appid=85029c8fdb784e1eb0bf08479d0cd595</u> Franklin County

Shoreline: https://www.co.franklin.wa.us/planning/smmp.html

Zoning:

http://franklinassessor.maps.arcgis.com/home/webmap/viewer.html?webmap=49e9b20342de4107b437dca9ef3a574f



Cultural Resources

State and federal law requires the protection of cultural, geological, and other non-renewable resources. Such resources may not be removed unless determined to be beneficial to wildlife, habitat, or scientific or educational purposes. WDFW coordinates with appropriate agencies and tribes for the protection of such resources if any activity affects cultural, archaeological, or historic resources. This includes the removal of various rock formations, Native American artifacts, plants, seeds, and other items. Wildlife area staff have received training in the importance of protecting the cultural resources on the wildlife area. Any action on the wildlife area that disturbs the ground requires a cultural resources survey. See Appendix E for a brief cultural history of the area.

Enforcement

Enforcement on the wildlife areas is provided by WDFW enforcement officers who have general authority peace officer status for the State of Washington. The Mission for WDFW enforcement officers is, "To protect our natural resources and the public we serve." WDFW's enforcement officers perform a wide range of duties to protect natural resources, the communities and economies that rely on them, and those who recreate outdoors. WDFW officers approach enforcement in four ways: enforcement, education, partnerships, and community involvement.

Their highest priority is enforcement of all fish, wildlife, and habitat laws under <u>Title 77 RCW</u>. Officers often deal with issues related to poaching, threatened and endangered species protection, habitat protection, and destruction of habitat. A core duty for WDFW Officers is protecting public safety in the outdoors, and they participate in a variety of enforcement activities related to this, including enforcing boating, off-road vehicle and snowmobile laws, and eradicating illegal drug growing and manufacturing. Officers work closely with emergency management agencies and play an important role in emergency management statewide.

Public conduct issues on this wildlife area include vandalism, littering and garbage dumping, especially on the western units, typically at water access sites and parking areas (I-82 Ponds, Sunnyside). Efforts by a volunteer citizens group, Eyes in the Woods, using surveillance cameras and signage have reduced but not eliminated illegal activities. WDFW is searching for innovative answers to curtail this issue, as litter pick takes an inordinate amount of staff time.

Signs posted on the wildlife areas are very specific to what activities are and are not acceptable. As in all wildlife areas, more enforcement presence is needed to handle all of the issues that occur. As part of this plan, WDFW will take actions to improve safety and security, and explore options of increasing citizen involvement in reporting illegal activities. The wildlife area staff will continue to encourage hunters and all other users to report suspicious or illegal behavior, and things that the managers should know about. Report illegal or dangerous activity to WDFW Enforcement at 1-877-933-9847, or 911 if you observe poaching in progress.



Research and studies

Consistent with WDFW's mission to preserve, protect, and perpetuate fish, wildlife, and habitat, WDFW supports independent studies to achieve wildlife area objectives.

Researcher	Date	Description
Jennifer Andreas - WSU extension Puyallup	Ongoing	Bio-control insects on noxious weeds, Sunnyside and Byron Units
WDFW	Summers	Waterfowl banding for Pacific Flyway and WDFW research
WDFW, Audubon Washington	2015-2019	Columbia Plateau shrubsteppe avian obligate songbird occurrence study
Michael Casazza – USGS Western Ecological Research Center	2017	Tracking blue wing and cinnamon teal with satellite transmitters
WDFW	2016-17	Statewide Monarch and Milkweed Surveys
WDFW	2016	Statewide Ferruginous Hawk Surveys
Marc Divens, Larry Phillips	2000	1998 Warmwater Fisheries Survey of Powerline Lake (Franklin County)
Marc Divens, Heather Woller, Randall Osborne	2003	Warmwater Fisheries Surveys of the I-82 Ponds (Yakima County) 2000 & 2001
Marcus J. Divens	2013	2012 Warmwater Fisheries Survey Of Mesa Lake (Franklin County), Washington
Chris Sergeant	1997	Clark Pond Fishery Analysis

Table 5: Research activities on Sunnyside-Snake River Wildlife Area

Recreation and stewardship

Recreation overview

WDFW wildlife areas provide fishing, hunting, and wildlife-related recreation opportunities, consistent with the agency's mission, the statewide wildlife area planning goals, and with the funding sources for each property. Public use is influenced by the character of the landscape, ability to access the area, fish and wildlife species present, as well as seasonal considerations and regional engagement from the local community. WDFW may place limitations on some activities in order to protect resources, preserve quality of experiences and infrastructure, and address the



Trout fishing at Powerline Lake Photo by Phillip Buser

safety of personnel and the public. The agency seeks to promote public enjoyment of fish and wildlife while managing and perpetuating them for future generations.



Washington State's population is growing, putting more pressure on wildlife areas across the state, including the Sunnyside-Snake River Wildlife Area. With more people comes a greater diversity of recreation interests, which can lead to conflicts between users. User conflicts can be detrimental to natural resources and can result in fewer quality recreational experiences. WDFW is developing a Statewide Recreation Plan to address these issues, which may lead to new rules, policies, and guidance to guide area management and to inform the public about where and how to recreate on WDFW lands. The strategy is expected to be completed in 2021.



Duck hunters at the Sunnyside Unit Photo by Alan L. Bauer

Recreation on the Sunnyside-Snake River Wildlife Area is predominately waterfowl hunting, along with upland bird, small game, deer and elk hunting, and fishing. There is a growing interest in bird watching, and visitors also enjoy walking, wildflower viewing, and horseback riding. No ATVs are allowed and some ADA hunter access is available at Sunnyside and Thornton units.

Wildlife area staff are responsible for managing public use. This includes providing accurate and up-to-date signage and information on the agency website, indicating what recreation activities are

permissible, and if any local restrictions apply. All state wildlife areas are governed by the agency's Public Conduct Rules <u>https://wdfw.wa.gov/about/wdfw-lands/public-conduct</u>), and may also have local requirements tailored to the area and its natural features, habitats and species. Additionally, many units of the wildlife require free on-site registration. Visitors must register on site when arriving, and report their activity before leaving. Registration provides usage and harvest data for future management of the unit. At the time of this plan, the units where registration is required are: Glover, Sunnyside, Hope Valley, Windmill, Mesa Lake, and Rattlesnake Slope. An objective of this plan to have registration at all the units.

Staff work with volunteers to make improvements to recreation areas, and collects public input to help prioritize funding needs (e.g. kiosks, parking areas, viewing platforms, etc.). The following table shows the major recreational opportunities and facilities by wildlife area unit.

Fishing: Fishing is popular on a few of the units, especially the I-82 ponds where anglers can catch bass and perch and other warmwater fish. Powerline Lake on the Windmill Unit is popular where largemouth bass, rainbow trout, black crappie, channel catfish, and yellow perch can be caught. On Mesa Lake bass, bluegill, and yellow perch can be caught.

Hunting: The primary hunting opportunities on the wildlife area are waterfowl, upland birds (pheasant, quail, and dove), and turkey in the spring. There is also limited small game deer and elk hunting opportunity. Pheasants Forever also helps administer some youth hunts. Always refer to WDFW's website for the current Hunting Pamphlet for seasons and rules. To protect other wildlife



species including waterfowl and raptors, non-toxic shot is required for all upland bird, dove, and band-tailed pigeon on all pheasant release sites statewide. This restriction applies to both shotshells and to loose shot for muzzleloading.

Target Shooting: Target shooting is allowed on the wildlife area only at the managed shooting range on the Rattlesnake Slope Unit. The land is leased to Benton County, who contracts with the Tri Cities Shooting Association to manage it as a multiple use, fenced range. There is a small fee and limited hours. There is no target shooting allowed anywhere else on the wildlife area.

Bird watching: The abundance of waterfowl, shorebirds, and migratory birds makes bird watching very good on most of the wildlife area. This is a result of the mixed wetland, riparian, upland, and shrubsteppe habitats across the wildlife area. Many



Pheasant release Photo by Jason Fidorra, WDFW

of the wetland areas maintain open water throughout the year. Migratory shorebirds and wading birds utilize the wetlands during the spring and summer. Waterfowl can be found across the wildlife area year-round. Many migratory songbirds can be observed using the riparian corridors during the spring, summer, and fall seasons. Shrubsteppe-obligate species are often observed in the uplands throughout the year. Each unit has something for the avid and novice bird watcher on any given outing.

Camping: Camping on the wildlife area is very limited and primitive in nature. Camping is allowed at only at Sunnyside, Thornton, Rattlesnake Slope, Thompson Seeps, and Esquatzel Coulee units. It is primitive camping, restricted to the parking lots, with no facilities.



Birdwatchers on Thornton Unit Photo by Jason Fidorra, WDFW

Table 6: Recreational use on Sunnyside-Snake River Wildlife Area

Wildlife	Primary hunting	Other	Water	Restrictions / conditions	Parking and other
area unit	and fishing	recreational activities	access		facilities
		uctivities	51105		
Bailie	Waterfowl & Upland bird Deer and elk (restricted) Fishing – private lease only			Registration required onsite. Privately owned ranch –WDFW has hunting easement. Open for hunting on Sat., Sun., Wed., and legal holidays. Limited fishing may be available by contacting the Bailie Youth Ranch. Limited creek crossing access. No overnight camping.	Two parking areas, each limited to five spots Kiosk
Benton City	Fishing	Birding Walking	2 water access sites	Managed by City of Benton City.	Parking lot, picnic tables
Byron - Main	Waterfowl & Upland bird Deer Warm water fishing (limited)	Birding Walking		Registration required onsite. Only open during the general hunting season.	Two gravel parking areas Kiosk
Byron - Reserve	No Hunting	Birding Walking		No hunting allowed. No guns, traps, or dogs. Open to walk-in visitors.	Unimproved parking area Kiosk, interpretive sign
Esquatzel Coulee	Waterfowl & Upland bird Deer Fishing	Walking Birding Horseback riding	Primitive		Gravel parking area
Glover	Upland bird	No other use		Surrounded by private land.	Gravel parking area Kiosk
Hope Valley (Clark Pond + Nipper)	Pheasant release site Waterfowl Upland birds Deer Fishing	Birding Wildlife viewing	Water access site and boat launch	Registration required onsite. Non-toxic shot only. Pheasant release site.	Two gravel parking areas Kiosks
I-82 Ponds	Waterfowl & Upland bird, turkey Fishing – Yakima River and borrow pits	Limited other uses	Water access (10)	Focus is on water access. Yakima River fishing requires a permit from the Yakama Nation.	10 parking areas associated with access sites, and some with facilities Kiosks
Mesa Lake	Waterfowl & Upland birds Fishing	Wildlife viewing Bird watching	Water access Boat launch	Registration required onsite.	Gravel parking area Kiosk
Rattlesnake Slope	Chukar, partridge Deer and elk	Horseback riding Hiking, Wildflowers Mountain biking Target shooting on range only		Walk in traffic only. No camping. Target shooting only on the range managed by third party. *	Gravel parking area Kiosk



Wildlife area unit	Primary hunting and fishing opportunities	Other recreational activities	Water access sites	Restrictions / conditions	Parking and other facilities
Sunnyside	Pheasant release site Waterfowl & Upland bird, dove Small game, turkey Deer	Walking Birding Horseback riding		Registration required onsite. Snipes Reserve is closed to all public entry. Non-toxic shot only. Yakima River fishing requires a permit from the Yakama Nation.	Multiple parking areas (six) Kiosks
Thornton	Upland bird, chukar, partridge Deer and elk	Birding Wildflowers Horseback riding Hiking		No fires. Master Hunters (special permits).	#1 Gravel parking area #2 Gravel drive-thru parking area for horse trailers Limited disabled hunter drive-in access is available by permit only
Vance-Ferry Road	Upland bird Fishing (limited access)	None	I mile of access along Yakima River	This site falls within a waterfowl hunting closure area. No waterfowl hunting is allowed within 1/4 mile of the Yakima River.	One parking area Kiosk
Thompson Seeps (WB-10 Wasteway)	Upland bird & Waterfowl Deer Fishing	Birding Hiking		The road/pond system is impacted by beaver and vehicle access may be impossible. Be prepared to walk in from Mt. Vista Road.	No specific parking area
Whitstran	Waterfowl & Upland bird (limited) Deer & elk Warm-water fish (limited) Fall Chinook salmon	Wildlife viewing	Water access on Yakima River	Access to this unit is only from Bunn Road.	Two gravel parking areas, one is seasonal only Kiosk
Windmill Ranch	Upland bird and waterfowl (restricted) Deer & elk Fishing & Ice fishing	Birding Wildlife viewing		Registration required onsite. 8 car limit during hunting seasons only. No overnight camping or open fires. See kiosk for further information/ restrictions. Foot traffic only.	Two gravel parking areas Wildlife viewing blind Kiosk

* The only place on the wildlife area target shooting is allowed at the shooting range.



Water access: WDFW manages more than 500 water access areas throughout the state for recreation associated primarily with boating and fishing. These areas occur within wildlife areas as boating or fishing facilities and beyond wildlife area boundaries as separately managed areas. This section provides information on water access areas that occur within the planning area.



Dock fishing at I-82 Ponds Photo by Alan L. Bauer

The wildlife area offers many opportunities to access the rivers, lakes, and ponds for fishing, boating, and water play. There is one fishing and boat launch access on the Columbia River, and multiple access points along the Yakima River, ponds off of the river, and ponds and lakes.

Powerline Lake, a popular walk-in fishing spot on the Windmill Ranch Unit, is not on the table below as it has no facilities. Information on fishing locations and regulations can be found at https://wdfw.wa.gov/fishing.



Boat launch at I-82 Ponds Photo by Alan L. Bauer



Sunnyside-Snake River Wildlife Area			Fishing and Boating Opportunities			Access Facilities				
County	Waterbody	Access	Unit	Public fishing easement	Fishing *	Hand launch	Trailered boat launch	Boat Ramp Surface	Toilet	ADA Parking
Benton	Yakima River	Benton City	Benton		•		•	Concrete	•	
		Duportail			•		•	Unimprove		
		Hyde Road			•		•	Gravel	•	
		Snively Road			•		•	Gravel	•	
Franklin	Clark Pond	Clark Pond	Hope Valley		•		•	Concrete		
	Columbia River	Ringold			•		•	Unimprove	• ADA	٠
	Mesa Lake	Mesa Lake	Mesa		•		•	Gravel		
Yakima	Buena Pond	Buena Pond	I-82		•	•			•	
	Ponds 1 & 2	Ponds 1 & 2	I-82		•				•	
	Pond 3	Pond 3	I-82		•				•	
	Ponds 4 & 5	Ponds 4 & 5	I-82		•	•			•	
	Yakima River	Granger			•		•	Gravel		
		Fitzsimmons	I-82		•	•				
		Mabton Bridge	Sunnyside		•		•	Concrete		
		Mellis Road	I-82	•	•	•			•	
		Zillah Bridge	I-82		•		•	Concrete	•	

* Fishing opportunities on department land. Refer to current WDFW sport fishing rules, as fishing seasons change and may not occur at all sites.

Stewardship and volunteerism

The Sunnyside-Snake River Wildlife Area has benefited from long-term participation of a number of groups and individuals who volunteer on a variety of projects to support the agency's conservation and recreation objectives. An important objective of this plan is to strengthen and continue to expand these partnerships and uncover more opportunities for interested parties to engage with the wildlife area.

WDFW staff are interested in exploring new opportunities and partnerships at the wildlife area that help highlight the educational and nature-based opportunities that the area could provide.

Developing nature interpretive signs and information, co-hosting nature hikes or wildflower walks, and bird watching and habitat enhancements are some of the ideas to explore in the next 10 years.

Table 8 shows some of the main ongoing volunteer opportunities on the wildlife area. Contact the



Pheasants Forever Volunteers Photo by Noel Ferguson, WDFW

wildlife area manager to propose ideas or to get involved with any on-going efforts.

Group	Types of activities
Washington Waterfowler's Assoc.	Building/maintaining duck blinds - late summer
Pheasants Forever	Tree and shrub planting in spring, assist with fall pheasant
Mule Deer Foundation	Potential collaboration on habitat projects - summer
National Wild Turkey Federation	Potential collaboration on habitat projects - summer
Richland Rod and Gun Club	Wood duck nest box cleaning - early spring
Various	Hunting blind clean up - early fall
Back Country Horsemen	Litter pick up, fence repair, other various projects
Runners of the Sage	Various as needed

Table 8: Ongoing volunteer opportunities Sunnyside-Snake River Wildlife Area

Conservation

Conservation of natural resources is the core mission of WDFW, and a driver for many actions on the wildlife area. Active management of water resources helps provide habitat for waterfowl and other wildlife. Restoration activities such as replanting after fires, weed control, restoring impacted areas to native habitat, maintaining a healthy riparian area and upland bird habitat, help to achieve more suitable habitats for wildlife. A key objective of this 10-year plan is to develop a long-term wetland management plan for habitat restoration and conservation.



Northern harrier Photo by Alan L. Bauer

Riparian areas and wetlands in an arid environment provide good habitat for birds and browse and cover for elk and mule deer. The wildlife area consists of variable sized parcels spread out over the landscape, so it is important to think on a landscape level, connect with other landowners, and work with partners on priorities. As part of the planning process, staff will identify priority areas for habitat restoration, and waters to improve and enhance for fishing.

The wildlife area supports three areas that provide refuge to wildlife, where no hunting, and sometimes no access, is allowed: 1) The Snipes

Reserve on the Sunnyside Unit; 2) the Byron Reserve on the Byron Unit; and 3) the Mabton Refuge on the Yakima River. One of the objectives of this management plan is to evaluate the function and benefits of the refuges and reserves to wildlife and recreationalists in order to promote their ecological value.



Sandhill cranes, Windmill Ranch Unit Photo by Alan L. Bauer



Wildlife area goals, objectives, and monitoring

Goals, objectives and performance measures

This plan sets management priorities for the Sunnyside-Snake River Wildlife Area for the next 10 years. The goals, objectives, and performance measures in this plan were developed by an interdisciplinary team of regional and headquarters staff, with input from the Wildlife Area Advisory Committee, tribes, the public, and other agency staff. They are consistent with WDFW's Mission and Strategic Plan. The plan goals, objectives, and performance measures will be reviewed and updated every two years. The objectives listed in this plan may or may not be fully funded. In many cases, successful outcomes will be dependent on

many cases, successful outcomes will be dependent on availability of funding.

Table 9 lists the goals, objectives, and performance measures of the plan. The "Tasks" column lists some of the steps that need to be taken or things to consider to achieve the planned objectives. While writing the goals and objectives, staff considered how projected changes in climate could impact the resources of the wildlife area and took note of opportunities that may help to mitigate or prepare for those impacts. These considerations are listed in the "Tasks" column were appropriate to the activity.

Monitoring and adaptive management

Wildlife area objectives will be measured annually based on the associated performance measures and through staff annual evaluations. On a biennial basis, wildlife area managers will review and work with staff leads to develop two-year updates with the advisory committee and district teams. Such reporting will allow the manager, staff, and regional office to modify tasks and timelines as necessary to meet plan objectives. Plan implementation may be affected by available funding.



Lupines on Thornton Unit Photo by Alan Bauer



Table 9: Goals, objectives, and performance measures

Goal	Objective	Unit	Performance measure	WDFW lead	Tasks
Statowida Goal: M	aintain or improve the ecological integri	ty of priority sites and sy	stome	and support	
Statewide Goal. IVI		ty of priority sites and sy	stems.		
1. Maintain or improve the ecological integrity of priority systems and sites.	A. Establish an ecological integrity baseline and associated goals for ecological systems of concern and priority systems and sites by 2024.	All	 Funding identified (Y/N) Baseline established (Y/N) Ecological Integrity goals established (Y/N) 	Ecological integrity Monitoring Team	 Work with WLA manager to design monitoring plan to achieve Objective A over 10-year planning term. Conduct data collection to determine baseline within 10-year planning term. Provide ecological integrity baseline report to WLA manager prior to start of subsequent 10-year planning term. Work with WLA manager to establish ecological integrity goals.
	B. Update and implement the 10- year plan for wetland management and restoration (Appendix D), which is focused on enhancements and also includes development of new wetlands on the wildlife area.	Sunnyside Byron Windmill Ranch Bailie Mesa Lake Thompson Seeps	 Number of acres managed annually Number of acres restored annually 	WLA Manager Habitat Biologist	 Prioritize areas for restoration. Work with partners to identify common goals. Identify funding sources. Work with conservation districts to help leverage opportunities with land owners. Tie in with state plans, conservation initiative, and other partners. Ensure not in conflict with upland bird habitat.
	C. Increase wetland function and value through vegetative restoration.	Sunnyside Byron Windmill Ranch Bailie Hope Valley	Vegetative restoration completed at: 1. Morgan Lake – Sunnyside Unit (Y/N)	WLA Manager	 Reduce emergent vegetation to provide 50 % cover / 50 % open water of wetlands. Utilize moist-soil management techniques to

Goal	Objective	Unit	Performance measure	WDFW lead	Tasks
			 2. Bridgeman Pond – Sunnyside Unit (Y/N) 3. All of the ponds at Byron Unit (Y/N) 4. Circle #3 Wetlands – Windmill Ranch Unit (Y/N) 5. V.I.P. Wetland – Bailie Unit (Y/N) 6. Bailie Lake – Bailie Unit (Y/N) 7. Clark Pond – Hope Valley Unit (Y/N) 	and support	increase natural food sources and improve overall wetland value. - Incorporate planted crops whenever practical to provide additional food and cover resources. - Manipulate water levels, as much as possible, to provide optimum recreational opportunities.
	D. Implement weed management plan (Appendix B) annually.	All	 Number of acres inspected Number of acres treated Annual weed control report produced (Y/N) 	WLA Manager	 Annually develop work plan using principles of IPM Complete annual reporting requirements. Respond to weed control needs after fires or other largescale disturbances. Coordinate with the Yakama Nation and County on purple loosestrife control. Coordinate w/ WSU Extension on Bio-Control efforts.
	E. Coordinate with Yakima Basin Integrated Plan (YBIP) work on Wapato Reach, which includes the I-82 ponds, as well as other areas of the Yakima River system.	I-82 Ponds	1. Number of projects on completed	WLA Manager Habitat Biologist	 Coordinate with YBIP on any projects on the Yakima R. Coordinate with the Yakama Nation.
	F. Coordinate with the City of Grandview to ensure	Byron	1. Flow to Byron Ponds improved (Y/N)	WLA Manager	- Work with City of Grandview wastewater



Goal	Objective	Unit	Performance measure	WDFW lead	Tasks
				and support	
	consistent water supply to Byron ponds for wildlife and recreation.		 Increased use by waterfowl and migratory birds (Y/N) Appropriate level of health 		treatment on water flow to Byron Ponds. - Coordinate with Mosquito
			restrictions applied (Y/N)		Control District. - Coordinate with Dept. of Health on access restrictions.
	 G. Develop a protocol for management of nuisance beavers on the wildlife area by 2020. 	Mesa Lake Hope Valley Windmill Ranch Sunnyside Bailie	1. Beaver Management Protocol Developed (Y/N)	Habitat Biologist Wildlife Biologist WLA Manager Enforcement	 Work with the irrigation district on plan. Work with the Yakama Nation. Work agency staff involved with this issue.
	 Identify upland areas with high potential for habitat improvement and develop action items by 2021. 	All	 Number of sites identified Number of action items implemented 	WLA Manager Habitat Bio	 Identify sites. Develop action items to protect habitat.
	 Maintain and restore floodplain functions throughout the Yakima River Valley. 	Benton City Whitstran I-82 Ponds Sunnyside	1. Number of floodplain restoration projects developed and implemented	Habitat Biologist WLA Manager Fish Biologist	 Coordinate with local partners on floodplain priorities (YBIP). Coordinate with Yakama Nation.
	J. Evaluate habitat loss from recent fires on Rattlesnake Slope and develop restoration strategy by 2020.	Rattlesnake Slope	 Habitat evaluated (Y/N) Develop strategy for habitat restoration (Y/N) Habitat restoration implemented (Y/N) 	WLA Manger Habitat Biologist	 -Work with Habitat Program to develop restoration strategy. - Secure funding for implementing restoration.
	K. Develop a prescribed burn plan for wetlands by 2020.	Sunnyside Byron Windmill Ranch Mesa Lake Thompson Seeps	 Plan developed (Y/N) Plan implemented (Y/N) Acres burned annually 	WLA Manager Burn Team Leader	- Work with fire districts and burn team to develop a burn plan.
	 Develop wetland habitat in Thompson Seeps Unit by 2022. 	Thompson Seeps	1. Overgrowth vegetation cleared (Y/N)	WLA Manager	 Work with Ducks Unlimited. Work with BOR. Work with Irrigation Districts.



Goal	Objective	Unit	Performance measure	WDFW lead	Tasks
Statewide Goal: Ac	chieve species diversity at levels consiste	ent with healthy ecosyste	 2. Low-head berms developed (Y/N) 3. Water control structures installed (Y/N) 	and support	
2. Achieve species diversity at levels consistent with healthy ecosystems.	A. Conduct survey for Species of Greatest Conservation Need in coordination with the Diversity Division by 2022.	All	1. Number of species surveys completed every 5 years	Diversity Wildlife Biologist	 Coordinate district priorities with Olympia Diversity staff annually. Contribute to Observations database.
	 B. Develop regularly scheduled waterfowl surveys for use in assessing management actions. 	I-82 Ponds Byron Sunnyside Windmill Ranch Bailie Mesa Lake Hope Valley Thompson Seeps	 Waterfowl survey developed (Y/N) Surveys conducted on a regular basis (Y/N) Number of times surveys used to determine success of management actions. 	Wildlife Biologist WLA Manager	 Determine appropriate information to collect. Develop and conduct surveys. Use information to evaluate management actions.
	 C. Identify and implement opportunities to enhance monarch butterfly, bumble bee, and other pollinator habitat by 2022. 	Hope Valley Mesa Lake Windmill Ranch Bailie Esquatzel Coulee Sunnyside Byron	 Number of units assessed for pollinator habitat Number of areas enhanced for pollinator habitat 	Wildlife Biologist WLA Manager	 Allow milkweed to grow undisturbed during the time period it supports monarchs. Conduct assessment of pollinator habitat. Collect data on the location of milkweed within the WLA. Develop action plan to enhance habitat. Determine which areas to plant shrubs, forbs, and native wildflowers.
	 D. Install nest platforms for ferruginous hawks to increase nesting opportunities by 2021. 	Thornton Rattlesnake Slope	1. Number of nest platforms installed	Wildlife Biologist WLA Manager	 Determine appropriate locations for nest platforms Install nest platforms



Goal		Objective	Unit	Performance measure	WDFW lead	Tasks
Statewide Goal: Su	рро	rt and maintain appropriate recreat	ion opportunities.		and support	
3. Enhance recreational experience through site development.	Α.	Focus recreation enhancement and management actions on the most popular units to improve recreational experience by 2021.	Sunnyside Rattlesnake Slope Mesa Lake Windmill Ranch Esquatzel Coulee Thornton	 Recreation enhancement projects identified by 2021 (Y/N) Number of funding opportunities applied for and received. Number of projects implemented. 		 Identify potential projects and viewing opportunities. Develop projects in coordination with regional staff and CAMP. Apply for RCO grant funding and/or capital funding. Recreation strategy may inform actions. Partner with others for Wildflower walks and birding tours.
	B.	Continue to improve recreational experience, user expectations, and support of the wildlife area by providing information online, at kiosks, in maps, and directional signage.	All, with a focus on the five priority units: Sunnyside Rattlesnake Slope Mesa Lake Windmill Ranch Esquatzel Coulee	 Number of kiosks installed or improved Number of signs installed Website content updated annually 	WLA Manager Public Affairs Enforcement	 Coordinate with other WLAs engaged in similar pursuit. Keep website current. Construct/erect kiosks and informational signs for all access points, trailheads and parking areas as staff time and funding allows. Where applicable consider interpretive signage that describes species, habitat types, unique features, restoration projects. Install signs as staff time and funding allows. Informed by Lands Showcase work, develop positive information messages (not all focused on the Nos).

Goal	Objective	Unit	Performance measure	WDFW lead	Tasks
				and support	- Work with local DOT for adding directional signs on appropriate highways and
	C. Address seasonal closures that affect both the wildlife area and water access sites annually.	All	 Number of agreements reviewed Number of clarifying changes made Information on closure posted (Y/N) 	WLA Manager Water Access Manager	byways. - Review agreement with landowners on easement and determine appropriate closure period (set closure dates). - Get closure dates added to the website.
	 D. Install footbridge across irrigation canal at Hope Valley Unit by 2022. 	Hope Valley	 Funding secured (Y/N) Project designed (Y/N) Construct complete (Y/N) 	WLA Manager	 Identify and apply for grant, secure funding. Coordinate with Irrigation District. Design and construct bridge.
	E. Work with user groups to improve the existing trail on Rattlesnake Slope by 2022.	Rattlesnake Slope	 User groups engaged in discussion (Y/N) Number of funding opportunities applied for Current trail system mapped (Y/N) Maps and kiosk info developed (Y/N) Maintenance plan developed with local support (Y/N) 	WLA Manager	 Work with user groups including, Runners of the Sage, to improve trails. Create maps and post on kiosks and online. Work with user groups to develop a maintenance plan.
	F. Improve waterfowl hunting and recreational bird watching by restructuring and maintaining berms that form rice paddies by 2020.	Sunnyside	 Berms restructured (Y/N) Maintenance systems developed (Y/N) 	WLA Manager Regional Staff	 Restructure berms that create rice paddies. Work with other regional staff on system to maintain berms.



Goal		Objective	Unit	Performance measure	WDFW lead	Tasks
					and support	- Ensure movement of water between each cell.
	G.	Improve waterfowl hunting by working by increasing food resources and water supply in the Johnson Wetland by 2020.	Sunnyside	 Food resources increase in Johnson Wetland (Y/N) Adequate water supplies maintained during hunting season (Y/N) 	WLA Manager	 Work with Agriculture Permit Holder on plan to increase food resources. Work with the Irrigation District on plan for adequate water supplies.
	H.	Evaluate the function of the refuges and reserves for hunting and recreation opportunities by 2021.	Sunnyside Byron Vance Ferry	 Evaluation developed (Y/N) Evaluation conducted (Y/N) Functions and values identified (Y/N) 	WLA Manager Habitat Biologist Game Division	 Coordinate with Game Division and regional staff on management. Work with managers of Toppenish Wildlife Refuge (USFWS) on regional management.
4. Improve fishing and hunting opportunities	A.	Assess potential to improve lakes and stock with fish (trout and/or warmwater) by 2024.	Priority units: Windmill Ranch Sunnyside Mesa Lake Also: Hope Valley I-82 Ponds Byron Thompson Seeps	1. Assessment completed (Y/N)	Fish Biologist	 Conduct a survey to identify possible fisheries. Use creel anglers and report catch.
	В.	Manage and improve upland bird habitat and nesting cover.	Priority units: Sunnyside Mesa Lake Esquatzel Coulee Also: Byron Glover Bailie Hope Valley	 Restoration projects developed projects (Y/N) Harvest data through self- registration cards (Y/N) Number of habitat improvements made 	WLA Manager	 Work with stakeholders to improve upland habitat. Minimize impact of other management activities during nesting season.



Goal	Objective	Unit	Performance measure	WDFW lead	Tasks
				and support	
	C. Identify opportunities to improve/increase waterfowl hunting by 2022.	All with waterfowl hunting	 Opportunity assessment completed (Y/N) Funding identified for opportunities (Y/N) Number of opportunities implemented 	WLA Manager District Biologist	 Identify areas for placement of blinds Identify field hunting improvement opportunities Survey area users to determine what improvements are desired.
	D. Improve fishing opportunities managing aquatic weeds in stocked lakes.	by I-82 Windmill Ranch	1. Number of weed control measures implemented	WLA Manager Fish Biologist Statewide WLA Weed Manager	 Collaborate with Fish Program on weed control efforts. Work with the Fish Program's fish stocking program (Ponds 1-6 and Powerline Lake). With Fish Program, evaluate other lakes/ponds/wetlands for potential stocking efforts.
	E. Expand Register to Use progra to all wildlife area units by 202	m All 1.	1. Number of sites enrolled in Register to Use	WLA Manager	 Install materials. Publicize change on website and through other means such as at meetings or through WAAC.
5. Improve access and other recreation opportunities	A. Improve infrastructure at wate access sites to provide moder amenities by 2024.	er All units with water access sites	 Number of parking improvements Number of restroom improvements Number of fishing enhancements Grants awarded, or projects funded 	WLA Manager Water Access Manager	 Prioritize list for improvements. Submit capital project requests or RCO grants.

Goal		Objective	Unit	Performance measure	WDFW lead	Tasks
	В.	Research areas where possible access or access agreements need to be updated by 2024.	Whitstran Esquatzel Coulee I-82 Ponds	1. Number of formal access agreements in place	Real Estate WLA Manager Water Access Manager	 Work with landowners and neighbors on agreements. Work with Real Estate
	C.	Maintain Disabled Hunter Access Program, including the Thornton ADA Special Permit for Elk hunting (limited entry special draw).	Sunnyside Thornton	 Disabled Hunter Access for upland bird implemented each year (Y/N) Disabled waterfowl hunter blinds maintained annually Y/N) 	WLA Manager	 Keep information up to date. Coordinate with ADA Coordinator. Maintain locations to keep them ADA accessible.
	D.	Open the west side of the Byron unit year-round	Byron	1. Westside of Byron open year-round (Y/N)	WLA Manager	- Work with the City of Grandview and the departments of Health and Ecology
Offer multiple and	varie	ed opportunities for stakeholder par	ticipation and engageme	nt.		
6. Offer multiple and varied opportunities for stakeholder participation and engagement.	А.	Coordinate and maintain a Wildlife Area Advisory Committee.	All	 Number of meeting(s) per year List of topics of concerns addressed Number of actions resulting from topics discussed 	WLA Manager	 Setup meeting time and place based on group members' availability. Draft agenda with attention to group interest and time constraints. Hold meeting and collect group comments and recommendations for consideration relative to future management actions (proposed or ongoing). Include meeting notes in wildlife area management plan updates and website.
	В.	Communicate with community groups about current wildlife area management activities.	All	1. Number of group/constituents contacted	WLA Manager	 Provide WLA information to local organizations, through email, telephone calls, community group meeting



Goal	Objective	Unit	Performance measure	WDFW lead and support	Tasks
					attendance and presentations, and written notices and newsletters.
	C. Engage with local schools and offer educational or service opportunities for youth on the wildlife area.	All	1. Number of school projects completed on the wildlife area	WLA Manager	-Coordinate with local schools - Coordinate with local organizations such as scouts -Identify projects or education opportunities.
	D. Continue to recruit new hunters and offer hunter education opportunities.	All	 Number or recruitment actions Number of hunter education opportunities offered 	Hunter Education WLA Manager	 Develop or distribute hunter recruitment information. Recruit at appropriate events. Offer hunter education.
	E. Work with local community tourism associations to communicate opportunities and benefits on the wildlife area.	All	1. Number of stories or events promoted on the wildlife area annually	WLA Manager Lands Messaging Team	 Work with internal Lands Messaging Team to develop messages, stories, and promotions on the wildlife area. Develop 1-2 stories each biennium or as opportunities arise.
Maintain productiv	e and positive working relationships wit	h local jurisdictions, com	munity neighbors, lessees, and p	ermittees.	
7. Maintain productive and positive working relationships with local jurisdictions, community, neighbors, lessees, and permittees.	A. Assess potential boundary issues on Rattlesnake Slope by 2020 and others as necessary.	Rattlesnake Slope	1. Rattlesnake Slope boundary issues assessed (Y/N)	WLA Manager	- Work with Lands Agent to determine legal boundary



Goal	Objective	Unit	Performance measure	WDFW lead	Tasks
	 B. Ongoing, as agricultural leases come up for renewal or for new ones developed, evaluate agricultural leases per agency policy for multiple benefits, especially to improve waterfowl forage for hunting opportunities. 	Sunnyside Byron Vance-Ferry Thornton Windmill Mesa Lake Hope Valley Glover	 Leases reviewed by District Team (Y/N) Modification made to lease if needed to provide additional wildlife benefit (Y/N) Monitoring for improvements conducted (Y/N) 	WLA manager District Team	 Review WDFW agriculture policies. Work with District Team to review and modify leases. Monitor effect of new lease modifications.
	C. Maintain strong communication and coordination with local jurisdictions on implementation of the plan.	All	 Appropriate local jurisdictions consulted on land use actions (Y/N) 	WLA Manager	- Consult with appropriate jurisdictions on actions that need their input.
	D. Maintain agreement with Benton City to manage the unit for recreation	Benton City	1. Conditions of agreement with Benton City fulfilled (Y/N)	WLA Manager	- Communicate with Benton City about unit management
Properly train, equ	ip, and license WLA staff to meet operation	ional and management n	eeds of the WLA.		
8. Identity reliable sources of funding.	 Identify reliable funding pool for restoration and operations and maintenance funding annually. 	All	 Sources identified (Y/N) Annual allocation distributed to proposals (Y/N) 	WLA Manager Lands Division Manager	 Explore grant opportunities. Apply for grants, if available. Explore endowments, revolving funds/pool agreements), etc.
	 B. Continue to work with BPA on operations and maintenance budget, securing adequate, sustainable funding, and reporting, and ensuring mitigation obligations are met. 	Sunnyside Thornton Rattlesnake Slope Byron	 Sufficient operations and maintenance funding secured (Y/N) Annual contract submitted for approval (Y/N) Annual report completed (Y/N) 	WLA	 Determine adequate operations and maintenance budget. Work with BPA on securing adequate funding. Fulfill mitigation obligations.
	C. Continue staff engagement after planning process through annual planning meetings.	All	1. Planning meetings held annually (Y/N)	WLA Manager Planning Team	- Coordinate with staff involved in planning and with actions in the plan.



Goal	Objective	Unit	Performance measure	WDFW lead	Tasks				
Maintain productiv	Maintain productive and positive working relationships with tribes.								
9. Maintain productive and positive working relationships with tribes.	A. Work with the Yakama treaty tribe to ensure the plan's management objectives of fish and wildlife are achieved while providing opportunities for the exercise of Treaty reserved rights.	All	1. Yakama Nation invited to discuss wildlife area plan management objectives and mutual concerns for wildlife resources (Y/N)	WLA Manager Lands Operations Manager Wildlife Program Manager Region 3 Director	 Coordinate with tribes on management actions that may be of interest to them. Extend annual invitation to Yakama Nation to discuss wildlife area management issues, wildlife resources, and other common interests. For management actions that could affect tribal interests, consider and evaluate impacts. 				
	 B. Protect tribal treaty rights and carefully evaluate and consider impacts to traditional hunting and gathering sites. Discuss mutual concerns for wildlife resources with the Tribes. 		 For management actions that could affect them, tribal treaty rights are considered and evaluated (Y/N) For management actions that could affect them, tribe is consulted (Y/N) 	WLA Manager Lands Operations Manager Wildlife Program Manager Region 3 Director	- For management actions that could affect tribal interests, consider and evaluate impacts.				
Maintain safe high	C. Assess project areas for culturally important plants before project implementation.	All	 Project area surveyed (Y/N) Culturally important plants protected or enhanced (Y/N) 	WLA Manager	 Be aware of potentially culturally important plants. Coordinate with the Yakama Nation. 				
Wantan Sarc, ngn		tration racintics and equi		1					
10. Maintain safe, highly functional, and cost effective administration facilities and equipment.	A. Remove derelict structures.	Sunnyside Mesa Lake	1. Number of derelict structures removed	WLA Manager Archeologist	- Consult with WDFW archeologist before removal for proper procedures.				
	 B. Maintain and upgrade water control structures as necessary. 	Sunnyside Windmill Ranch	1. Number of structures inspected	WLA Manager	- Coordinate with Ducks Unlimited when possible.				



Goal	Objective	Unit	Performance measure	WDFW lead	Tasks
				and support	
		Mesa Lake	2. Replacement/upgrade at Giffin Lake (Sunnyside) (Y/N)		 Seek grants and additional funding from other sources
			3. Replacement/upgrade at Bos Lake (Sunnyside) (Y/N)		
			4. Replacement/upgrade at Pie-Wedge Wetland (Mesa Lake) (Y/N)		
			5. Replacement/upgrade at Windmill Lake (Windmill Ranch) (Y/N)		
			6. Replacement/upgrade at Northeast Wetland (Windmill Ranch) (Y/N)		
	C. Annually inspect 20 % of fencing and all gates; repair and replace as needed and as funding allows.	All	 Amount of miles of fencing inspected or repaired (Y/N) Number of gates inspected and repaired Number of miles of new fence or replaced. 	WLA Manager	 Inspect 20 % of fence annually. Prioritize replacement of old fence. Complete repairs as needed. Submit Capital Funding requests for replacement of old fence.
	D. Explore best option for long-term management of the Vance-Ferry Unit.	Vance Ferry	 Options for potential disposition of Vance-Ferry Unit explored (Y/N) Decision reached on Vance Ferry Unit management (Y/N) 	Real Estate Lands Division Manager Wildlife Regional Manager Lands Operations Manager WLA Manager	 Review legal documents related to the property. Assess if it is meeting obligations. Determine final disposition.
	E. Continue working with BOR on agreements pertaining to BOR lands.	All BOR lands	1. Participation continued in BOR/WDFW quarterly coordination meetings (Y/N)	Real Estate Lands Division Manager	 Wildlife area manager, lands operation manager and wildlife regional program



Goal	Objective	Unit	Performance measure	WDFW lead	Tasks
			2. MOU revised (Y/N)	Wildlife Regional Manager Lands Operations Manager WLA Manager	manager, Lands Division Manager meet with BOR staff to negotiate MOU.
	F. Continue to work on strategy and funding to replace aging wildlife area equipment.	All	 Priority replacement list created (Y/N) Replacement plan developed (Y/N) Funding identified (Y/N) 	WLA Manager	 Work with Lands Division to identify need. Secure funding through Lands Division or BPA.
	 G. Update wildlife area facility information in centralized database quarterly. 	All	 Central facilities database updated annually (Y/N) 	WLA Manager	 Use agency facility inventory tool to update facilities information.
	 H. Review and update information on the wildlife area web pages annually. 	All	 Wildlife area web pages reviewed and updated annually (Y/N) 	WLA Manager	 Keep information available to the public on the web pages current.
	 Create a database on all the irrigation information, including diversions, wells, pumps, and maps of use areas, as well as instructions on operation by 2021. 	All	1.Database developed (Y/N)	WLA Manager	 Work with WDFW water rights staff, and Dept. of Ecology

Physical characteristics

Geology and soils

The geology of the wildlife area is quite varied. The mountainous areas in the western part of Yakima County consist of many different types of rock, including basalt and andesite. The principal rock in the central and eastern parts is Yakima Basalt, which is the younger flow of Columbia River Basalt. This basalt originated from large fissures or rifts where the fluid lava swelled to the surface and spread in all directions. Soils such as those in the Ritzville, Starbuck, Shano, and Bickleton series formed in areas where loess is underlain by basalt. Soils such as those in the Bakeoven, Lickskillet, Kiona, McDaniel, and Rock Creek series formed in colluvium and residuum derived from basalt. Overlying the Yakima Basalt in many areas that flank foothills and ridges are the light-colored tuffaceous sandstone, siltstone, and conglomerate of the Ellensburg Formation. This old stream-deposited sediment was derived from volcanic material ejected during the early development of the Cascade Range. This formation occurs extensively in the Wenas Valley, in the lower reaches of the Naches Valley, in areas west of Yakima, and along the southern part of Rattlesnake Ridge. The formation is more than 1,800 feet thick in places (Lenfesty and Reedy, 1985).

The upper and lower parts of Yakima Valley have been filled with material that was deposited by normal stream activity and glacial outwash. These areas include low terraces and floodplains. Extensive areas in the lower part of Yakima Valley are mantled by loess underlain by lake sediment that was deposited during glacial flooding in the late Pleistocene. This sediment occurs at elevations of as much as 1,000 feet in the survey area (Lenfesty and Reedy, 1985).

Yakima County units

I-82 Unit

Soils in the I-82 Unit consist mainly of the Weirman sandy loam series with some Zillah and Yakima series. These consist of excessively drained sandy loam soils on the low terraces and floodplains. It forms in mixed alluvium with slopes 0-5%. The native vegetation is mainly grasses, forbs and shrubs. This unit is subject to frequent periods of flooding in the spring. There are small patches of Yakima and Logy silt loam series that

are more suitable for grasses and forbs and was used in the recent past for crop production.

Sunnyside Unit

Soils in this unit consist of the silt loam series with most in the Zillah, Umapine, and Esquatzel series. This very deep, artificially drained soil is on floodplains and formed in recent alluvium. The native vegetation is mainly water-tolerant trees, sedges, and forbs. Included are areas of soils that have been artificially drained and areas of salt and alkali affected soils. Salt grasses and greasewood shrubs dominate these areas. These soils are prone to wetness and flooding.



Sunnyside Unit Photo by Justin Haug, WDFW



Byron Unit

Soils in this unit consist mainly of the Starbuck series. This shallow, well-drained soil is on the uplands and formed in loess. Soil depth is about 16 inches with basalt underlying it to a depth of 12-20 inches. The native vegetation is mainly grasses, forbs and shrubs. Native grass is mainly bluebunch wheatgrass and Sandberg bluegrass. The main shrub component is big sage and predominately rabbitbrush. Rock outcrops occurs in areas as exposed bedrock.

Vance-Ferry Road Unit

This unit lies adjacent to the Yakima River with silt loam and loamy fines soils predominating. The series are made up of Fiander silt loam, Kittitas silt loam, Umapine silt loam drained (2-5% slope) and Quincy loamy fine sand (0-10% slope). These are generally well drained soils on floodplains formed on terraces and on alluvium. The Quincy loamy fine sand was formed in eolian sand. Also included are very deep, artificially drained, salt and alkali-affected soil on the floodplains and low terraces and formed in alluvium. Native vegetation is mainly salt and alkali tolerant grasses, forbs and shrubs. Soil blowing can be a major problem and should have a constant cover crop of vegetation.

Benton County units

Whitstran / Benton City units

These units are located along the Yakima River. The Benton City Unit consists of the Finely fine sandy loam (2-5% slope) association while the Whitstran Unit has the Burbank loamy fine sand (0-2% slope) series as well as the Scootney stony silt loam (0-30% slope) and Scootney silt loam, gravelly subsoil (0-2% slope) series. These soils formed on alluvial terraces and on bottomlands along intermittent streams and the Yakima River. They developed under bunch grasses in stony and gravelly alluvium and in silty, windblown deposits.

Thornton Unit

The area includes six south facing parallel ridges separated by one permanent stream (Snipes Creek) and several intermittent waterways. Soils range from bare rock to over three feet deep allowing for the establishment of sagebrush stands and grasslands. Steep canyons bottom out at just under 1,500 feet in the southernmost part of the area while ridge top elevations extend to 2,400 feet on the north side of the unit. The unit consists of silt loam soils in the Kiona stony silt loam series, Lickskillet very stony silt loam series, Starbuck silt loam 2-15% slope series, Ritzville silt loam 15-30% slope series and the Moxee silt loam 15-30% series. These soils are generally well drained on uplands and formed in loess and colluvium with some derived from basalt. In the Starbuck series soil depth is about 16 inches with basalt underlying it to a depth of 12-20 inches. Rock outcrops occur in areas as exposed bedrock. In the Moxee silt loam 15-30% slope series the soil is well drained on uplands and is shallow over a hardpan and formed in loess. Lime and silica cemented hardpan is at a depth of about 18 inches and ranges from 10–20 inches. The hardpan is commonly underlain by basalt.

Native vegetation consists of grasses, forbs, and shrubs. These include Bluebunch wheatgrass, Sandberg bluegrass, and Thurber needlegrass. As production of grasses decrease the proportion of sagebrush and rabbitbrush increases. Dustiness can be a problem and the sites should be disturbed as little as possible.



Rattlesnake Slope Unit

Soils series in this area are Ritzville silt loam (0-5% slope), Willis silt loam, shallow (0-15% slope), Kiona very stony silt loam (30-65% slope), and Willis silt loam (0-5% slope). These soils occur on the uplands and developed under bunch grasses in silty windblown deposits mixed with small amounts of volcanic ash. The Willis series are underlain by a lime-silica hardpan overlying basalt bedrock. The Willis silt loam, shallow series are most often found on broad ridge tops.

Franklin County units

Soil reports were not available for Franklin County however, soil maps were. Using other county descriptions the following series were available.

Hope Valley Unit

Soil type is the Quincy-Hezel-Burbank association and Warden-Sagemoor-Kennewick association. These are dry sandy soils on terraces and dunes that have formed under sparse dune vegetation or shrubsteppe vegetation in wind deposited sand or silt and loam over glaciolacustrine deposits from cataclysmic glacial outburst floods; most have low water holding capacity. Native vegetation is mainly grasses and shrubs.

Soil type for the most part is the Warden-Sagemoor-Kennewick association. Soil description is dry, silty, and loamy soils that formed in glaciolacustrine deposits from cataclysmic outbursts floods in the lower part and loess in the upper part. Warden soils are on terraces and are very deep and well drained. They formed in lacustine deposits that have a mantle of loess. The surface layer is silt loam and the subsoil is are silt loam and very fine sandy loam. Sagemoor soils are on terraces and are very deep and well drained. They formed in lacustrine deposits that have a mantle of loess. The surface layer and subsoil are silt loam. Kennewick soils are on terraces and are very deep and well drained. They formed in lacustrine deposits that have a mantle of loess. The surface layer and subsoil are silt loam. Kennewick soils are on terraces and are very deep and well drained. They formed in lacustine deposits are silt loam. Kennewick soils are on terraces and are very deep and subsoil are silt loam. Kennewick soils are on terraces and are very deep and subsoil are silt loam. Kennewick soils are on terraces and are very deep and subsoil are silt loam. Kennewick soils are on terraces and are very deep and well drained. They formed in lacustine deposits and are silt loam throughout. Native vegetation is mainly grasses and shrubs.

Windmill Ranch /Bailie Units

Soil type is the Starbuck-Schawana-Prosser association. Soils of the Channeled Scablands are shallow, stony soils formed in loess over cata scoured basalt and occur in complex landscape patterns with moderately deep soil. Sandy or cobbly flood sediment formations are small areas of very deep loessial affected alluvial soils. The Starbuck soils are on benches, hillsides, and ridgetops. These are shallow and well drained and formed in loess and in material derived from basalt. Surface layer is very fine sandy loam with subsoil a silt loam. Schawana soils are shallow, somewhat excessively drained and on benches and hillsides. They formed in eolian deposits and material derived from basalt. The surface is cobbly loamy fine sand and underlying material is a gravelly very fine sandy loam. Prosser soils are on benches and hillsides and are moderately deep and well drained. They formed in loess and are very fine sandy loam. Native vegetation is mainly grasses and shrubs.

Hydrology and watersheds

I-82/Sunnyside/Byron/Glover/Thornton/Rattlesnake Slope/Vance-Ferry Road/Whitstran/Benton City units

All these units except Glover, Thornton, and Rattlesnake Slope lie along the Yakima River stretching from Union Gap to Benton City. Thornton lies along the south slope of the Rattlesnake Hills and Rattlesnake Slope



lies along the east slope of Rattlesnake Hills. Both eventually drain into the Yakima River, upstream and downstream of Benton City, respectively. All lie within Watershed Resource Inventory Area 37, the Lower Yakima River Watershed.

Thornton is bisected by the upper Snipes Creek and its small tributaries and lies adjacent to Sharp Road. Springs feed Snipes Creek part of the year with intermittent flows late in the summer with no fish known to reside in this section. Rattlesnake Slope Unit is bisected by three major canyons, none of which carry any water other than seasonal snowmelt or periodic, violent rainstorms.

The I-82 Unit has a relatively stable shoreline (Yakima River) due to the low bank and mature, woody riparian vegetation. The Sunnyside Unit for the most part is bordered by a very high vertical bank and is unstable. The exceptions are outward bends in the river, which continue to create deposition sites on the inside of the bends. Willows are the first woody species to become established on these bars. Annual runoff events scour the banks heavily against the outside bends of the river. It's not uncommon for the river to cut 20 feet off these banks in a single year, adding many tons of sediment to the river. The vertical nature and height of these shoreline areas makes it virtually impossible to stabilize them.



Yakima River waterways, I-82 Ponds Unit

The Sunnyside Unit lies mostly within the Yakima River floodway and much of it was underwater during severe flood events. The unit is laced with old river oxbows, isolated from the river as it has meandered across the valley over the years. These oxbows are still influenced by groundwater and some are artificially filled with irrigation wastewater to create more wetland habitat. Three major irrigation drains enter the Sunnyside Unit on their way to the Yakima River. At least two of these drains existed as natural streams before they were channelized and rerouted during the development of the irrigation project. Now, irrigation drain water and natural runoff are

commingled in these drains. An artificial and natural series of wetlands on the Sunnyside Unit is used to improve the quality of this water before it re-enters the river.

The Vance-Ferry Road Unit contains about three-quarters of a mile of river frontage and the shoreline condition is similar to that of the Sunnyside Unit. A shoreline buffer was established soon after the property was purchased. The Byron Unit has only two tenths of a mile of direct river frontage (similar to Vance-Ferry) and receives water from irrigation tailwater, drainage, or filtered wastewater from the City of Grandview. There is at least one small spring that enters the unit along the south boundary. Regardless of the origin, water from all these sources fills natural swales throughout the area, creating open water habitat, mixed with emergent vegetation. The water from the City of Grandview remains contained in a number of small basins, while the irrigation runoff leaves the property and eventually re-enters the Yakima River about a mile beyond the eastern boundary.

Both the Whitstran and Benton City units have Yakima River frontage. Both have relatively stable shorelines, minimal cutting, and support fair to excellent woody riparian cover.


Windmill Ranch/Bailie /Hope Valley Units

These properties lie within the Columbia Basin watershed with the bulk of water coming from irrigation wastewater that forms artificial lakes and wetlands creating fish habitat and recreational hunting and fishing opportunities. All of these units existed entirely of shrubsteppe habitat before the Columbia Basin Irrigation Project came on line in the early 1950s. Now, over 60 years of irrigation seepage out of canals and underground fractures in underlying basalt, the areas have taken on a completely different complexion. All of the units have at least some low lying depressions and swales that have collected water and become open water ponds and/or areas with hydric soils that support wetland vegetation. These units lie within Watershed Resource Inventory Area 36, the Esquatzel Drain.

The Bailie Unit has a substantial irrigation wastewater creek that flows year around along its entire length (over five miles). The channeled scabland nature of the property has produced several basins that are full of water on a year around basis. Some of the ponds are filled by underground flow and some are filled by direct surface flow. The Windmill Ranch has similar ponds and wetland habitat but most of it is created by seepage from the Potholes Canal, underground flow, or a side benefit of direct irrigation flow used to irrigate crops. The Windmill Ranch has three major water bodies that provide fishing and hunting recreation. The largest, Powerline Lake, is 35 acres in size, over a mile long and 70 feet deep.

Part of the Hope Valley Unit lies in a valley, which remains saturated by irrigation runoff most of the year. The gradient in this part of the valley is level and the presence of water has created highly alkaline soils.

Weeds and emergent vegetation, typical of this hydrology, are present on all but the west half, which is at a slightly higher elevation. As the water leaves the north end of the unit, it collects in Clark Pond, an old borrow pit. The water is then consolidated into a ditch as it continues to the south end of the unit. The south end is mostly an upland site but has three swales that contain wetland habitat and a small amount of open water, a result of seepage from the adjacent irrigation canal.

Climate

The wildlife area is mainly located in the Central Basin district climatic area. The following summary comes from the Western Regional Climate Center: Climate of Washington – Central Basin.

"The Central Basin includes the Ellensburg valley, the central plains area in the Columbia basin south from the Waterville Plateau near Chelan to the Oregon border and east to near the Palouse River. The elevation increases from approximately 400 feet at the confluence of the Snake and Columbia Rivers to 1,300 feet near the Waterville Plateau and 1,800 feet along the eastern edge of the area. This is the lowest and driest section in eastern Washington. Annual precipitation ranges from seven inches in the drier localities along the southern slopes of the Saddle Mountains, Frenchman Hills and east of Rattlesnake Mountains, to 15 inches in the vicinity of the Blue Mountains. Summer precipitation is usually associated with thunderstorms. During July and August, it is not unusual for four to six weeks to pass without measurable rainfall.

The winter season snowfall is from 10 to 35 inches. Snow can be expected after the first of December and to remain on the ground for periods varying from a few days to two months between mid-December and the last of February. Other than in the Ellensburg valley, snow depths seldom exceed eight to 15 inches. The Central Basin is subject to "chinook" winds which produce a rapid rise in temperature. A few damaging hailstorms are reported in the agricultural areas each summer.



The average January maximum temperature is near 30 degrees Fahrenheit in the colder localities in the Columbia Basin and 40 degrees Fahrenheit in the lower Yakima valley, and minimum temperatures are between 15 to 25 degrees Fahrenheit. Minimum temperatures between 0 to minus 10 degrees Fahrenheit are recorded almost every winter and temperatures from minus 15 degrees Fahrenheit to minus 30 degrees Fahrenheit have been recorded.

In July the average maximum temperature is in the lower 90s, and the minimum temperature is in the upper 50s. The recorded high temperature for the state, 118 degrees Fahrenheit, was recorded on July 24, 1928, at Wahluke, located along the southern slope of the Saddle Mountains and again on August 5, 1961, at Ice harbor Dam on the Snake River. Maximum temperatures reach 100 to 105 degrees Fahrenheit on a few afternoons each summer. The last freezing temperature in the spring occurs during the latter half of April in the Yakima valley and the latter half of May in the colder localities of the Columbia Basin. The first freezing temperature in the fall is usually recorded between mid-September and mid-October (https://wrcc.dri.edu/Climate/narrative_wa.php)."

Yakima River Corridor Units

The I-82 Ponds, Sunnyside, Byron, Vance Ferry Road, Whitstran, and Benton City units all lie along the Yakima River Corridor. Elevations range from 448 to 2,446 feet (at Thornton), and there is little variation in climate. As recorded at Sunnyside, the area receives an average annual precipitation in rainfall of seven and a half inches. Most precipitation occurs between October and May. Snow falls mainly in December and January, with an average annual snowfall of 10 inches, though it can vary from year to year. The average temperature is about 53 degrees Fahrenheit. Temperatures can go over the 90s in July and August, and down to the 20s in December and January (U.S Climate Data).

Units North of Pasco

The Hope Valley, Bailie, Mesa Lake, Ezquatzel Coulee, Thompson Seeps, and Windmill Ranch units north of Pasco also exhibit little change in elevation, from about 632 to 1,000 feet. As recorded at Connell, the area receives an average annual precipitation in rainfall of eight and seven tenths inches, with no snowfall. Most precipitation occurs between October and May. The average temperature is about 51 and a half degrees Fahrenheit. Temperatures can go over the high 80s in July and August, and down to the 20s in December, January, and February (U.S Climate Data).

Ecological values

Ecological systems and ecological integrity

WDFW's strategic objectives include protecting and restoring the ecological integrity of critical habitats consistent with DNR's Natural Heritage Program's Ecological Integrity Monitoring (EIM). The statewide goal is to restore and protect the integrity of priority ecological systems and sites. Ecological Integrity Assessments (EIA) and EIM are used to direct and measure achievements towards that goal.

Ecological integrity is defined as the ability of a system to support and maintain a community of organisms that has species composition, diversity, and functional organization comparable to those of natural habitats. EIM is a tool to evaluate ecological integrity, and changes over time, within priority systems and sites on the wildlife areas. Similar to species classifications grouped according to level of threat and potential inability to support sustained populations, habitats are grouped by type, including those that are priorities for



preservation and conservation. The complete classification system document, including descriptions of all ecological systems, can be found on this web page: <u>https://www.dnr.wa.gov/NHPecologicalsys</u> and summarized in the framework.

The planning process for Sunnyside-Snake River Wildlife Area identified seven National Ecological Systems of Concern to manage for ecological integrity. Table 10 summarizes these systems for the wildlife area, as described in DNR's Natural Heritage Program website. Listed are the systems considered "Critically Imperiled" and "Imperiled".

Appendix A contains the list of Species of Greatest Conservation Need (SGCN) believed to be present on the wildlife area and their relationships with ecological systems of concern. Actions associated with ecological integrity are included in the goals and objectives section (page 90), and include determining a baseline for ecological integrity for each of these systems and devising a monitoring plan to evaluate progress over time.



Ecological system of concern	Wildlife areas	Estimate acres	Description
Columbia Basin Foothill Riparian Woodland and Shrubland Critically imperiled	I-82 Ponds, Sunnyside, Thompson Seeps, Windmill Ranch, Byron, Bailie, Mesa Lake, Esquatzel Coulee, Vance-Ferry Road, Benton City, Whitstran, Hope Valley, Rattlesnake Slope, Glover	1945	Low-elevation riparian system found along the mainstem of the Columbia River and associated major tributaries on the periphery of the mountains surrounding the Columbia River Basin at and below lower tree line. Found in low-elevation canyons and draws, on floodplains, or in steep-sided canyons, in narrow V-shaped valleys with rocky substrates.
Inter-Mountain Basins Semi-Desert Shrub Steppe Critically imperiled	Bailie, Esquatzel Coulee, Windmill Ranch, Sunnyside, Byron, Mesa Lake, I-82 Ponds, Benton City, Whitstran, Hope Valley, Thompson Seeps, Rattlesnake Slope, Vance-Ferry Road	110	This semi-arid shrubsteppe is typically an open shrub to moderately dense woody layer and a strong graminoid layer (>25% cover but rarely closed). The woody layer is often a mixture of shrubs and dwarf- shrubs, although it may be dominated by a single shrub species. Characteristic species include Grayia spinose or Krascheninnikovia lanata with Ericameria nauseosa. Artemisia tridentata may be present but typically does not dominate although it will increase with disturbance.
Columbia Basin Foothill and Canyon Dry Grassland Critically imperiled - imperiled	Rattlesnake Slope, Thompson Seeps, Bailie, Esquatzel Coulee, Windmill Ranch, I-82 Ponds, Mesa Lake, Byron, Sunnyside, Thornton, Hope Valley, Vance- Ferry Road, Benton City, Whitstran	945	Foothill herbaceous vegetation found on steep open slopes, in the canyons and valleys of the Columbia Basin, particularly along the Snake River canyon, the lower foothill slopes of the Blue Mountains, and along the main stem of the Columbia River. Settings are primarily long, steep slopes of 328 feet to well over 1,300 feet, and slope failure is a common process.
Inter-Mountain Basins Big Sagebrush Steppe Imperiled	Rattlesnake Slope, Bailie, Esquatzel Coulee, Thornton, Thompson Seeps, Windmill Ranch, Byron, Mesa Lake, Sunnyside, Hope Valley, I-82 Ponds, Whitstran, Benton City, Vance- Ferry Road	10,215	This system is grassland with shrubs. Shrubs are dominated by Artemisia spp., and/or Purshia tridentata in an open to moderately dense shrub layer and with at least 25% total perennial herbaceous cover. The natural fire regime of this ecological system maintains a patchy distribution of shrubs, so the general aspect is that of grassland. P. tridentata is present almost always in association with tree cover, not out in the open.

Table 10: Ecological systems of concern on the Sunnyside-Snake River Wildlife Area (2015 Rocchio)



Columbia Plateau Steppe and Grassland Imperiled	Bailie, Thompson Seeps, Windmill Ranch, Sunnyside, Esquatzel Coulee, Thornton, Mesa Lake, Hope Valley, I-82 Ponds, Byron, Glover, Vance- Ferry Road, Whitstran, Benton City,	3,624	Extensive grasslands, not grass-dominated patches within sagebrush shrubsteppe ecological system, dominated by perennial bunch grasses and forbs, sometimes with a sparse shrub layer. Often forms a landscape mosaic with the Columbia Plateau Shrubland ecological system. Very little exposed bare ground due to mosses and lichens carpeting the area between plants, comprising a biological soil crust that is a very important characteristic in this
North American Arid West Emergent Marsh Imperiled	Sunnyside, Bailie, Byron, Thompson Seeps, Windmill Ranch, Mesa Lake, I-82 Ponds, Hope Valley, Esquatzel Coulee, Vance-Ferry Road, Whitstran, Thornton	948	Forests and tall shrublands that are linear in character, occurring on low-elevation, alluvial floodplains. Confined by valleys and inlets or lower terraces of rivers and streams.

Additionally, there are about eight acres of Northern Rocky Mountain Lower Montane Riparian Woodland and Shrubland (*imperiled*), about five acres of Northern Rocky Mountain Lower Montane, Foothill and Valley Grassland (*borderline status*), over one acre of Inter-Mountain Basins Greasewood Flat (*critically imperiled*), and over 100 acres of Columbia Plateau Scabland and Shrubland (*secure*). The remainder, about 3,300 acres, is not ranked and is a combination of cultivated cropland, open water, introduced vegetation, and developed areas.

Habitat connectivity

Fish and wildlife survival depends in part on the ability to move through the environment to find food and reproduce. The degree to which land protection and condition supports these necessary movements is called habitat connectivity. WDFW is a member of the Washington Wildlife Habitat Connectivity Working Group (WHCWG)

(<u>http://waconnected.org/</u>). This group represents a

science-based collaboration of land and resource



Byron Unit Photo by Alan L. Bauer



management agencies, non-governmental organizations, universities, and Washington Treaty Tribes.

Key wildlife habitat connectivity linkage networks at the statewide level were identified by the WHCWG in 2010 which looked at 16 focal species. A second examination of wildlife habitat connectivity linkages within the Columbia Plateau occurred two years later and looked at 11 species, WHCWG (2013). The Columbia Plateau is the largest ecoregion in Washington, occupying nearly one-third of the state. The Sunnyside-Snake River Wildlife Area is in this ecoregion. See the Columbia Plateau Ecoregion Analysis for more information.

These two connectivity efforts have some species in common. The Columbia Plateau Connectivity Analysis however, was performed at a finer scale since it was focusing on a subset of Washington State, not the entire state. (The default is to the Columbia Plateau Analysis when there is species overlap between the two studies.) The linkage networks, comprised of suitable habitats and the linkages connecting them, were derived from two modeling approaches: focal species and landscape integrity. The focal species approach identified important habitat areas for the species. The landscape integrity approach was used to help define the best linkages between habitat areas for each wildlife focal species found on or near the Sunnyside-Snake River Wildlife Area.

Focal species were carefully selected to represent the connectivity needs of a broader assemblage of wildlife (WHCWG 2012). The best linkages provided the least resistance to movement between habitat areas for that animal in that area. This means that some of the linkages may not be comprised of ideal habitat, but provide opportunities for movement through a human-modified landscape. The landscape integrity approach identified core habitat areas that were relatively free from human modification and the least human-modified linkages between them (WHCWG 2012).

Habitat connectivity information will be used to inform management decisions on the wildlife area. Habitat restoration and management projects will seek to maintain or improve linkages between habitat blocks on the Sunnyside-Snake River Wildlife Area for tiger salamander, badger, mule deer (statewide and Columbia Basin), black-tailed jack rabbit, white-tailed jack rabbit, western rattlesnake, Townsend's ground squirrel, Washington ground squirrel, least chipmunk, beaver, and elk (statewide). Habitat concentration areas and linkages for these species can be found online (see link above). The connectivity findings are a useful tool to assess important locations for the movement or migration of animals so they can reach the wildlife areas and move between wildlife area units.

Species management

WDFW's mission is to preserve, protect, and perpetuate fish, wildlife, and ecosystems while providing sustainable fish and wildlife recreational and commercial opportunities. The agency carries out this mission according to state and federal laws (including the Endangered Species Act or ESA) and funding requirements (from property acquisition and/or funds used for ongoing operations and maintenance), which direct many management activities on WDFW's wildlife areas. Other guidance comes from statewide plans for species and/or habitats, and other scientific approaches recommended by internal and external parties (e.g. The Washington State National Heritage Program's Ecological Integrity Assessments). Management actions may also be influenced by collaborative work undertaken with other conservation organizations, including tribal governments, land trusts and other land management organizations, academic research programs, and even the specific interests of volunteers if they fit within WDFW's mission, budget and wildlife area goals.



Species management overview

Consistent with WDFW's mission, the agency manages species on wildlife areas for two primary purposes: 1) conservation and protection to manage sustainable populations; and 2) provision of recreational and



American avocet, Sunnyside Unit Photo by Noel Ferguson, WDFW

commercial opportunities.

The Wildlife Area Management Planning Framework describes how species are classified – including species listed at the state or federal level as threatened or endangered, as well as other designations such as Species of Greatest Conservation Need (SGCN). SGCN species are summarized in the 2015 State Wildlife Action Plan (https://wdfw.wa.gov/species-habitats/atrisk/swap) and defined as species already listed as threatened, endangered or sensitive, as well as additional species thought to need conservation attention. The framework also incorporates goals from WDFW's Game Management Plan, which includes protecting, sustaining, and managing hunted wildlife, providing stable, regulated recreational hunting to all

citizens, protecting and enhancing wildlife habitat, and minimizing adverse impacts to residents, other wildlife, and the environment. The wildlife area plan integrates these plans and priorities, and, in the goal and objectives section (page 90), defines specific actions to achieve them.

The Sunnyside-Snake River Wildlife Area supports a broad variety of game and non-game species (also known as diversity species). Mammals include several species of bats, black-tailed jackrabbit, white-tailed jackrabbit, Townsend's ground squirrel, and possibly Washington ground squirrel, American badger, Rock Mountain elk, and Rocky Mountain mule deer. The area provides important breeding habitat for waterfowl, including mallards, wood duck, cinnamon teal, hooded merganser, and redhead, as well as stopover sites and winter habitat for a vast number of



Black-necked stilt, Sunnyside Unit Photo by Noel Ferguson, WDFW

duck and geese species. Upland birds include game species like California quail and pheasant as well as bald eagle, burrowing owl, loggerhead shrike, sage thrasher, and sagebrush sparrow.

The wildlife area provides habitat for the sagebrush lizard and possibly two other SGCN reptiles. With all of the wetlands, the Woodhouse's toad and tiger salamander find habitat. The monarch butterfly is on many of the units where milkweed is present, and the Columbia clubtail and white-belted ringtail dragonflies are found along the Yakima River in Benton County.



Several species of conservation concern likely utilize the Sunnyside-Snake River area due to its importance as supporting various natural and created wetlands in an otherwise arid environment, as well as providing a refuge of shrubsteppe habitat (sagebrush and bunchgrass) in an increasingly developed agricultural landscape. Two federally threatened species in the Yakima River are Mid-Columbia DPS bull trout and steelhead. The one federal candidate species, the greater sage grouse, historically used the area, but is not present now. The state endangered lesser sandhill crane uses the area for foraging during migration. Of the three state threatened species listed, the ferruginous hawk and American white pelican are known to be on the wildlife area, while the greater sage grouse is only historical.

Of the species that could be expected to appear on the wildlife area, there 21 State Candidate species, 45 SGCN species, and 40 PHS species (Table 11).

Table 11: Species conservation status

State and federal conservation status, WDFW Priority Habitats and Species (PHS) and SGCN criteria and priority areas that may occur on the Sunnyside-Snake River Wildlife Area

Common Name	Scientific Name	Federal Status State Status SGCN, PHS - 2019	General Location and Potential Wildlife Area Unit
MAMMALS			
Merriam's shrew	Sorex merriami	SGCN, PHS	Possible in shrubsteppe habitats, unconfirmed
Bat roosting concentrations	Eptesicus fuscus, Myotis spp., Antorzous pallidus	PHS	
Hoary bat	Lasiurus cinereus	SGCN	Possible on some units, unconfirmed
Silver-haired bat	Lasionycteris noctivagans	SGCN	Possible on some units, unconfirmed
Townsend's big-eared bat	Corynorhinus townsendii	SC,SGCN, PHS	Possible on some units, unconfirmed
Spotted bat	Euderma maculatum	SGCN	Possible on some units, unconfirmed
Black-tailed jackrabbit	Lepus californicus	SC,SGCN, PHS	Shrubsteppe habitats throughout
White-tailed jackrabbit	Lepus townsendii	SC,SGCN, PHS	Likely on Rattlesnake and Thornton
Townsend's ground squirrel (S of Yakima R.)	Urocitellus townsendii	SC,SGCN, PHS	Present. Benton/Yakima County upland units
Washington ground squirrel	Urocitellus washingtoni	SC,SGCN, PHS	Possible on Franklin County Units, unconfirmed
American badger	Taxidea taxus	SGCN	Present in upland habitat
Rocky Mountain elk	Cervus canadensis	PHS	Present in low numbers throughout; winter concentrations on Thornton and Rattlesnake Slope units
Rocky Mountain mule deer	Odocoileus hemionus	PHS	All
BIRDS			
American white pelican	Pelecanus erythrorhynchos	ST,SGCN, PHS	Foraging/staging areas on open water
Tundra swan	Cygnus columbianus	PHS	In winter on open water
Snow goose	Anser caerulescens	PHS	Spring congregations on open water
Black-crowned night- heron	Nycticorax	PHS	All river and wetland sites
Great blue heron	Ardea herodias	PHS	All river and wetland sites



Common Name	Scientific Name	Federal Status State Status SGCN, PHS - 2019	General Location and Potential Wildlife Area Unit
Western grebe	Aechmophorus occidentalis	SC,SGCN, PHS	Foraging/staging areas on open water
Black-necked stilt	Himantopus mexicanus	PHS	Breeding at wetland sites
Sandhill crane (lesser)	Antigone canadensis	SE, PHS	Present as foraging/staging areas during migration for Lesser subspecies
Cavity-nesting ducks: Wood duck, Barrow's goldeneye, common goldeneye, bufflehead, hooded merganser	Aix sponsa, Bucephala islandica, Bucephala clangula, Bucephala albeola, Lophodytes cucullatus	PHS	All Wood duck and Hooded Merganser nest boxes/cavities
Waterfowl concentrations	(Anatidae - excluding Canada geese in urban areas)	PHS	All
Cinnamon teal	Anas cyanoptera	SGCN	All Breeding at wetland sites
American avocet	Recurvirostra americana	PHS	Breeding at wetland sites
Wilson's phalarope	Phalaropus tricolor	PHS	Present at Wetland sites, possible breeder
Bald eagle	Haliaeetus leucocephalus	SGCN, PHS	All Present in riparian areas
Golden eagle	Aquila chrysaetos	SC,SGCN, PHS	Possible foraging in winter
Ferruginous hawk	Buteo regalis	ST,SGCN, PHS	Recent nest sites in Franklin County, possible any upland habitat
Peregrine falcon	Falco peregrinus	FSC, PHS	All
Short eared owl	Asio flammeus	SGCN	All Present in upland sites
Western screech owl	Megascops kennicottii	SGCN	All
Burrowing owl	Athene cunicularia	SC,SGCN, PHS	Present in upland habitats
Prairie falcon	Falco mexicanus	SGCN, PHS	Breeding sites on WLA
Loggerhead shrike	Lanius ludovicianus	SC,SGCN, PHS	Present in upland habitats
Chukar	Alectoris chukar	PHS	All
Greater sage grouse	Centrocercus urophasianus	FC,ST,SGCN, PHS	Historical/Recent unconfirmed sightings
Ring-necked pheasant	Phasianus colchicus	PHS	All
Sage thrasher	Oreoscoptes montanus	SC,SGCN, PHS	All Breeding Thornton Unit, large areas of sagebrush.
Sagebrush sparrow	Artemisiospiza nevadensis	SC,SGCN, PHS	Breeding on Thornton Unit, large areas of sagebrush.
AMPHIBIANS			
Tiger Salamander	Ambystoma tigrinum	SGCN	Likely in Franklin County units, unconfirmed
Woodhouse's toad	Anaxyrus woodhousii	SGCN	Breeding on Bailie Unit, possible any wetland unit



Common Name	Scientific Name	Federal Status State Status SGCN, PHS - 2019	General Location and Potential Wildlife Area Unit
REPTILES			
Desert nightsnake	Hypsiglena torquata	SGCN	Possible in shrubsteppe, unconfirmed
Sagebrush lizard	Sceloporus graciosus	SC,SGCN, PHS	All
Pygmy horned lizard	Phrynosoma douglasii	SGCN	Possible in shrubsteppe, unconfirmed
INVERTEBRATES			
Dragonfly: Columbia clubtail	Gomphurus lynnae	SC,SGCN, PHS	WA occurrence along Yakima River in Benton Co., not searched for on WLA
Dragonfly: White-belted ringtail	Erpetogomphus compositus	SGCN	WA occurrence along Yakima River in Benton Co., not searched for on WLA
Monarch butterfly	Danaus plexippus	Fed Review	Present throughout where milkweed present
Snail: Shortface Lanx	Fisherola nuttalli	SC,SGCN, PHS	Hanford Reach water access sites
Snail: Ashy (Columbia) Pebblesnail	Fluminicola fuscus	SC,SGCN	Hanford Reach water access sites
Western ridged mussel		SGCN	Yakima River species
Columbia River tiger beetle	Cicindela columbica	SC,SGCN, PHS	
Morrison bumble bee	Bombus morrisoni	SGCN	All
Western bumble bee	Bombus occidentalis	SGCN	All
Suckley cuckoo bumble bee	Bombus suckleyi	SGCN	All
FISH			
Bull trout (Middle Columbia River DPS)	Salvelinus confluentus	FT,SC, SGCN	Yakima River units
Steelhead (Middle Columbia River DPS)	Oncorhynchus mykiss	FT,SGCN	Yakima River units
Pacific lamprey	Entosphenus tridentatus	FSC, SGCN	Yakima River units
Mountain Sucker	Catostomus platyrhynchus	SC,SGCN	Yakima River units

Abbreviations:

State endangered (SE), State threatened (ST), State Candidate for listing (SC), State Sensitive (SS), Species of Greatest Conservation Need (SGCN), Priority Habitats and Species (PHS)

Federal endangered (FE), Federal threatened (FT), Federal candidate (FC), Federal species of concern (FSC) PHS Criteria: 1: State listed candidate species; 2: Vulnerable aggregations; 3: Species of recreational, commercial, or tribal importance.



Game species overview and management

The main big game species on the wildlife area are Rocky Mountain mule deer, which can be found on most units and attract hunters to the wildlife area. Rocky Mountain elk are also present in small and less predictable numbers on units in Benton and Franklin counties. Most hunters likely come to the wildlife area for small game, including excellent waterfowl and upland bird hunting (pheasant, quail, and dove). Some of the units of the wildlife area are too small to support sustainable game populations independently, and most species are dependent upon surrounding land, which is primarily private agricultural land. WDFW's 2015-2021 Game Management Plan (http://wdfw.wa.gov/publications/01676/) details management objectives and goals for Washington's game species. The overall goals support sustaining populations and providing recreational opportunities.

The wildlife area units often provide islands of cover for game species and opportunity for hunters in an otherwise mostly privately owned agriculture dominated landscape.

Most wildlife area units are open to hunting, but with specific restrictions at some units. Within the wildlife area, management actions can help maintain broader populations and harvest objectives for game species. These actions include invasive weed control, moist soils management, agricultural leases for forage and cover, habitat plots and restoration, pheasant releases for hunting, surveys, and research.

Hunting season dates and harvest restrictions are species specific within the state and across regions, with seasons and regulations evaluated and updated each year. Species populations receiving higher hunting pressure are monitored more intensely than those with lower participation rates, therefore season changes may occur more frequently. The specific regulations pertaining to individual species and hunting seasons are found on WDFW's website (<u>http://wdfw.wa.gov/hunting/regulations/</u>). Additional information on harvest history and population status are located in WDFW Game Harvest Reports (<u>https://wdfw.wa.gov/hunting/management/game-harvest</u>) and the Hunting Prospects published annually for District 4 and District 8: (<u>https://wdfw.wa.gov/hunting/prospects/</u>).

The Sunnyside-Snake River Wildlife Area units fall into the following Game Management Units (GMUs):

- 372 Rattlesnake Hills (units in Benton and Yakima County, aside from those listed for 373)
- 373 Horse Heaven Hills (Vance-Ferry and Byron Units)
- 379 Ringold (all units in Franklin County, aside from the Esquatzel Coulee Unit)
- 381 Kahlotus (Esquatzel Coulee Unit)

Game species

Mule Deer (Odocoileus hemionus)

Mule deer, associated with dry open expanses of eastern Washington, are year-round residents on many units of the Sunnyside-Snake River Wildlife Area. While white-tailed deer (*Odocoileus virinnnianus*) are occasionally present and make up an insignificant portion of deer (less than 1%) in the area. While white-tailed deer seasons are present to allow legal harvest during mule deer season, it is not a species hunters should expect to encounter in this area.

While WDFW does not conduct population surveys specific to the wildlife areas, surveys of the Columbia Plateau Mule Deer management zone indicate stable populations in key wintering areas. The Mule Deer Management Plan sets objectives and conservation priorities for the overall zone:

https://wdfw.wa.gov/publications/01755/. Objectives for the Columbia Plateau are stable populations with



a post-hunt sex ratio of 15 to 19 bucks per doe. Due to the open country and low buck escapement in the Columbia Plateau, a three-point-minimum antler restriction on harvest is in place for mule deer general seasons in the area.

Intact sagebrush and shrubsteppe habitat provides important forage and cover for mule deer. Management practices that promote healthy sagebrush stands will help provide winter forage and cover. Several shrub plantings and cover plots installed on the wildlife area can help increase overwinter survival and escapement of deer. Restoration of shrub components after a fire is also critical as sagebrush can take decades to reestablish after a burn, or may never outcompete non-native vegetation without assistance post-fire.

Fencing can be a hazard for deer across the landscape. Significant amounts of derelict and unnecessary fencing have already been removed from the wildlife area. Managers should continue to identify fencing for removal, and ensure any new fencing projects incorporate wildlife friendly designs to reduce risk to wild ungulates.

Agriculture leases on the wildlife area are being reviewed to maximize benefits for wildlife while maintaining sustainable funding for land management actions. Crop type, harvest methods, and forage left onsite can be altered to benefit deer and other wildlife and create great forage and cover in close proximity as well as enhanced hunting opportunities.

Elk (Cervus Canadensis)

Rocky Mountain elk in eastern Washington stem from reintroductions by sportsman's groups in the 1910s with elk transported from Yellowstone National Park. Elk are occasionally found on a few units of the Sunnyside-Snake River Wildlife Area, and their source population is in most instances likely the Rattlesnake Unit of the Hanford Reach National Monument managed by the USFWS. Elk were very rare in the mid-Columbia basin area until recently. Beginning with an initial report of a handful of elk on Rattlesnake Mountain in the 1970s, the population has grown within the refuge of the monument boundaries to its current size of over 1,100 individuals, despite a population objective of 350 elk. Elk now regularly leave the protected area may be found on wildlife area units including Thornton and Rattlesnake Slope. The lower elevations of the Rattlesnake Slope Unit can host many hundred elk during winter months. Small numbers of elk also cross the Columbia River and make their way onto the wildlife area in the Ringold GMU. Harvest still requires extreme luck and usually only a handful of elk are harvested during the general season in the Ringold or Rattlesnake Hills GMU. Most harvest during general seasons on the wildlife area likely occurs on the Thornton Unit.

Due to conflicts with crop producers and the over population of the Hanford Monument, there are no management actions specifically intended to increase elk populations on any unit of the Sunnyside-Snake River Wildlife Area.



Waterfowl

The location of the Sunnyside-Snake River Wildlife Area amidst the Yakima and Columbia rivers, along with



Mallards and Canada geese at Windmill Ranch Unit Photo by Alan L. Bauer

multiple irrigation projects and canals, provides an abundance of opportunity across an otherwise arid environment for breeding, staging, and wintering waterfowl and the hunters that pursue them.

The main breeding species on the wildlife area include mallard, gadwall, wood duck, and teal. Small numbers of redhead, ring-necked ducks, common and hooded mergansers, wigeon, Canada goose, and other species may nest here as well. During fall and winter, the full variety of western inland waterfowl can be found. Most open water is in shallow wetlands hosting puddle ducks, but some deeper lakes such as Powerline Lake on the Windmill Unit host diving ducks.

With the purchase of an amphibious

maintenance craft in 2015, wetland management actions such as mowing, spraying, and excavating became more feasible. Recent efforts have targeted invasive native and non-native aquatic species, with the intent to open choked up wetlands and improve waterfowl use and hunter access on wetlands lacking water control. At other sites, we use moist soil management to control weeds, and foster germination of seedbanks of forage plants like smartweed. We have also been experimenting with forage crops in recent years to provide more food for waterfowl and other species. The department has planted and left unharvested sunflowers, buckwheat, and other forage crop species to attract and feed wildlife in or around wetlands in small patches. Nest boxes for wood ducks and other cavity nesters have been installed, but maintenance is likely needed.

Regular monitoring of wetlands through the season is the best way to quickly respond to avian disease outbreaks. Carcass collection is a key to control of avian cholera and botulism, both of which have been reported from the Tri-Cities area but have not occurred on the wildlife area.

More information on waterfowl hunting can be found at: (<u>https://wdfw.wa.gov/hunting/regulations/migratory-waterfowl-upland-game</u>).

Upland Game Birds

Upland game birds are common on many units of the wildlife area. California quail, ring-necked pheasant, and mourning dove are the primary upland game available for hunters. Some hunters may encounter gray partridge or chukar in a few locations, such as Thornton and Rattlesnake Slope. Habitat plots, food plots, and agriculture lease management are all strategies that we employ on various units of the Sunnyside-Snake River Wildlife Area to promote upland bird populations.

The Sunnyside Unit and the Hope Valley Unit are both pheasant release sites. Planted birds are released prior to the youth and general pheasant seasons and sporadically throughout the season as a way to increase hunter participation. The birds released are all roosters and the purpose is solely for harvest. More



information on upland game birds and hunting can be found at: <u>https://wdfw.wa.gov/hunting/regulations/migratory-waterfowl-upland-game</u>.

Diversity species overview and management (Non-game)

The Sunnyside-Snake River Wildlife Area protects critical habitat in a region heavily converted to agriculture. Both riparian areas and shrubsteppe are considered critical habitats in the counties the wildlife area resides. Having an arid climate, the abundance of water on the wildlife area adds to the diversity of species the wildlife area helps support.

Very few surveys have been conducted on the Sunnyside-Snake River Wildlife Area, and species occurrence data is lacking for many species groups. Incidental observations have generally not been recorded in state databases, so even presence data is unknown for many species. A priority over the next few years should be to train and equip wildlife area staff and volunteers with the tools to log incidental observations through District Wildlife Biologists for submittal to state databases. Additionally, species specific surveys should be conducted to identify species groups such as reptiles, amphibians, odonates (dragonflies and damselflies), small mammals, and birds that are hypothetically using the wildlife area but have not been documented to date.

Diversity Species (non-game)

Sandhill crane (Antigone canadensis)

At the time of writing (2019), sandhill crane is the only state-endangered species known to utilize the wildlife area. The species is listed as endangered primarily because of the critically low breeding population

of Greater Sandhill cranes (*A. c. tabida*) that nest in Klickitat County. However, there are also greater sandhill cranes breeding in Canada mixed in with migrating lesser sandhill cranes (*Antigone canadensis*) in the Columbia Basin, which hosts important stopover sites for cranes in eastern Washington, including the Sunnyside-Snake River Wildlife Area. Agricultural fields on the Windmill Unit can host over 1,500 sandhill cranes during early spring during their northward migration.

While agricultural fields are currently not limited in the area during migration, the wildlife area can begin to manage agriculture leases for crops that are in attractive conditions to cranes during March.

Ferruginous hawk (Buteo regalis)

Ferruginous Hawks are a threatened species in decline in Washington. They naturally nest amidst vast open areas in arid shrubsteppe habitat primarily on cliffs, rim rock ledges, lone trees, and intact habitat. They have been successful ground nesters on bluffs or steep slopes. Some pairs have taken to artificial structures placed in open areas. The Esquatzel Coulee Unit of the wildlife area has had nesting ferruginous hawks in the past, and hawks have also nested in close proximity to the Thornton, Rattlesnake Slope, and Mesa Lake units. Any





Sandhill cranes at Windmill Ranch Unit Photo by Alan L. Bauer

site with suitable prey, especially ground squirrels, could be attractive to ferruginous hawks for foraging sites.

Control of ravens (nest predators and nest site competitors) may help ferruginous hawk populations, but reestablishment of prey species is likely the key factor in protection of the Washington population. The birds are also considered to be sensitive to disturbance.

Sites on the wildlife area where suitable habitat and prey exist, but cliff or nest substrate are lacking, would be ideal sites for installing nest platforms. Currently, a mitigation project is being planned which will install one or two platforms on raised poles on the Thornton and Rattlesnake Slope units. Continued monitoring of historic nest sites is also needed, and protection of occupied sites from March-July from disturbance would be needed.

Monarch butterfly (Danaus plexippus)

The monarch is a species whose decline has led to a petition for listing by the USFWS. Currently under federal review, the species' decline may be linked to habitat loss, systemic pesticides (such as neonicotinoids), and habitat destruction on wintering grounds. The species is an obligate breeder on milkweed plants. In 2016-2017, WDFW conducted monarch and milkweed training of wildlife area staff and biologists and surveys for the insect and its host plant. Monarchs and milkweed were located on many units of the Sunnyside-Snake River Wildlife Area, and often in dense patches. Larva were found on most units in



Monarch butterfly Photo by Justin Haug, WDFW

Franklin County that were surveyed. The training and attention has resulted in interest by local staff who are now protecting milkweed patches when encountered. Catastrophic declines in the western population continued and in 2019 surveys, no monarchs were found at the breeding sites in Franklin County.

Generally, milkweed is allowed to grow in areas where it will not be disturbed during the time period it supports monarchs (late May – early September). In areas where milkweed disturbance cannot be reduced or eliminated, milkweed should be controlled outside of the period it supports monarchs.

Pesticides and seeds used on the wildlife area by

staff, contractors, or leaseholders should be reviewed to avoid those containing neonicotinoids, which unnecessarily cause nectar of the planted species to be toxic to insects. Native bee species are also presumed to be in decline because of these pesticides.

Ground squirrels

Ground squirrels of the genus *Urocitellus* in the area include the Washington ground squirrel in Franklin County, and Townsend's ground squirrel in Benton and Yakima counties. The Washington ground squirrel is a protected species in Washington and illegal to hunt, while Townsend's ground squirrel is not protected, though shooting and poisoning are not encouraged. Both squirrels provide potentially important prey sources for ferruginous hawks and badgers.



Washington ground squirrels are not currently known on any units of the wildlife area in Franklin County;

however a site was reported in the early 2000s as being active on the Esquatzel Coulee Unit. No formal survey effort has been conducted on the wildlife area. The species can be hard to detect as they are active mostly from March through June and otherwise estivate to avoid the hot dry summer and cold winter. Often more easily heard than seen, the very highpitched thin whistle call is inaudible to many. Habitat includes shrubsteppe sites, usually with some sagebrush component.

Townsend's ground squirrels have been reported on units in Yakima County, including I-82 Ponds, Sunnyside, and Byron units. The species is fairly common in the area and likely occurs in more locations but incidental reports and surveys have not been



Townsend's ground squirrel, Byron Unit Photo by Alan L. Bauer

completed. Often persecuted for damaging irrigation ditches, managers should work with Irrigation Districts on the wildlife area to seek for alternatives to lethal removal such and improvements to canal structures.

WDFW and USFWS have worked to translocate ground squirrels in Adams County to reestablish populations in suitable protected habitat. If their methods work, some sites in Franklin County may be assessed as suitable release locations in the wildlife area.

Bats

Four bat Species of Greatest Conservation Need (SGCN) are likely present in the vicinity of the wildlife area, but there aren't good data to support this. These include Townsend's big eared bat which is a State Candidate Species for listing, spotted bat, hoary bat, and silver-haired bat. The only bat data in Wildlife Survey Data Management database from the wildlife area is a bat study done in 2007 by USFS staff that captured hoary bats and *myotis sp*. on the Sunnyside Unit.

Reptiles and amphibians

Two amphibian and four reptile Species of Greatest Conservation Need may occur on the Sunnyside-Snake River Wildlife Area. However, surveys of reptiles and amphibians on the wildlife area have not occurred, and few species have been reported. Tiger salamander (*Ambystoma tigrinum*) likely breeds in fishless wetlands on the units in Franklin County, and Woodhouse's toad (*Anaxyrus woodhousii*) have been found on the Bailie Unit but no reports exist elsewhere. Any expansion of the fish stock programs should address risk to amphibians.

Pygmy horned lizard (*Phrynosoma douglasii*), sagebrush lizard (*Sceloporus graciosus*), and side-blotched lizard (*Uta stansburiana*) are also likely to exist on suitable upland sites but no records exist. Desert nightsnake (*Hypsiglena torquata*) is likely to be found if appropriate searches were conducted and incidental sightings reported.



Water birds

In addition to waterfowl, the wetlands of the Sunnyside-Snake River Wildlife Area provide habitat for well over 100 species of birds. SGCN species include the state-threatened American white pelican, which uses the wildlife area for forage and roosting. Several sites provide PHS listed breeding sites for American avocets

(*Recurvirostra Americana*), black-necked stilts (*Himantopus mexicanus*), and possibly Wilson's phalaropes (*Phalaropus tricolor*) if water levels are adequately maintained. Tundra swans (*Cygnus columbianus*), snow geese (*Anser caerulescens*), and western grebes (*Aechmophorus occidentalis*) are SGCN that sporadically use the larger wetlands, sometimes in large numbers. During March of 2018, thousands of snow geese roosted on the lake in the Windmill Unit. Nesting sites of great blue herons (*Ardea Herodias*) and black-crowned night herons (*Nycticorax nycticorax*) may exist in riparian areas, but have not been recorded. With the added ability to manage wetlands with amphibious equipment, more monitoring and documentation of species occurrences are needed to measure management results across a suite of species and seasons.



White pelican, Sunnyside Unit Photo by Alan L. Bauer

Promoting public use of the wildlife areas during non-hunting seasons for birdwatchers and photographers provides the added benefit of possibly capturing species data through citizen science portals such as iNaturalist (https://www.inaturalist.org/) and eBird (https://ebird.org/home).

Shrubsteppe songbirds

Away from the wetland sites, sagebrush and bunchgrass provide native shrubsteppe habitat for many species of conservation concern. Sagebrush obligate species such as sagebrush sparrow (*Artemisiospiza*

nevadensis) and sage thrasher (*Oreoscoptes montanus*) can be found reliably breeding on and around the Thornton Unit, famous locally to birdwatchers. Loggerhead shrikes (*Lanius ludovicianus*) breed in similar habitat and have been reported on many wildlife areas units. Burrowing owls nest where burrowing mammals, including badger and coyote, create nesting sites for them, and have been reported at Rattlesnake Slope, Thompson Seeps (WB10-Wasteway), and Esquatzel Coulee units.

Protection and restoration of shrubsteppe habitat, especially from fire in arid areas such as Thornton, Rattlesnake Slope, and many Franklin County units is a priority. Control of nonnative vegetation, fire breaks, and post-fire restoration can all help reduce the frequency and impact of fire on the shrubsteppe landscape. See the Climate Change section for more information on shrubsteppe species.



Sagebrush sparrow Photo by Jim Cummins



Invertebrates

Several presumed rare, understudied, and underappreciated species that may occur on the wildlife area may either be protected, or negatively impacted by actions on the Sunnyside-Snake River Wildlife Area. Two of Washington's SGCN dragonflies are known from only a couple of reports from the state which have been made along the Yakima River. The Columbia clubtail (*Gomphurus lynnae*) and white-belted ringtail (*Erpetogomphus compositus*) have not been studied well in Washington, and surveys along units adjacent to the Yakima River may be interesting. The western ridged mussel (*Gonidea angulate*) is a SGCN bivalve also present only in the Yakima River. WDFW water access areas along the Columbia River Hanford Reach may impact or provide survey opportunities for two SGCN mollusks: shortfaced lanx (*Fisherola nuttalli*) and ashy pebblesnail (*Fluminicola fuscus*).



Lupine with bee, Thornton Unit Photo by Alan L. Bauer



Fish species overview and management

Fish species overview

Of the 15 units comprising the Sunnyside-Snake River Wildlife Area, 11 provide opportunities for recreational angling, either fishing the lakes located within the units or providing access to the Yakima River. There are 20 ponds located within the wildlife areas that support fish populations of sufficient size to provide angling opportunity. Fish communities within these lakes are all mixed populations of warmwater (non-native) species except for six ponds that in addition to the warmwater species, are planted annually with rainbow trout from area WDFW hatcheries. The planted lakes include ponds 1, 2, 3, 4, and 6 in the I-82 unit and Powerline Lake, located within the Windmill Ranch Unit. The majority of the fish species within these waters are non-native and have been introduced by managers or anglers or have been migrated into these waters via surface water connections including the Columbia Basin irrigation system and Yakima River. Largemouth bass, yellow perch, bluegill, and black crappie are the predominant species in the majority of these ponds. Angler effort varies by location and is seasonal.

The six wildlife area units providing shoreline access to the public along the Yakima River are the I-82 Ponds, Sunnyside, Byron, Vance-Ferry Road, Whitstran, and Benton City. Anadromous species present in the Yakima River include spring, summer, and fall runs of Chinook salmon, sockeye, coho, steelhead, and Pacific lamprey. ESA-listed fish species present in the Yakima River bordering the wildlife areas include Middle Columbia River steelhead and bull trout. There are seasonal sport fisheries for spring Chinook, fall Chinook, and coho salmon in the Yakima River when returns are in excess of conservation needs or the proportion of hatchery returns are elevated. To reduce impacts of incidental take (catch & release) of both juvenile and adult anadromous species in the Yakima River, angling for gamefish is opened on a seasonal basis.

Fish species

A total of 33 distinct species can be found within the Sunnyside Snake River Wildlife Management Unit: 20 native and 13 non-native species. The dominant species observed in the inland lakes are largemouth bass, yellow perch, black crappie, bluegill, and common carp. Table 12 lists all known species, origin, current status, and the management unit where they have been documented. The Yakima River supports at least 48 species of anadromous, resident native, and non-native fish, but many of these are not present in the lower river bordering the management units.

Common Name	Scientific Name	Origin	Federal/State Status/SGCN	Wildlife Area Unit*
Black Crappie	Pomoxis nigromaculatus	Nonnative		I82,ML,PL,WB, YRU,
Bluegill	Lepomis macrochirus	Nonnative		I82,ML,WB,YRU
Brown Bullhead	Ictalurus nebulosis	Nonnative		182,PL
Bull Trout	Salvelinus confluentus	Native	FT, SC, SGCN	YRU
Bridgelip Sucker	Catostomus columbianus	Native		ML,YRU
Channel Catfish	Ictalurus punctatus	Nonnative		182,YRU
Chinook Salmon spring/summer/ fall	Oncorhynchus tshawytscha	Native		YRU
Chiselmouth	Acrocheilus alutaceus	Native		182,YRU

Table 12: Fish species on the Sunnyside Snake River Wildlife management areas



Common Name	Scientific Name	Origin	Federal/State Status/SGCN	Wildlife Area Unit*
Coho Salmon	Oncorhynchus kisutch	Native		YRU
Common Carp	Cyprinus carpio	Nonnative		ML,YRU
Goldfish	Carassius auratus	Nonnative		ML
Lake Whitefish	Coregonus clupeaformis	Nonnative		ML
Largemouth Bass	Micropterus salmoides	Nonnative		All*
Largescale Sucker	Catostomus macrocheilus	Native		ML,182,YRU
Pacific Lamprey	Entosphenus tridentatus	Native	FSC, SGCN	YRU
Longnose Dace	Rhinichthys cataractae	Native		YRU
Longnose Sucker	Catostomus catostomus	Native		ML
Mountain Sucker	Catostomus platyrhynchus	Native	SC, SGCN	YRU
Mountain Whitefish	Prosopium williamsoni	Native		YRU
Northern Pikeminnow	Ptychocheilus oregonensis	Native		YRU
Peamouth	Mylocheilus caurinus	Native		YRU
Pumpkinseed	Lepomis gibbosus	Nonnative		I82,ML,WB,YRU
Rainbow Trout	Oncorhynchus mykiss	Native		182,PL,YRU
Redside Shiner	Richardsonius balteatus	Native		YRU
Sculpin	Cottus (various species)	Native		ML,YRU
Smallmouth Bass	Micropterus dolomieu	Nonnative		YRU
Sockeye Salmon	Oncorhynchus nerka	Native		YRU
Speckled Dace	Rhinichthys osculus	Native		YRU
Steelhead DPS	Oncorhynchus mykiss	Native		YRU
3-spine Stickleback	Gasterosteus aculeatus	Native		YRU
Walleye	Sander vitreus	Nonnative		ML,YRU
Yellow Bullhead	Ictalurus nebulosis	Nonnative		YRU
Yellow Perch	Perca flavescens	Nonnative		ML,PL, WB, WL,YRU

BoP Bounds Pond BrP Bridgeman PondP5 Byron Ponds ΒP Clark Pond СР Giffin Lake GL HP Haystack Ponds

I-82 Pond 5

ML

I-82 Ponds (1-6) JW

Mesa Lake

MGL Morgan Lake

HL Horseshoe Lake

ΡL Powerline Lake WB WB Wasteway Ponds

Johnson Wetland

WML Windmill Lake

Worth Lake WL

YRU Yakima River units



There are eight focal species/stocks (Table 13) in the Yakima River evaluated by the Yakima Subbasin Fish and Wildlife Planning Board.

Focal Species Criteria	Bull trout	Steelhead/ Rainbow trout	Spring Chinook	Summer Chinook	Fall Chinook	Sockeye	Coho	Lamprey
ESA Status	Threatened	Threatened	None	None	None	None*	None	None
Has Ecological Significance	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Has Cultural Significance	Yes	Yes	Yes	Yes	Yes	Yes	Yes	
Anadromous and/or Resident	R	A & R	A	A	A	А	А	А
* Sockeye were extirpated from the Yakima Subbasin ca 1920								

Table 13: Focal fish species in the Yakima River Basin

Yakima Basin bull trout populations were listed as threatened under the ESA, effective July 10, 1998. Nine distinct bull trout stocks have been identified in the Yakima Basin by WDFW. Of these nine stocks, six are classified as "Critical," one as "Depressed," one as "Healthy," and one as "Unknown".

Spring Chinook salmon populations have been dramatically reduced from pre-1850s abundance levels. Introductions of spring Chinook from the Cle Elum Supplementation and Research Facility (CERSF) have increased the abundance of spawning fish in the Upper Yakima spring Chinook population.

Fall Chinook salmon populations have also been dramatically reduced from pre-1850s abundance levels. There are two genetically distinct stocks recognized in the Yakima Basin. The mainstem stock is found throughout the lower Yakima River (roughly the lower 100 miles), and a stock is endemic to the Marion Drain, a man-made drainage ditch for the Wapato Irrigation Project. Environmental conditions have changed since the early 1930s, which has resulted in decreased production of fall Chinook and a shift in juvenile outmigration shifting to earlier in the year.

Steelhead and rainbow trout are widely but thinly distributed across the Yakima Basin and have been dramatically reduced from pre-1850s abundance levels. Yakima Basin steelhead were listed as threatened under the ESA, effective May 24, 1999. Production of steelhead within the Yakima Basin is heavily weighted towards Satus and Toppenish creeks, which have healthy populations. Anadromy in rainbow trout populations, and the overall size of the population in the Upper Yakima River, is presently much decreased from historic levels.

The historical total run size of Yakima River sockeye salmon has been estimated at either 100,000 or 200,000. Sockeye were extirpated following the completion of impassible storage dams below all natural rearing lakes in the late teens and early 1920s. Because sockeye salmon were extirpated from the Yakima Subbasin so long ago, little is known about genetic or life history variation that may have occurred in individual stocks or populations.

Pacific lamprey, once an important food source for Native Americans in the subbasin, is a Washington State species of concern and is under consideration for ESA listing by USFWS. They are currently found in the mainstem Yakima and Naches rivers, but fewer than 15 have been observed in the Yakima system since 1992.



Fish management

Inland lakes

Fish populations in the lakes located on the wildlife area have been relatively stable long term with normal cyclic and seasonal variations in numbers. This stability in populations requires limited oversight and management. Many of the lakes have surface water connections to the Columbia Basin irrigation system or are subject to inundation from the Yakima River during flooding which allows immigration and emigration of fish species. This connectivity reduces management options for management of or supplementation of these lakes. The WDFW Warmwater Program has regional teams that conduct population assessments of WDFW-controlled lakes. These surveys have been conducted on the I-82 Ponds, Mesa Lake, Thompson Seeps, Worth Lake, and Powerline Lake. These management briefs are located on the WDFW website at: (https://wdfw.wa.gov/publications/). Only a limited number of the lakes located within the wildlife area are suitable for rainbow trout due to depth, water temperature, and connectivity. Most lakes in south central Washington cannot support rainbow trout through the summer months due to lethal water temperatures, with Powerline Lake as the only exception. Rainbow trout have been shown to "carry over" for multiple years producing fish in excess of four pounds. WDFW district biologists allocate rainbow trout plants to each of the six lakes annually based on availability from the hatchery and recommendations from WDFW fish and wildlife staff.

Yakima River

Sockeye, summer Chinook, and coho salmon were extirpated from the Yakima Basin but have been reintroduced. Bull trout is a focal species, but historical abundance estimates are unavailable. In March 25, 1995 (64 FR 14517) the National Marine Fisheries Service listed summer steelhead in the Mid-Columbia ESU, which includes the Yakima Subbasin, as threatened under the ESA. Endemic coho stocks were extirpated by 1980 although naturalized production resulting from hatchery releases have been documented since 1989. Endemic summer Chinook were last observed in the early 1970s and were considered extirpated.

Sockeye were historically abundant, but were extirpated following the completion of impassible storage dams below all natural rearing lakes in the late teens and early 1920s. Pacific Lamprey are a U.S. Fish and Wildlife Service category two candidate species, and in the Yakima Basin, have become very rare. Historically, 500,000-900,000 adult salmon and steelhead returned to the Yakima Subbasin annually. This total was comprised of spring, summer, and fall Chinook, coho, sockeye, and steelhead. Summer Chinook, sockeye, and native coho were extinct in the subbasin until recent reintroduction efforts. The number of returning adults is greatly reduced from historic levels. Over the last ten years, returns of spring Chinook have varied from a low of 645 to highs of 25,000. Fall Chinook returns average 2,000 to 4,000, while coho runs have been in the 1,000 to 2,000 range.

Habitat management

This section provides a description of habitat management activities that occur on the Sunnyside-Snake River Wildlife Area, wetland management, weed management, fire management and history, and habitat restoration.



Wetland management overview

The majority of wetland habitats, from vegetated and shallow moist soil impoundments to deeper open water habitats that occur on the Sunnyside-Snake River Wildlife Area, are the result of the U.S. Bureau of Reclamation's Columbia Basin Irrigation Project. A very small number of wetlands are seasonally influenced by subsurface Yakima River flows and the Byron Unit is part of the effluent management from the Grandview Wastewater Treatment Plant. Wetland management occurs both as a part of normal day-to-day operations on the units as well as through developing larger collaborative wetland projects with partners.

Water management of shallow wetlands to promote wetland vegetation is important for native and migratory birds as well as for providing recreational opportunities for waterfowl hunters and bird watchers. Periodically (typically about every two to three years), shallow wetlands will need some type of management to reduce vegetation cover while still providing habitat for wildlife and opportunities for recreation. Since most of these wetlands have some form of water control structures that control incoming and outgoing water, wetlands can essentially be dried out to allow equipment to safely enter the wetland basin for mowing and spraying. In some instances, fire is used as an initial treatment to remove years of decedent vegetation growth prior to spraying. With careful planning, these wetlands can provide suitable year round habitat and seasonal recreational opportunities.

WDFW is collaborating with partners, including Ducks Unlimited, Yakima County, and Yakama Nation to partially remove or alter a levee near ponds 4 and 5 on the I-82 Ponds Unit, located on the Wapato Reach of the Yakima River. This action will reconnect floodplain and side-channel habitat, which will improve existing wetland habitat and create additional wetlands. Ponds 4 and 5 originated from gravel pits excavated by the Washington State Department of Transportation during the construction of the I-82 Highway through Yakima River floodplains. The lower section of the levee built around the ponds is failing. Removing the failing section or reconfiguring the levee presents opportunity to improve habitat conditions, maintain recreation, and protect other infrastructure. Pond 5 is about 18 acres and the meadow south of the pond is about 17 acres.

Design alternatives under discussion and investigation revolve around breaching or lowering the levee, enhancing a portion of Pond 5, adding swales and depressions to the meadow for additional wetland habitat, and potentially enhancing the upland portions of the meadow. Earthwork actions would be included to restore as much floodplain function as possible given the altered state of the river system and the presence of the highway. No changes are planned for Pond 4 (Draft Wapato Reach Action Plan, 2017).

The Yakima River Basin Integrated Water Resource Management Plan (Integrated Plan), prepared by the Bureau of Reclamation and the Washington State Department of Ecology identified the Wapato Reach as a priority area for ecological restoration and management (Reclamation and Ecology 2011). Integrated Plan.

Future phases include enhancing habitat in the perennial side channel downstream of Pond 5; install riparian plantings, remove left bank rip-rap and levee to reconnect the side channel.

Fire history and management

Fire history

Periodic fires, both human-caused and natural (lightening) have and continue to affect all habitat types in the Sunnyside-Snake River Wildlife Area. Data from 2000 to present indicate that 55% of the fires were human caused and only 22% were from lightening (Table 14).



Fire Name	WLA Unit	Year	Cause	Acres Burned/Type
Duck Club Fire	Sunnyside	2014	Human	250
April Fire	Sunnyside	2014	Human	200
N/A	Mesa Lake	2015	Human	63
N/A	I-82	2000	Lightening	100
N/A	Sunnyside	2000	Controlled burn	unknown
N/A	Thornton	2000	Human	900 / shrubsteppe
N/A	Thornton	2003	Lightening	1,200 / shrubsteppe
N/A	Rattlesnake Slope	2018	Target Shooting	185
N/A	Rattlesnake Slope	2018	Electrical	5,080

Table 14: Fires history – fires greater than 50 acres from 2000 - 2018

Prescribed fires are used as one of many land management tools to control vegetation and promote plant growth. On the Sunnyside-Snake Wildlife Area prescribed fire is primarily used to remove excessive vegetation litter buildup in wetlands. This opens up vegetation areas and can provide efficient delivery and effectiveness of herbicides that may be used post fire to control resprouting vegetation or to provide more open water habitats. Annually, one or two wetlands typically get burned at the Sunnyside Unit, depending on environmental conditions and personnel availability. Local fire districts assist with conducting prescribed burns and all prescribed fires comply with local and state regulations. Additional prescribed burns are being planned for the Thompson Seeps, Windmill Ranch, and Mesa Lake units.

Fire management

Wildland fires ignited on or in the vicinity of the Sunnyside-Snake River Wildlife Area are responded to by county fire districts and DNR. Several units of the wildlife area that are nearer to urban centers (Benton City, Prosser) and are part of fire protection districts/associations, but more rural units have no formal fire protection coverage. USFS fire crews also provide protection primarily in areas of federal ownership and DNR has fires crews throughout the area that may also provide fire resources. In addition, wildlife area staff maintain fire suppression qualifications and make equipment accessible for controlling wildfire when it is needed. Wildlife area staff coordinate with Fire Districts, DNR, and USFS as resource advisors and landowner representatives to minimize habitat loss, protect resources, and meet fire suppression needs.

Weed management

Managing weeds is a significant part of the staff's workload to establish and maintain diverse native plant communities that support fish and wildlife populations. Weed management must also meet legal obligations and reduce the likelihood of spread of noxious weeds to adjacent private lands. Invasive plants and noxious weeds can infest high quality native plant communities and convert them to low quality monocultures that reduce wildlife value. The weed management plan (see Appendix B) identifies species and management practices to control weeds. Weeds of primary concern on the wildlife area include: purple loosestrife, poison hemlock, kochia, thistles, knapweeds, puncturevine, rush skeletonweed, and phragmites. WDFW will continue coordinating with county and the Yakama Nation on purple loosestrife control. As part of noxious



weed mandate, when conducting planned projects, preference is given to replacing pollen and nectar rich noxious weeds with native pollinator-friendly forage plants when appropriate to meet management goals.

Climate change approach

Purpose

The purpose of this section is to evaluate how projected changes in climate will affect the resources of the Sunnyside-Snake River Wildlife Area and to highlight opportunities to mitigate or prepare for those effects. This section also summarizes work by the wildlife area planning team to review the management objectives (see Goals and Objectives section) take into consideration factors to ensure that objectives are robust to future changes.

This work is consistent with the directives of a 2017 WDFW policy titled "Addressing the Risks of Climate Change", which states that WDFW will "manage its operations and assets so as to better understand, mitigate, and adapt to impacts of climate change".

Projected climate change impacts

Increasing greenhouse gases will lead to warmer temperatures throughout this century for the Pacific Northwest. The most direct impacts of climate change to this area will be in the form of warmer winters (3-6 degrees within 15 years) and dryer summers (Climate Impacts Group 2013). For summer months, a majority of models projected decreases in precipitation, with the average declining 16% by the 2080s. A majority of models projected increases in winter precipitation, with an average value reaching over 9% by 2080 (Mote and Salathé 2009). Other key impacts are highlighted below.

Forests in the northwest also will likely be affected by climate-driven changes in disturbance regimes, such as wildfire (Littell et al. 2010), insect outbreaks (e.g., mountain pine beetle; Logan et al. 2003), disease (e.g., Swiss needle cast; Black et al. 2010), and drought (van Mantgem et al. 2009; Knutson and Pyke 2008). The areas burned by fire in the Columbia River Basin is projected to triple by the 2040s relative to median for 1916-2006 (Littell et al. 2010, 2012). Wildfire suppression costs have increased as fire seasons have grown longer and the frequency, size, and severity of wildfires has increased due to changing climatic conditions, drought, hazardous fuel buildups, insect and disease infestations, nonnative invasive species, and other factors. Funding has not kept pace with the cost of fighting fire. Over the last 10 years, adjusting for inflation, the USFS has spent an average of almost \$1.13 billion on suppression operations annually.

Vegetation models of sagebrush-steppe systems in eastern Washington and Oregon simulate large declines in current distributions of shrublands under future climate conditions (Neilson et al. 2005; Rogers et al. 2011), with shrubs largely replaced by woodland and forest vegetation. The response to climate change of grassland and shrubland systems throughout the northwest will be influenced by invasive species that are currently present in these systems or may be able to expand into these systems as climate changes (Dennehy et al. 2011).

Impacts to wildlife area resources and potential adaptation

The wetlands in the Sunnyside-Snake River Wildlife Area are dependent on irrigation water, and for this reason may be more resilient than other "natural" wetlands in the region, given the shifts in precipitation and hydrologic regimes expected with climate change. It also means that the ecological significance of these



wetlands could potentially become more important over time, if wetlands dependent on natural seeps and springs become less productive. This concept could be further developed and potentially used to secure funding or other enhancements. Agricultural practices are already changing to become more water efficient, which results in less irrigation water for the wetlands in the wildlife area. Climate change may force additional efficiencies and may impact the water availability for habitat needs. Similarly, crop suitability may change with increasing drought, heat, and wildfires.

Monitoring changes in species and habitat composition, as well as changes in how species are using habitat (including the timing of key ecological events, such as bud burst, first and last frosts, and migration) are critical tools for understanding how and when climate is impacting resources on the wildlife areas. There is a potential role for a citizen science effort on this and other wildlife areas to help monitor ecological indicators.

The climate nexus identified for several of the objectives points to an overall context of changing future conditions – away from historic natural variability. Prioritization of where to restore or develop new wetlands, or improve upland habitat, should be done from an understanding of how future conditions may change. For example, the UW Climate Impacts Group recently conducted a study on climate change impacts on Columbia Basin wetlands, and studies such as these should be consulted as new initiatives are developed.

Species of concern with high vulnerability to climate change

Eight Species of Greatest Conservation Need (SGCN) potentially on the Sunnyside-Snake River Wildlife Area have been ranked by the climate vulnerability assessment to have a moderate-high vulnerability to climate change, and with high confidence in the data. These include: black-tailed jackrabbit, white-tailed jackrabbit, Townsend's ground squirrel, Washington ground squirrel, Townsend's big-eared bat, golden eagle, and greater sage grouse. None of these species are on the WDFW "preliminary climate watch list". Only SGCN were considered in this assessment and it does not include climate sensitivities for other species that may be associated with the wildlife area. See Appendix A for terrestrial SCGN and relationship to ecological systems of concern.

Species of greatest conservation need	Population size/trend	Conservation concern
Black-tailed jackrabbit	Low/declining	Once abundant and broadly distributed in eastern Washington, the species is now rare and sparsely distributed due to habitat loss from fragmentation and possibly disease.
White-tailed jackrabbit	Low/declining	Once abundant and broadly distributed across the bunchgrass communities of eastern Washington, the species is now rare and sparsely distributed due to the loss, degradation, and fragmentation of habitat and possibly disease and competition with black-tailed jackrabbits
Townsend's ground squirrel	Unknown/ unknown	Population status unclear. Significant declines have occurred in many areas, yet this species is common at a number of human-modified locations.

Table 15: Species of greatest conservation need with high to moderate vulnerability and high confidence



Species of greatest conservation need	Population size/trend	Conservation concern
Washington ground squirrel	Low/declining	Associated with shrubsteppe and steppe in eastern Washington and is threatened by a number of factors, especially habitat loss, degradation, and fragmentation.
Townsend's big-eared bat	Low/stable	Occurs in small to moderately-sized aggregations at sites throughout the state, where it may be vulnerable to human disturbance during the breeding and wintering periods.
Golden eagle	Low/unknown	Species is of concern due to declines in the distribution and abundance of its primary prey species, jackrabbits and ground squirrels; across its range additional mortality factors include continued exposure to lead in the environment and collisions at wind energy facilities.
Greater sage grouse	Low/stable	Requires large landscapes of sagebrush steppe, much of which has been degraded, fragmented, or lost. The primary threat is the combined impact of habitat loss, fragmentation, and degradation.

Shrubsteppe habitat and species associations

Over 10,000 acres of the Sunnyside-Snake River Wildlife Area are in a shrubsteppe system called "Intermountain Basins Big Sagebrush Steppe", which is classified as imperiled (see Table 10 - Ecological systems of concern). Eight species, which are also species of greatest conservation need, are closely associated with this system. These species are American badger, burrowing owl, ferruginous hawk, greater sage grouse, sage thrasher, sagebrush sparrow, Woodhouse's toad, and the sagebrush lizard. WDFW and EcoAdapts's 2016 fact sheet "How Will Climate Change Affect Shrub-Steppe Ecological Systems and Species in Washington?" is

summarized in Table 16 for species that may occur on the Sunnyside-Snake River Wildlife Area.

Shrubsteppe systems are sensitive to changes in precipitation and soil moisture, temperature, drought, and altered wildfire regimes. Changes in precipitation can lead to shifts in species composition or vegetation structure. More frequent fire could result in conversion to annual grasslands, which would be adversely impact many species. Shrubsteppe habitats and species will likely be adversely affected under projected future climate conditions. The level of certainty is high that the summers will get drier and hotter in the Columbia Plateau. Fall, wing, and spring will be wetter and also



Burrowing owls Photo by Doug Kuehn

hotter. Inter-mountain Basins Big Sagebrush Steppe is projected to decline by the end of the century. About 4% is projected to remain stable, and 70% to become climatically unsuitable.



SGCN species closely Vulnerability **Actions needed** Threats associated with rank shrubsteppe Requires large landscapes of sagebrush steppe, much of which High sensitivity / Protect and restore key has been degraded, fragmented, or Greater sage moderate habitat. Replant sagebrush. lost. The primary threat is the grouse confidence Augment populations. combined impact of habitat loss, fragmentation, and degradation. Low sensitivity -Increased temperatures and Protect and restore key **Burrowing owl** moderate changes in precipitation may lead to habitat. Conduct surveys to range contractions confidence assess status and trends. Low sensitivity -Warmer temperatures may benefit Protect and restore Ferruginous moderate if grasslands expand, however shrubsteppe habitat. Protect hawk confidence droughts may lead to decline in prey nests from disturbances. Conduct research and modeling of habitat using High sensitivity / Reductions in shrubsteppe findings of habitat American badger moderate grasslands and riparian areas could associations from badger confidence lead to altered prey availability surveys. Restore populations of prey species. Moderate - high Reductions in sagebrush habitat will sensitivity / Protect sagebrush habitat. Sage thrasher impact forage, nesting and moderate Control cheatgrass. reproduction confidence Protect core areas of habitat; identify degraded Moderate – high Reductions in sagebrush habitat will habitat for restoration and Sagebrush sensitivity / low impact forage, nesting and sparrow establish connectivity confidence reproduction between core areas. Control cheatgrass. Research, surveys and monitoring to understand Moderate - high Reductions in shrubsteppe species distribution and Woodhouse's sensitivity / grasslands and riparian areas could status. Protect native toad moderate cause adverse impact shrubsteppe habitat from confidence conversion and degradation due to agriculture. Monitor populations to Moderate - high Reductions in sagebrush habitat will make sure their habitat Sagebrush lizard sensitivity / low impact forage, nesting and remains suitable and the confidence reproduction populations persist. Protect sand dune habitat.





Making the goals and objectives of the wildlife area plan climate resilient

The information listed in Table 17 is a list of the Sunnyside-Snake River Wildlife Area goals and objectives potentially affected by climate change, or those with a "climate nexus". Actions and considerations are listed to ensure climate impacts are addressed in implementation of the wildlife area management plan. Opportunities are summarized below for the list of objectives on page 90

Objective	Climate Nexus				
GOAL: Maintain or improve the ecolog	gical integrity of priority sites.				
Establish an ecological integrity baseline and associated goals for ecological systems of concern/priority systems by 2024.	 Include climate change future conditions in planning and monitoring. Develop indicators such as water and air temperature. Measure what is most sensitive to climate change. Use climate tools to evaluate goals and objectives. Evaluate for attainability in light of climate change and maximize outcomes. 				
Implement the 10-year plan for wetland management and restoration. Increase wetland function and value through vegetative restoration.	 Use climate change trends as a criteria for prioritizing where to restore or create new wetlands. Monitor ground water and analyze for trends. Consider how other ecologically important wetlands in region may be affected by climate change – importance of these managed wetlands may increase with a longer term perspective. 				
Implement weed management plan annually.	 Consider monitoring for invasive species expected to increase under climate change. Plan for possibility of new weeds. 				
Develop protocol on how to manage nuisance beavers on the wildlife area.	 Regionally, beavers may be an important resilience strategy on some landscapes. Evaluate effort to relocate beavers in the larger context. 				
Identify upland areas with high potential for habitat improvement and develop action items.	Upland habitat is important for climate refugia.				
Maintain and restore floodplain functions.	 Consider impacts of future flow regimes and extreme weather events in prioritizing locations and in project design. 				
GOAL: Achieve species diversity at levels consistent with healthy ecosystems.					
Conduct survey for Species of Greatest Conservation Need in coordination with the Diversity Division.	 May need to adjust surveys to accommodate species range shifts and/or phenological shifts. Adjust survey timing to match species instead of static window each year. 				
Address habitat loss from recent fires on Rattlesnake Slope.	Consider climate change in restoration strategy.				

Table 17: Plan objectives with climate nexus



Objective	Climate Nexus					
Identify opportunities to enhance monarch butterfly, bumble bee, and other pollinator habitat.	•	If promoting new plantings, prioritize locations based on suitable current and future climatic conditions.				
Develop wetland habitat in Thompson Seeps Unit.	•	Habitat goals should be designed around future conditions due to climate impacts.				
GOAL: Support and maintain appropriate recreation opportunities						
Improve recreational experience user expectations, and support of the wildlife area by providing information on the web, at kiosks, in maps, and directional signage.	•	As new signage and interpretive material is developed, keep in mind opportunities to include climate change, which can increase the knowledge of the ecological importance of these wetlands.				
Assess potential to improve lakes and stock with fish (trout and/or warm water)	•	Habitat suitability for certain species may change.				
Manage and improve upland bird habitat and nesting cover.	•	Consider fire impacts/potential in habitat projects.				
Identify opportunities to improve/increase waterfowl hunting opportunities.	•	Criteria should include consideration of any future climatic changes, water temperatures, groundwater or surface hydrology.				
GOAL: Identity O&M funding sources.						
Investigate and identify resources for additional O&M funding.	•	Consider value of these wetlands in the context of climate change, as a selling point for grants. Use species vulnerability assessment results to help demonstrate the value of wildlife area to species – help them to adapt to climate change (consistent water availability, etc)				



References

Climate Impacts Group. 2013. Washington State of Knowledge Report – Climate Change Impacts and Adaptation to Washington State: Technical Summaries for Decision Makers, Climate Impacts Group. Seattle, WA.

Governor's Office of Indian Affairs. Treaty with the Yakama, 1855. <u>https://goia.wa.gov/tribal-government/treaty-yakama-1855</u>

Lenfesty, C.D. and T.E. Reedy. 1985. Soil Survey of Yakima County Area, Washington. United States Department of Agriculture, Soil Conservation Service. Washington D.C.

ICF International, in association with R2 Consultants. 2012. Wapato Reach Assessment Report. April 26. (ICF 00703.11.) Seattle, WA. Prepared for Yakama Nation, Toppenish, WA. Washington Department of Fish and Wildlife. 2015. Washington's State Wildlife Action Plan: 2015 Update. Washington Department of Fish and Wildlife, Olympia, Washington.

Mote and Salathé. 2009. Future Climate in the Pacific Northwest. Chapter 1 in: The Washington Climate Change Impacts Assessment: Evaluating Washington's Future in a Changing Climate. Seattle, WA

Sandisen, Derek and Christiansen, Wendy, et al. 2012. Yakima River Basin Integrated Water Resource Management Plan Final Programmatic Environmental Impact Statement

Schroeder, Michael A., A.J. Shirk, A. Well, L.A. Robb. 2015. GNLCC Final Report. Habitat Occupancy and Movement by Greater Sage-Grouse in Washington State.

Wapato Action Plan Steering Committee, Yakima River Basin Water Resource Management Plan Workgroup. 2017. Draft Wapato Reach Action Plan.

U.S. Fish and Wildlife Service. 2015. Recovery plan for the coterminous United States population of bull trout (*Salvelinus confluentus*). Portland, Oregon. xii + 179 pages.

Washington Department of Fish and Wildlife. 2015. Washington's State Wildlife Action Plan: 2015 Update. Washington Department of Fish and Wildlife, Olympia, Washington.

Washington Department of Fish and Wildlife and EcoAdapt. 2016. How Will Climate Change Affect Shrub-Steppe Ecological Systems and Species in Washington? <u>http://ecoadapt.org/data/documents/EcoAdapt WDFW_SHRUB-STEPPE_Final.pdf</u>

Western Regional Climate Center. Climate of Washington. <u>https://wrcc.dri.edu/Climate/narrative_wa.php</u>

Yakama Nation. 2012. Wapato Reach Assessment Report. Prepared by ICF International.



Appendices

- A. Species and habitat information
- B. Weed management plan
- C. Fire response information
- D. Wetlands restoration summary/plan
- E. Cultural resources summary
- F. Public response summary



Appendix A. Species and habitat information

Terrestrial SGCN Relationship with Ecological Systems of Concern

SGCN Relationship with Ecological Systems of Concern	Columbia Basin Foothill and Canyon Dry	Columbia Basin Foothill Riparian Woodland and	N Rocky Mtn Lower Montane Riparian Woodland	Columbia Plateau Steppe and Grassland	Inter-Mountain Basins Big Sagebrush Steppe	Inter-Mountain Basins Greasewood Flat	Inter-Mountain Basins Semi-Desert Shrub Steppe	North American Arid West Emergent Marsh
Black-tailed jackrabbit				√	√			
White-tailed jackrabbit	✓			~	✓			
Merriam's shrew	✓			~	✓			
Hoary bat	✓	~	√	✓	✓	✓		✓
Townsend's big-eared bat	✓	\checkmark	√	~	✓		√	✓
Spotted bat		√		\checkmark	√	\checkmark		√
American badger	✓			~	✓	\checkmark		
Townsend's ground squirrel				√	✓			
WA ground squirrel	✓			✓	\checkmark		✓	
American white pelican								✓
Western grebe								
Greater sage grouse				\checkmark	✓	\checkmark	\checkmark	
Ferruginous hawk	✓	\checkmark		\checkmark	\checkmark		\checkmark	
Golden eagle	✓	~	\checkmark	\checkmark	✓	\checkmark	\checkmark	
Bald eagle	✓	\checkmark	\checkmark					\checkmark
Peregrine falcon	✓		✓		✓	\checkmark		✓
Cinnamon teal				√	√		√	√
Sage thrasher				√	√			
Sagebrush sparrow				√	✓			
Burrowing owl	✓			√	✓	√	✓	
Western screech owl								
Mountain quail	✓		√	√	√			
Loggerhead shrike	✓	√		√	√	√	√	
Tiger salamander	✓		√	√	√			√
Woodhouse's toad				✓	✓			✓
Desert nightsnake	✓			\checkmark	✓			
Sagebrush lizard					✓			
Pygmy horned lizard				\checkmark	✓			
Columbia clubtail dragonfly		✓						
White-belted rintail dragonfly								
Columbia River Tiger Beetle								
Western Bumble Bee								
Morrison's Bumble Bee	√	√	✓	√	\checkmark		√	
Suckley Cuckoo Bumble Bee								
Shortface Lanx								



Priority habitats in Yakima, Benton, and Franklin counties Source: WDFW Priority Habitats and Species, 2017

Yakima	Benton	Franklin		
Aspen Stands	Aspen Stands	Aspen Stands		
Biodiversity Areas & Corridors	Biodiversity Areas & Corridors	Biodiversity Areas & Corridors		
Inland Dunes	Inland Dunes	Inland Dunes		
Old-Growth/Mature Forest				
Eastside Steppe		Eastside Steppe		
Oregon White Oak Woodlands				
Riparian	Riparian	Riparian		
Shrubsteppe	Shrubsteppe	Shrubsteppe		
Freshwater Wetlands & Fresh	Freshwater Wetlands & Fresh	Freshwater Wetlands & Fresh		
Deepwater	Deepwater	Deepwater		
Instream	Instream	Instream		
Caves	Caves	Caves		
Cliffs	Cliffs	Cliffs		
Snags and Logs	Snags and Logs	Snags and Logs		
Talus	Talus	Talus		



Appendix B. Weed management plan

Weed control goals at Sunnyside-Snake River Wildlife Area

The goals of weed control on Department lands at the Sunnyside-Snake River Wildlife Area, which is composed of Bailie, Benton City, Byron, Esquatzel Coulee, Glover, Sunnyside, Hope Valley, I-82 Ponds, Mesa Lake, Rattlesnake Slope, Thornton, Vance-Ferry Road, Thompson Seeps, Whitstran, and Windmill Ranch units, are to maintain or improve the habitat for fish and wildlife, meet legal obligations, and protect adjacent private lands.

WDFW focuses land management activities on the desired plant species and communities, rather than on simply eliminating weeds. Control for listed species is mandated by state law (RCW 17.10 and 17.26) and enforced by the County Noxious Weed Board. WDFW strives to meet its legal obligation to control for noxious weeds listed according to state law (Class A, B-Designate, and county listed weeds). Importantly, the wildlife area will continue to be a good neighbor and partner regarding weed control issues on agency and adjacent lands. The agency believes the best way to gain long-term control is to work cooperatively on a regional scale. As funding and mutual management objectives allow, WDFW will find solutions to collective weed control problems.

WDFW uses integrated pest management (IPM), which is defined in RCW 17.15.010 as "a coordinated decision-making and action process that uses the most appropriate pest control methods and strategy in an environmentally and economically sound manner to meet agency programmatic pest management objectives."

WDFW's weed management objectives

Staff will formulate a plan for the upcoming season. Prior to the beginning of each field season, wildlife area staff will review the previous year's weed control efforts. They will address any new issues and identify focal areas.

Wildlife area staff will routinely identify areas of upland habitat for restoration improvements and increased nesting and brood rearing cover. These areas will be evaluated for weed control efforts necessary for successful restoration efforts.

Survey up to 500 acres annually on <u>Rattlesnake Slope</u> to determine post-fire weed control requirements in order to maintain and enhance forage for elk utilizing the area. Reseed any areas burned by wildfire in the first winter following the fire. Use native seed mixes (local ecotypes when possible) that include grasses, forbs and shrubs. Use techniques proven to enhance and speed recovery from fires before competitive weeds invade.

Reseed non-irrigated fields at the <u>Sunnyside Unit</u> to more native grass, forb and shrub species, as additional funding allows, helping recover native habitats for native wildlife.

Plant scattered blocks of sagebrush on <u>Rattlesnake Slope</u> grasslands to provide nursery areas for natural reproduction of shrub cover.

Check <u>Windmill Ranch</u>, <u>Sunnyside</u>, and <u>Mesa Lake</u> roads and treat annually for maintenance needs. Besides general weeds, species such as kochia and Canada thistle area problematic and pose a risk of spreading to new areas if not treated and controlled. These sites require annual maintenance.



Check wetland and riparian areas annually, especially in areas managed for waterfowl, for purple loosestrife, reed phragmites and Russian olive. Remove 40 acres of Russian olive at the Sunnyside Unit each year until 2023 to restore additional wetlands. Control Canada thistle, poison hemlock and Russian knapweed in river oxbows and swales, and plant to native species.

Staff will take all necessary precautions to avoid incidental treatment of plant species important for pollinators. Weed control efforts will be evaluated against the benefit of wildflowers and forbs on a site by site basis.

Staff will report weed control efforts annually to the Agency's Statewide Noxious Weed Coordinator.

Weed species of concern on the wildlife area include but are not limited to:

Bristly foxtail (*Hordeum jubatum*), Canada thistle (*Cirsium arvense*), cheatgrass (*Bromus tectorum*), cocklebur (*Xanthium spinosum*), diffuse knapweed (*Centaurea diffusa*), hairy whitetop (*Cardaria pubescens*), houndstongue (*Cynoglossum officinale*), jointed goatgrass (*Aegilops cylindrical*), kochia (*Bassia scoparia*), poison hemlock (*Conium maculatum*), puncturevine (*Tribulus terrestris*), purple loosestrife (*Lythrum salicaria*), reed phragmites (*Phalaris australis*), rush skeletonweed (*Chondrilla juncea*), Russian knapweed (*Rhaponticum repens*), Russian olive (*Elaegnus angustifolia*), Russian thistle (*Salsola spp.*), spotted knapweed (*Centaurea stoebe*), white bryony (*Bryonia alba*) and yellow starthistle (*Centaurea solstitialis*).

Weeds occurring on the wildlife area and associated units are listed in Table 1. The table also describes the weed's classification, an estimate of the acreage affected by the weed, how many acres were treated, the relative density of infestation, the general trend the weed infestation has been exhibiting, the control objective and/or strategy for the weed and finally, which wildlife units have the weed present.

Detailed descriptions and natural history information for each of the above state-listed weed species listed above can be found at the Washington State Noxious Weed Control Board web site:<u>http://www.nwcb.wa.gov/search.asp</u>. Information on other species contained in the list can be found at the University of California's IPM Online web site: <u>http://www.ipm.ucdavis.edu/PMG/weeds intro.html</u>.

Weed management information for individual weed species can be found at the PNW Weed Management Handbook link at: <u>http://pnwhandbooks.org/weed/control-problem-weeds</u> and on WDFW's weed management website at: TBD.

Resources

Detailed descriptions and natural history information for each of the above state-listed weed species listed above can be found at the Washington State Noxious Weed Control Board web site at http://www.nwcb.wa.gov/search.asp.

Information on other species contained in the list can be found at the University of California's IPM Online web site at <u>http://www.ipm.ucdavis.edu/PMG/weeds intro.html</u>.

Weed management information for individual weed species can be found at the PNW Weed Management Handbook link at <u>http://pnwhandbooks.org/weed/control-problem-weeds</u>


Table 1: Sunnyside-Snake River Wildlife Area weed table including the weed class and unit location on the wildlife area.

Weed Species	Benton	Franklin	Walla Walla	Yakima	2016 Estimated Affected Acres	2016 Treated Acres	Annual Trend	Control Objective/ Strategy	Wildlife Area Unit Weed Distribution
Bristly foxtail		NA			60	8	No Change	Control	Windmill Ranch (2013)
Broadleaf weeds		NA			134	125	Decreasing	Control	Bailie Youth Ranch, Hope Valley, Mesa Lake, Windmill Ranch
Canada thistle		C			41	36	Decreasing	Control	Hope Valley (biocontrol 2013), Mesa Lake, Windmill Ranch
Cheatgrass		NA			24	24	Decreasing	Control	Mesa Lake, Windmill Ranch
Cocklebur		С			7	7	Decreasing	Control	Mesa Lake, Windmill Ranch
Diffuse knapweed		В			3	0.5	Decreasing	Control	Windmill Ranch (2013), Mesa Lake (biocontrol 2013)
General Weeds	NA	NA		NA	874	210	No Change	Control	Sunnyside, I-82, Mesa Lake, Rattlesnake Slope, Windmill Ranch
Hairy whitetop			C		60	20	No Change	Control	Sunnyside
Houndstongue			В		30	1	No Change	Control	Windmill Ranch (2013)
Jointed goatgrass		С			<0.1	<0.1	Unknown	Eradicate	Mesa Lake (2013)
Kochia		В		В	392.5	94	Decreasing	Control	Hope Valley, Mesa Lake, Windmill Ranch, Sunnyside
Poison hemlock		В			18	16	Decreasing	Control	Windmill Ranch
Puncturevine		В			1	1	Decreasing	Control	Bailie (2013)
Purple loosestrife				В		10	Unknown - Decreasing	Control	Mesa Lake, Windmill, Hope Valley, Bailie (biocontrol 2013), Sunnyside and Byron (biocontrol 2015)

Weed Species	Benton Franklin	Walla Walla	Yakima	2016 Estimated Affected Acres	2016 Treated Acres	Annual Trend	Control Objective/ Strategy	Wildlife Area Unit Weed Distribution
Reed phragmites	C			30	10	Increasing	Control	Mesa Lake, Windmill Ranch
Rush skeletonweed	В			4	4	Decreasing	Control	Hope Valley (2014)
Russian knapweed			В	80	10	Decreasing	Control	Byron
Russian olive			С	50	8	No Change	Control	Sunnyside
Russian thistle	NA			12	10	Decreasing	Control	Hope Valley, Windmill Ranch
Spotted knapweed	В			1	1	Decreasing	Controls	Windmill Ranch (2013)
White bryony	В			0.5	<0.1	Increasing	Eradicate	Windmill Ranch (2013)
Yellow starthistle	В			3	3	Decreasing	Control	Bailie, Windmill Ranch (2014)



Appendix C. Fire response information

Agency	Units Covered	Phone Number
Yakima County FD 5	I-82 Ponds, Glover, Sunnyside, Vance-Ferry Road, Byron	(509) 829-5111
Prosser FD 3 /West Benton Fire Rescue	Thornton, Whitstran	(509) 786-3873
Benton City / Benton County FD 2	Benton City, Rattlesnake Slope	(509)588-3212
Basin City / Franklin County FD 4	Bailie, Thompson Seeps	
Connell / Franklin County FD 1	Hope Valley, Mesa Lake, Windmill Ranch, Esquatzel Coulee	(509) 234-5451
Franklin County - FD 3	Hope Valley	(509) 547-0961

Department of Fish and Wildlife contacts - contact in order listed

Contact	Phone Number
Pat Kaelber, Wildlife Area Manager	(509) 545-2028
Ross Huffman, Region 3 Lands Operations Manager	(509) 457-9313
Scott McCorquodale, Regional Wildlife Program Manager	(509) 575-2740

Fire district

The Sunnyside-Snake River Wildlife area is covered by six Fire Districts: Yakima County FD 5, Prosser FD 3, Benton City, Franklin County FD 4 (Basin City), FD 1 (Connell), and FD 3 (See Maps 20 and 21). There is some overlap on the Rattlesnake Slope Unit (Benton City FD) and Thornton Unit (Prosser FD 3) between the Department of Natural Resources (DNR) and the fire districts. The wildlife area is with the DNR Southeast Region (509) 547-0961.



Figure 20: Fire District Boundaries - West





Figure 21: Fire District Boundaries - East



Washington Department of Fish and Wildlife

Appendix D. Sunnyside-Snake River Wildlife Area Wetland Management Plan

Introduction

This wetland plan is based on current project area conditions and reflects common goals resulting from discussions between wildlife area staff, district biologists, and comments from the public. An adaptive plan is required to address the dynamic nature of the Sunnyside / Snake River Wildlife Area wetland systems, and continued updates will be necessary to maintain the value of this document. The purpose of the Sunnyside / Snake River Wildlife Area Wetland Management Plan is to provide guidance to field staff to help maintain and improve wetland habitat and recreational value throughout the next ten years. This plan supports the mission of Washington Department of Fish and Wildlife, provides a strategic framework for continued management efforts and aligns with the rest of the goals and objectives of the ten-year management plan.

Part I: Wetland Management

Wetland Types

Wetlands of the Sunnyside / Snake River Wildlife Area are classified under three main wetland types – riverine, lacustrine, and palustrine. Riverine wetlands occur in a river channel and lack trees, shrubs and emergent vegetation. Riverine wetland habitat on the wildlife area is found on units along the Yakima River – I-82 Ponds, Sunnyside, Byron, Whitstran, and Benton City. Lacustrine wetlands are characterized by topographic depressions or old river channels, are greater than eight hectares in size, and lack emergent vegetation (open water) on more that 30% of the surface area. Lacustrine wetland habitat is found on the Sunnyside, Bailie, Hope Valley, and Mesa Lake units. Palustrine wetlands have emergent vegetation, are less than 8 hectares in size, are less than two meters deep, and lack wave-formed shorelines. Palustrine wetland habitat is found on all units except Rattlesnake Slope and Thornton.

The Sunnyside Valley Irrigation Canal was completed in 1923. The Yakima Project built six reservoirs in the Yakima River headwaters by 1933. The Columbia Basin Irrigation Project was developed following the completion of the Grand Coulee Dam in 1942. Most of the wetlands in the wildlife area are enhanced by irrigation water delivery. All wetlands associated within the wildlife area are impacted by the various irrigation district operating-regimes. This creates an unknown level of uncertainty for wetland planning because factors such as timing, rates, levels, and unanticipated shut-offs are outside of the control of WDFW managers. The Columbia Basin and Yakima Valley had far fewer wetlands prior to the establishment of irrigation. The original wetlands functioned naturally by recharging during fall and winter rains and spring snowmelt and gradually receding during the hot, dry summer. Natural wetlands are dynamic and typically experience wet-dry cycles. This periodic drying is critical for maintaining wetland productivity and habitat value. The irrigation-influenced wetlands within the wildlife area have a relatively consistent hydro-period throughout the growing season (and entire year in some locations). This consistency can compromise wetland function and accelerate issues such as invasive species, sedimentation, and succession.

Ecological Integrity

Many of the wetlands throughout the wildlife area are not natural. Their existence is a direct result of irrigation development throughout the Columbia Basin and the Yakima Valley. The network of designed irrigation canals and ditches either supply water directly to wetlands, or seepage from these waterways



moves to nearby low areas creating wetland habitat. These wetlands provide habitat to a wide variety of wildlife and offer outdoor recreational opportunities. However, they are not a priority for ecological monitoring at this time.

The Yakima Basin Integrated Plan is a product of multiple City, County, State, Federal, and Non-Governmental entities working together for a common goal of water and resource protection. The Yakima River corridor, and in particular that stretch adjacent to the wildlife area units (Sunnyside and I-82 Ponds), is of considerable importance. The Integrated Plan has identified this stretch of the Yakima River for protection of existing habitat and improving natural hydrology. Wildlife area staff will work with representatives of the Integrated Plan to assure wildlife area needs are met, and habitat conditions are improved where possible.

<u>Risk Management</u>

The greatest risk to any wetland is the availability of water. As many of the wetlands across the wildlife area receive water from irrigation sources, they are at less risk than other wetlands in other regions. However, as water conservation becomes more and more promoted, and irrigation efficiencies continue to improve, the "excess" water that has previously been supplied to the wildlife area wetlands will likely decrease. As these decreases become reality, wildlife area staff will have to assess management decisions to provide the most habitat and wildlife benefits throughout the year, and from year to year.

The natural wear and tear of time on infrastructure is also an important risk to consider. All infrastructure has a limited lifespan. Many water control structures across the wildlife area are near, or past, that intended lifespan. Wildlife area management staff must evaluate all of the existing infrastructure and develop a replacement schedule based on critical need. Replacement designs need to support anticipated future water regimes and potential management activities, and adequate funding to replace and maintain the infrastructure needs to be secured.

Recreation

The wetlands and wetland habitat of the wildlife area supports a variety of outdoor recreational opportunities. The most participated activity is waterfowl hunting. Over 1,000 hunter visits occur each year. Wetland habitat provides travel corridors and cover for mule deer and elk, along with other small game species. Larger wetlands, ponds, and lakes support populations of fish for anglers. The plethora of wildlife also invite individuals out for wildlife viewing. Over 200 species of birds are known to utilize the wildlife area. Many species of wetland plants are considered important to Native Peoples for collection and harvesting.

All of these considerations pose issues to wildlife area management staff. Wetlands need to periodically dry in order to maintain their overall productivity. Historically these dry periods would occur during the summer. These dry periods allow annual plants to grow and produce seeds, which are important food sources during the fall and winter. However, water regimes of irrigation districts typically produce the highest flows during the hottest, driest stretches of summer. These summer water regimes support breeding populations of wildlife that may not otherwise stay in the region. Summer water regimes also support unusually higher populations of mosquitos that have been found to carry West Nile Virus, thus putting visitors at risk.

Management actions that provide the most benefit to one form of recreation don't necessarily benefit other recreation activities. Wildlife area management staff must evaluate all wetlands on an annual basis to



determine which management activities will be implemented for each wetland. Not all wetlands will support all recreational activities every year, but the recreational opportunities will still be available each year.

Part II: Units

<u>Sunnyside Unit</u>

The Snipes Reserve portion of the unit is closed to the public and provides nesting and brood rearing habitat for waterfowl during the spring and summer, as well as resting and loafing sanctuary during the fall hunting season. Within the confines of the Snipes Reserve are Bound's Lake and Horseshoe Pond. Part of Bos Lake extends onto the wildlife area from adjacent private property.

On the east end of the unit is the Johnson Wetland and pond. This wetland has potential to provide nesting and brood rearing habitat for waterfowl. It is a popular hunting location during the fall hunting season. Upstream is Morgan Lake, and then Bridgeman Pond. These two wetlands also have potential for nesting and brood rearing waterfowl, as well as provide hunting sites during season. Closest to the office are the North Haystack Pond and the South Haystack Pond. Downstream is Giffin Lake. This is the largest wetland on the unit. An electric pump station delivers water to the Rice Paddy Wetlands. This is a complex of man-made wetland cells and the most popular hunting wetlands on the unit. Soil types do not retain water for very long after the pumps are shut down. Cost of electricity determines how long and when water is supplied to these wetlands. Repairs need to be made to the berms that separate the individual cells so they will hold more water during the hunting season. The 90-acre Field also receives water from pumping out of Giffin Lake. The soil types prevent this area from holding water without pumping. This wetland area provides ideal nesting habitat, and some hunting opportunities throughout the season. The Brady Wetland provides excellent nesting and brood rearing habitat for waterfowl, and is a popular hunting wetland. All of these wetlands can receive water from flood events of the Yakima River. Their primary source of water is irrigation water. At the west end of the unit is the South Emerald Oxbow. This wetland does not have any consistent water supply, only flood water. This wetland does provide nesting and brood rearing habitat. This unit is the first property



Sunnyside Unit Wetlands



acquired by WDFW in the Sunnyside / Snake River Complex. Wetland management has been ongoing since then and much of the infrastructure is old and in need of replacement or upgrades.

Byron Ponds Unit

The Byron Ponds can provide excellent nesting and brood rearing habitat for waterfowl, along with tremendous waterfowl hunting opportunities. Consistent water supply during critical time periods and the ability to manage weeds effectively dictate the high or low level of productivity in the past. Water is supplied from an irrigation ditch, and from excess water released by the neighboring Grandview Water Treatment Plant. In recent years, the amount of water available has decreased, causing wetland water levels to decrease, and allowing emergent vegetation to increase above desired amounts.



Byron Unit wetlands

<u>Mesa Lake Unit</u>

Mesa Lake is a long finger lake that is supplied by irrigation water. Although there is a water control structure on the downstream end of the lake, water level management is primarily controlled by the local irrigation district. Some nesting and brood rearing occurs around the lake. In the northwest corner of the property is the 7-acre Wetland. This wetland receives water from an irrigation ditch, and groundwater seepage from the adjacent hillside. Even though the irrigation water can be periodically diverted away, the amount of groundwater makes management efforts difficult. Also north of Mesa Lake, are the Upper and Lower Pigeon ponds. These two wetlands receive water from the same irrigation ditch as the 7-acre wetland. Water can be diverted at the ditch to the 7-acre wetland or the pigeon ponds, or split between both wetlands.

There are three different wetlands on the south side of the property. The Hilltop Wetland is east of the privately-owned orchard (on top of the mesa). This wetland receives runoff water from the orchard and there is no water control structure for water level management. This wetland also lies within the city limits of Mesa. **The Serpent Wetland** typically supports only some nesting and little brood rearing. The water for



this wetland is supplied by a pump system shared with the neighboring orchard. Soil types within the wetland allow water to seep away, which makes year-round water cost prohibitive. The seepage allows for a



Mesa Lake Unit wetlands

natural moist soil draw-down. The Pie Wedge Wetland sits within the agricultural crop field on the south end of the property. This wetland currently does not have a water control structure and efforts will be made to install one in the future.

Windmill Ranch Unit

At the north end of the property is Powerline Lake. This lake is very deep, with steep rock sides. Nesting and brood rearing is limited. Management actions are limited. Some waterfowl hunting does occur. This body of water is focused primarily on fish recreation. At the west end of the lake is the Phase 1 Wetland. This wetland receives water from Powerline Lake through a gravity-fed gate. In the northeast corner of the property is the Northeast Wetland. This wetland currently does not have a water control structure for water level management. This needs to be addressed before any management activity can occur. South of the Circle #1 crop field is the North Irrigation Pond. This wetland provides excellent nesting and brood rearing habitat. This wetland receives water directly from the irrigation canal and feeds additional wetland habitat downstream. The Circle #3 Wetlands sit inside crop field circle #3. This is a series of small wetlands with grass borders, terraced downhill through the crop field. They provide excellent nesting and brood rearing habitat. These wetlands receive water directly from the irrigation pivot. In the middle of the property is Windmill Lake. This lake currently has an undersized overflow pipe, and no water control structure for water level management. A water control structure will need to be installed. At the far south end of the property is the Phase 3 Wetlands. This is a series of small wetlands terraced downhill. Water is supplied to these wetlands from a diversion on the irrigation ditch. Across the service road is Worth Lake. This lake is mostly deep, open water with steep banks. Worth Lake is supplied by overflow water from Windmill Lake,



further upstream. Adjacent to Worth Lake is the Circle #8 Wetland. This wetland sits in crop field #8 with a grass border. This wetland is supplied directly by the irrigation pivot.



Windmill Ranch Unit wetlands



Bailie Ranch Unit

The Bailie Unit is a privately owned and operated ranch. WDFW has a perpetual easement to allow public hunting. The easement allows WDFW to participate in wetland management with coordination from the ranch. Bailie Lake is a large lake at the south end of the property. The east half of the lake is open water, but the west half is a dense stand of cattails. An irrigation ditch flows north to south through the unit. The amount of water is variable throughout the year, but it is constant.

There are four main wetlands in the middle of the Bailie Youth Ranch. The Stick Wetland is the furthest east. This wetland is supplied by a culvert that diverts water from the main irrigation ditch. The V.I.P. Wetland is a large wetland where the irrigation ditch opens up into a large basin. The Big Bend Wetland receives backwater from the irrigation ditch. This wetland is a depression formed within the basalt rock at the ground surface. The Warm Water Wetland is the furthest west. This wetland receives spring water from the adjacent ridge. The constant flow of spring water adds to the difficulty in managing this wetland. Another wetland is situated in the northeast corner of the property. This is Crescent Lake. This lake receives a combination of spring water and backwater from the irrigation ditch.



Bailie Ranch Unit wetlands



Hope Valley Unit

The Hope Valley Unit contains of a single, large irrigation water supply lake, Clark Pond. The south half of the lake is open water, but the north half is a dense stand of cattails. Water enters the property from the north via an irrigation ditch. The lake has a single outlet at the south end. All water management is conducted by the irrigation district.



Hope Valley Unit wetlands



Esquatzel Coulee Unit

There is only one lake on this property. Esquatzel Lake is a deep open water lake with steep banks. The lake is shallow at the north end where irrigation water enters the lake, and at the south end where water flows out of the lake. There is no man-made infrastructure for managing the water level of this lake. Emergent vegetation is minimal and management actions available are minimal. Although many people access Railroad Lake from this unit, this lake is on private property.



Esquatzel Coulee wetlands



Thompson Seeps Unit

This unit consists of a wide canyon that collects wasted irrigation water and seepage from a large irrigation canal. As water meanders down through the canyon, it collects in multiple open areas creating wetland habitat. Previously there were two lakes on the property created by beaver dams and debris flow. Changes in the water regime by the irrigation district and the wear of time caused the lakes to diminish. There are no manageable wetlands on this unit at this time, but efforts are already being made to create a plan for developing a series of manageable wetlands on this unit.



Thompson Seeps Unit wetlands



I-82 Ponds Unit

The I-82 Ponds unit is a series of land parcels situated between the Yakima River and Interstate 82. There are six ponds on the unit. These ponds were initially borrow pits for fill material during the construction of the interstate. These ponds have been intended to provide public fishing opportunities. The ponds are stocked with fish and the parking areas are managed by the Access Program. The rest of the property consists of riparian habitat and old river oxbows, and provides access to the Yakima River. The wetland habitat on this unit is not actively managed. The high frequency of flooding precludes the installation of water control structures.



I-82 Ponds Unit wetlands



Part III: Management Actions

Management actions identified for wetland management during the next 10 years include:

- Implement a 10-year plan for wetland management and restoration, which is focused on enhancements
 and includes development of new wetlands on the wildlife area. This will include prioritizing areas for
 restoration, working with partners to identify common goals, identifying funding sources, working with
 conservation districts to help leverage opportunities with other land owners, minimize conflicts with
 upland bird habitat, and tie in with other state plans, conservation initiatives, and other partners.
- Coordinate with the Yakima Basin Integrated Plan and their efforts in the Wapato Reach, as well as other areas of the Yakima River system.
- Coordinate with the City of Grandview to ensure a consistent supply of water to the Byron Ponds for wildlife and recreation. This will also include coordination with the Mosquito Control District and the Washington Department of Health.
- Develop a protocol for management of nuisance beavers on the wildlife area by 2022. This will include coordination with irrigation districts, the Yakama Nation, and other organizations.
- Maintain and restore floodplain functions throughout the Yakima River Valley. Coordination with irrigation districts, the Yakima Basin Integrated Plan, County Agencies, and other conservation organizations.
- Evaluate the function and benefits to wildlife and recreationalists of the refuges and reserves for
 promoting the greatest ecological value. The evaluation should be completed by 2022. There are
 established reserves on the Sunnyside and Byron units, and a waterfowl refuge on the Yakima River
 affecting the Vance-Ferry unit. The evaluation with include coordination with the Game Division,
 Toppenish National Wildlife Refuge, and other regional staff.
- Develop a prescribed burn plan for wetlands by 2022. Input will be received by the Agency Burn Team and local fire districts.
- Develop regularly scheduled waterfowl surveys for use in assessing management actions. Wildlife area
 and regional staff shall determine the most appropriate information to collect, and then establish
 protocols and schedules for the surveys.
- Identify opportunities to improve / increase waterfowl hunting opportunities. This will include identifying areas for placement of blinds, improving field-hunting opportunities, and collecting data for user-driven desires.
- Improve fishing opportunities by managing aquatic weeds in stocked lakes. Currently trout are stocked in Ponds 1-6 of the I-82 Ponds unit and Powerline Lake on the Windmill Ranch unit. Wildlife area staff will work with the Fish Program to collaborate on aquatic weed control efforts and evaluate other lakes/ponds/wetlands for potential stocking efforts.



- Water level management is crucial to wetland management. Wildlife area staff will maintain and upgrade water control structures as necessary. Wildlife area staff will coordinate with Ducks Unlimited whenever possible, as well as seek grants and additional funding from other sources. Water control structure replacement / upgrade is needed at:
 - Giffin Lake Sunnyside Unit
 - Bos Lake Sunnyside Unit
 - Pie-Wedge Wetland Mesa Lake Unit
 - Windmill Lake Windmill Ranch Unit
 - Northeast Wetland Windmill Ranch Unit
- Many wetlands are overgrown with emergent vegetation, thus reducing the wetland value and function.
 Wetlands identified for vegetative restoration efforts include:
 - Morgan Lake Sunnyside Unit
 - Bridgeman Pond Sunnyside Unit
 - All of the ponds at Byron Unit
 - Circle #3 Wetlands Windmill Ranch Unit
 - V.I.P. Wetland Bailie Unit
 - Bailie Lake Bailie Unit
 - Clark Pond Hope Valley Unit
 - The Rice Paddies are popular wetlands for waterfowl hunting and recreational bird watching. The low-head berms that create the paddies are in need of restructuring. Wildlife area staff will work with other regional staff to develop a system for maintaining the each berm and providing for movement of water between each cell.
 - The Johnson Wetland is a popular wetland for waterfowl hunting. Staff will work with the local Agriculture Permit holder to increase food resources in the wetland. Staff will also work with irrigation districts to maintain adequate water supplies throughout the hunting seasons.
 - Staff will work with Ducks Unlimited, the Bureau of Reclamation, and local irrigation districts to
 continue developing wetland habitat throughout the Thompson Seeps Unit. The unit is a waste way
 for excess irrigation water. Decadent vegetation and the lack of water management capabilities
 hinders the unit from reaching its full wetland potential. Clearing old overgrowth, developing lowhead berms and installing a few water control structures would improve the wetland function
 throughout the unit.



Appendix E. Cultural History Summary

Early history

Indigenous peoples inhabited the Columbia Basin and Lower Yakima Valley for at least 12,000 years. The Tribes and Bands of the Yakama Nation inhabited the Lower Yakima Valley associated with the western units of the wildlife area. The Waunapum, Wawyukma, and Chamnapum inhabited parts of the Columbia Basin associated with the eastern units of the wildlife area. Settlements consisted of both permanent and winter villages, and spring campsites. Native Peoples fished, harvested root crops, collected berries, gathered other culturally significant plants, and hunted. The Yakamas and others later adopted horses, which aided with hunting big game animals. Native peoples in the region had an extensive trade network for exchanging resources.

Arrival of immigrants

Regular contact between Euro-Americans and Native Peoples began in the 19th century. Trading posts were established by the Pacific Fur Company, the North West Company, and the Hudson Bay Company. Missionaries followed the traders. In the later decades of the 19th century, settlers began arriving in the region in large numbers. The Yakima Treaty created the Yakama Indian Reservation in 1855.

The Indians of Washington were affected by a gradual loss of territory, autonomy, and sometimes of personal and cultural identity. However, due to the initially large number of the Yakama, and their long resistance to and separation from the encroaching Euro-American culture, the Yakama have been able to preserve and use their native language within their communities.

Homestead settlement

The earliest "settlers" to the Columbia Basin were merely passing through on their way west of the Cascades. After tensions with Native Peoples began to subside, settler numbers began to increase. Grazing cattle on the lush bunch grasses of the basin was the economic factor that prompted people to stay in the area.

Due to the low seasonal precipitation of the basin, farming did not become economically feasible until irrigation projects began in the late 1860s. Early irrigation efforts were private companies, and small in scale. The Newlands Reclamation Act of 1902 permitted the Federal Government to participate directly in land reclamation and irrigation projects, and by 1910, Yakima County had the highest percentage of irrigated land in the state.

In 1943, the federal government took control of much of the land north of Benton City and Richland. The purpose was unknown to the residents at the time. Local landowners were forced to sell their property, and the towns of Hanford and White Bluffs were condemned. The federal government established the Hanford Nuclear Reservation for the Manhattan Project of World War II.

The communities of the Tri-Cities and Yakima are some of the fastest growing in Washington State, and the country. These communities are still greatly affected by the Hanford Nuclear Reservation, and agriculture continues to grow and support the local economy.



Wildlife Management

The Columbia Basin and Lower Yakima Valley fish and game populations are quite diverse. The combination of wetland, riparian, riverine, and upland shrubsteppe habitats supports a variety of large and small terrestrial and aquatic mammals; as well as birds, reptiles, amphibians, and insects.

In 1972, a small group of elk colonized the shrubsteppe habitat of the Hanford Nuclear Reservation. This herd has grown rapidly because of high reproductive success. The topographically diverse habitat of the reservation, and the minimal harassment associated with the high security of the nuclear reservation are also believed to contribute to the elk population growth.

Today, people primarily use the Sunnyside-Snake River Wildlife Areas for hunting and fishing. Waterfowl hunting is the most popular recreational activity. Big game (deer and elk) hunting and upland bird hunting are also popular. Two units (Sunnyside and Hope Valley) are designated pheasant release sites. The wildlife area provides access to the Yakima River for hunting and fishing, and a half dozen ponds and lakes are stocked with trout. The various units also attract hikers, bird watchers, mountain bikers and horseback riders.

Cultural Resource Surveys

State and federal laws require WDFW to review sites for archeological resources on any grounddisturbing project. The agency then evaluates the findings and devises a cultural resource protection plan to protect the integrity of historic properties. WDFW has conducted an extensive cultural resource survey of the wildlife area. Five units of the wildlife area provide mitigation credits to Bonneville Power Administration for dams built on the Columbia River. Three other units were purchased as habitat mitigation to the U.S. Army Corps of Engineers for dams on the Snake River.



Appendix F. Public Response Summary (SEPA)

Sunnyside-Snake River Wildlife Area Management Plan

SEPA comments and other comments. SEPA Review Period: September 9 – October 21, 2019

Date & Source	Comment	WDFW Response
9/20/19 SEPA Desk Dale and LouAnn Schielke <u>dahlialo</u> <u>u@msn</u> .com	Thompson Seeps ponds project needs to be actively managed for a timely project completion. The ponds at lower Thompson Seep have historically been excellent angler and hunting destinations. Beaver dams are believed to have provided the impoundments that produced the waterfowl and fish haven. However the water levels fluctuated significantly with beaver activity and subsequently dropped to the current "mudhole" levels. The area is not conducive to waterfowl and the limited remaining muddy water is infested with carp. Ducks Unlimited has produced plans for restoring the ponds with controlled water level structures. Now we just need to implement the plan. These ponds were once (20 plus years ago) so productive that anglers from Seattle area would come to fish them. I urge management of the project in a timely manner to restore this opportunity.	Work on the Thompson Seeps Unit is identified in the draft plan, Appendix D: Wetland Management Plan (page 160). Changes in the water regime by the irrigation district and changes over time have diminished the lakes. WDFW is working with Ducks Unlimited, the Bureau of Reclamation, and other partners to evaluate potential site improvements, and are in the preliminary stages of planning. It will take coordination, time, and funding to develop a plan and begin implementing these projects.
9/6/19 SS-SR email Chris Albertson Christopher. albertson@ srd.edu	Hello, I received the Wildlife Area Management plan and left a voice message as well. My name is Chris Albertson and I'm a Math Teacher at Richland High School. I started an organization called Runners of the Sage <u>www.runnersofthesage.com</u> a couple years ago and our group has done a lot of group runs out in the Rattlesnake area. We put on events throughout the tri-cities with the goal of encouraging all to live more active, sustainable, and environmentally aware lifestyles. You can look at all we are doing on the website above. Our group is growing steadily and we absolutely love the Rattlesnake Slope Preserve and would like to help that area. We have an army of people willing to do work and raise funds and I would love to talk to the right people about what we are thinking. Thanks for your time. Chris Albertson, 503-756-8621	Thank you for your interest in Rattlesnake Slope. WDFW welcomes and appreciates efforts of volunteers, and these efforts need to be coordinated with the wildlife area manager, who will be in contact with you.



SEPA Desk	TO: Scott McCorquodale, Region 3 Wildlife Program Manager; Ross Huffman,	Thank you for the support of the wildlife area
10/7/19	Region 3 Lands Operations Manager; Pat Kaelber, Wildlife Area Manager;	management plan. Weeds, encroachment, and
	Patricia Jatczak, WDFW Lands Planner; Noel Ferguson, Assistant Manager	reduction of habitat and recreational
	SSWMA; Cynthia Wilkerson, Lands Division Manager; Mike Ritter, District	opportunities are of concern to WDFW.
	Habitat Biologist; Members of the Sunnyside-Snake River Wildlife Area	Wildlife area managers will implement the
	Advisory Committee	Weed Management Plan (Appendix B) and the
		Wetlands Management Plan (Appendix D) of
	RE: Public Input - Sunnyside-Snake River Wildlife Management Plan	the management plan to tackle some of these
	I offer this letter in lieu of my personal attendance at the up-coming public	concerns.
	meeting at the Benton County PUD October 10 and any others prior to the	
	Public Input deadline October 21. Please log and distribute this letter formally	Appendix D of the management plan is the
	as Public Comment under the State Environmental Policy Act (SEPA) toward	"Sunnyside-Snake River Wetland Management
	the Sunnyside-Snake River Wildlife Management Plan.	Plan". This plan will be updated by mid-2020,
		and will focus on the Windmill, Bailie, Mesa
	First, a big thanks to each of you and many others named in the production of	Lake, Sunnyside, and Byron units. It will
	the Draft Report. As an outdoor enthusiast and grateful resident of	include a treatment plan and implementation
	Washington, I observe all over the state the breadth and challenges of the lands	schedule.
	you manage. While I am writing on specific matters below, please know that I	
	appreciate the work you all do every day.	
	My comments today are focused on the worsening condition of habitat	
	vegetation access and safety in wetland areas of the SSR-WMA Since L began	
	frequenting the units of the SS-WMA in 1992 I have witnessed a steady decline	
	in the overall quality of these four crucial vectors of public lands management.	
	While I am not familiar with every waterbody in this WMA. I am intimately	
	familiar with a number of them listed here:	
	Glover Unit (51 Acres); Bos Lake; Giffen Lake; Vance-Ferry Road Unit; Morgan	
	Lake; Bridgeman Pond; Johnson Wetland	
	These units in particular, as you have articulated on p. 160 of the Draft Plan,	
	suffer from unrestrained infill of cattails and other aggressive aquatic	
	vegetation in their wetland assets. I am not a biologist but I observe, in	
	wetlands such as Morgan Lake and Bridgeman Pond, a dramatic loss of open-	



water acreage. As a waterfowler I perceive that this loss means migratory waterfowl have fewer open-water resting options adjacent to the Yakima River and probably a less-than-ideal nesting situations for local populations.Several years ago on these units, the density of the aquatic vegetation came to occlude what I consider, reasonable access for even 'very physically fit' hunters. The impenetrable margins of these waters increases loss of downed birds, and risks of injury and drowning. The SS-WMA use to be a great place to introduce young hunters to waterfowling - but now there are much safer (if farther) places to go in the state for that. All of these lakes are soft-bottom - due to St. Helens ash, farming and local geology - this makes wading treacherous and it's only compounded with 20-50 yards of thick cattails and rushes to open water. Small, non-motorized watercraft are not a viable option due to the distances from parking areas and then through entangle vegetation to access open water. Even engineered canals and flood- gate mechanisms have become choke with aquatics.The wee wee uses wildI was hopeful a few seasons ago when I saw an amphibious weed-control machine parked on the southeast side of Morgan Lake – but no significant impact was made with that or other machines.WD water in its nice, broad basin, despite the WMA reportedly having unlimited access to Yakima District irrigation water (D-SSR WAMP, p.29). This alone seems like it is such a simple thing to improve conditions for wildlife and quality hunting.WD weet weet significant pressure of public use. I urge you to escalate the management of aquatic vegetation and percentage of open water as a highest, impactful priority in the Plan. I don't favor aerial spraying – but controlled burning, mechanical removal or other diligent management to maintained ba	e Marsh Master is one of the tools used for ed control and beaver management. It is ed on many wetland areas across the Idlife area. DFW has identified where water is being Id up and not getting to the Johnson tland. The updated wetland management in (Appendix D) will provide treatment tails.
--	---

	habitat, access and safety – along with adequate budget allocations will make these assets better and safer all around. With enthusiasm and confidence in all of you, Sincerely Richard Larson 2847 NE 178th Street. Lake Forest Park Washington 9815	
SS-SR email	Greetings, I have been asked by many citizens of Yakima County "why the state agencies don't have to control noxious weeds " I would like to discuss with the powers that be, scotch thistle and houndstongue control efforts in the lower Yakima valley, and need to know who to contact. Also some discussion of toxic plant controls, such as Poison Hemlock which isn't on the County's designation to control, however we ask landowners to be diligent in preventing the spread of it if possible. It is along the Yakima river system and directly boarders pastures and croplands posing potential risk to livestock and farm employees working near it. Any information you can provide as to contacts for regions in the Yakima valley is greatly appreciated. Have a wonderful week, Susan, Maintenance & Outreach Specialist Yakima County Noxious Weed Control Board Cell (509)945-3357 Office: (509)574-2180 Email susan.bird@co.yakima.wa.us	This request has been responded to through email with a link to the Weed Management Plan of the draft Sunnyside-Snake River Wildlife Area Management plan, as well as forwarded to the wildlife area manager for a direct response.
10/10/19 at the Public Meeting Hard copy presented to staff	To: Pat Kaebler, Sunnyside Snake River Wildlife Area Manager CC: Cynthia Wilkerson, Lands Division Manager Kelly Susewinde, Director, Washington State Department of Fish and Wildlife October 10, 2019 Area Manager Kaebler, I am representing the Lower Columbia Basin Audubon Society (LCBAS) with support from Yakima Valley and North Central Washington Audubon chapters that participated in a six year long Sagebrush Songbird Survey. We want to congratulate the Washington State Department of Fish and Wildlife for its	Thank you for the support of the wildlife area management plan and WDFW's efforts to maintain and enhance habitat in this wildlife area. We look forward to seeing the results of the sagebrush songbird survey.



ambitious work in the preparation of this DRAFT 2019 Sunnyside-Snake River	
Wildlife Area Management Plan. We support the Agency's mission conserving	
our state's natural habitats and wildlife and the roles that wildlife areas play in	
conservation as well as recreation for the benefits of all state residents.	
Audubon recognizes that riparian and wetland habitats, though small in	
acreage in the Sunnyside-Snake River Wildlife Area, are critical for the	
maintenance of many species. We support the agency in maintaining	
and enhancing riparian and wetland buffers on the river systems, and,	
providing public access when appropriate. We also support the partnerships to	
accomplish restorations.	
Our comments will address primarily shrub steppe habitat and species because	
we just completed our community science fieldwork for the Sagebrush	
Songbird Survey. Audubon has invested six years and tens of thousands of	
fieldwork hours by hundreds of volunteers gathering data in order to survey	
the Columbia Plateau shrub steppe for obligate songbirds. The Sagebrush	
Songbird Survey is a partnership between Audubon and WDFW. The results	
will be available through the WDFW Wildlife Program, Science Division in	
2020. Meanwhile, we offer comments based on our field experience over the	
years in and around the Sunnyside-Snake River Wildlife Area.	
We support WDFW management objectives to maintain, enhance and restore	
shrub steppe habitat. We surveyed on four of the Sunnyside-Snake River units	
(Hope Valley, Rattlesnake, Thornton, and Windmill).	
Our comments focus on the value of the Thornton Unit.	
Page 42—44 Thornton Unit. This area has some of the best shrub steppe	We appreciate your interest in the Thornton
habitat in the Sunnyside-Snake River Wildlife Areas. We would like to see even	Unit. The Thornton Unit does have excellent
more contiguous quality shrub steppe habitat for this unit. As noted in the plan,	shrubsteppe habitat, and it is important to
the presence of breeding Sagebrush Sparrows indicates the need for large	WDFW to maintain it. The adjacent parcel (10
contiguous patches of habitat. In our survey data for the area, we found this	North 25 East Section 15) is already state
species present as well as another Priority Habitat Species, the Sage Thrasher.	land, owned by Department of Natural
The adjacent DNR parcel (10 North 25 East Section 15) would be a valuable	Resources (DNR), so it does provide that
addition. It would form a connection between the northern and central WDFW	connection between the northern and central
parcels. Adding section 15 is valuable for habitat connectivity, climate refugia,	WDFW parcels for habitat connectivity,
breeding sites, as well as migration stopover sites.	



The DNR parcel shares a common 2-mile border with the WDFW Thornton	climate refugia, breeding sites, as well as
Unit. Audubon is currently working up a Trust Land Transfer parcel	migration stopover site.
nomination with our local chapter, Lower Columbia Basin Audubon Society.	
We will work with WDFW and any other partners to develop a proposal or	WDFW follows a set of principles and strategic
method of acquiring this section of land for addition to the Thornton Unit. We	acquisition priorities
suggest immediate acquisition of the Conservation Reserve Plan portion of the	<u>(https://wdfw.wa.gov/about/wdfw-</u>
Section 15, and acquisition of other two-thirds portion at the end of the current	lands/land-acquisitions#acquisition-
private grazing lease. We suggest talking to the lessee to see if they want out of	priorities) when considering purchasing new
the lease early.	lands. WDFW's rigorous review process is
	designed to determine which properties will
We request that WDFW explore disposition options for Vance-Ferry Unit to	best meet the state's conservation goals and
include a DNR exchange, as well as using potential disposition funds for	recreational priorities. For this reason, the
additional Thornton acquisition. (Page 101, Table 9,10.D).	process is employed even in situations when
We request that you please add language for acquisition to the primary	property owners seek to donate their land.
management section for Thornton:	
We recommend moving Section 15 ownership and management from DNR to	WDFW will discuss with DNR ways to ensure
WDFW. This section has high wildlife values for birds, and for elk and upland	that grazing is conducted in accordance with
bird hunting. WDFW is the only state agency that manages exclusively for	the state ecosystem standards for grazing,
wildlife values rather than multi-uses that can degrade the habitat.	found in <u>chapter 4, Laws of 1993 sp. sess.</u>
Thornton contains four of the six systems of concern:	<u>ecosystem standards</u> .
Columbia Basin Foothill and Canyon Dry Grassland (Critically imperiled-	
imperiled), Inter-Mountain Basins Big Sagebrush Steppe (imperiled).	WDFW does have an objective to determine
Columbia Plateau Steppe and Grassland (Imperiled)	how to manage the Vance Ferry Road Unit. We
North American Arid West Emergent Marsh (Imperiled)	will edit Vance-Ferry Unit objective 10.D to
Thank you for the opportunity to make comments for this important public	include "long-term management".
process.	
Kathy Criddle, Lower Columbia Basin Audubon Society	
Board Member & Sagebrush Songbird Survey Volunteer	





Washington Department of Fish and Wildlife

10/21/19 SEPA Desk	 To: Pat Kaebler, Sunnyside Snake River Wildlife Area Manager CC: Cynthia Wilkerson, Lands Division Manager Kelly Susewinde, Director, Washington State Department of Fish and Wildlife October 21, 2019 Area Manager Kaebler, Audubon Washington with our 25 affiliated chapters and 33,000 members congratulate the Washington State Department of Fish and Wildlife for its ambitious work in the preparation of a DRAFT 2019 Sunnyside-Snake River Wildlife Area Management Plan. We support the Agency's mission conserving our state's natural habitats and wildlife and the roles that wildlife areas play in conservation as well as recreation for the benefits of all state residents. 	Note: The first part of this response is identical to the response to the letter above dated 10/10/2019. Thank you for the support of the wildlife area management plan and WDFW's efforts to maintain and enhance habitat in this wildlife area. We look forward to seeing the results of the sagebrush songbird survey.
	Audubon recognizes riparian and wetland habitats, though small in acreage in Sunnyside-Snake River Wildlife Area, are critical for the maintenance of many species. We support the agency in maintaining and enhancing riparian and wetland buffers on the river systems, and, providing public access when appropriate. We also support the partnerships to accomplish restorations.	We appreciate your interest in the Thornton Unit. The Thornton Unit does have excellent shrubsteppe habitat, and it is important to WDFW to maintain it. The adjacent parcel (10 North 25 East Section 15) is already state land. owned by Department of Natural
	Our comments will address primarily shrub steppe habitat and species because we just completed our community science fieldwork for the Sagebrush Songbird Survey. Audubon has invested five years and tens of thousands of fieldwork hours by hundreds of volunteers gathering data in order to survey the Columbia Plateau shrub steppe for obligate songbirds. The Sagebrush Songbird Survey is a partnership between Audubon and WDFW. Results will be	Resources (DNR), so it does provide that connection between the northern and central WDFW parcels for habitat connectivity, climate refugia, breeding sites, as well as migration stopover site.
	available through the WDFW Wildlife Program, Science Division in 2020. Meanwhile, we offer comments based on our field experience over the years in and around the Sunnyside-Snake River Wildlife Area.	WDFW follows a set of principles and strategic acquisition priorities (<u>https://wdfw.wa.gov/about/wdfw-</u> lands/land-acquisitions#acquisition-
	We support WDFW management objectives to maintain, enhance and restore shrub steppe habitat. We surveyed on four Sunnyside-Snake River units (Hope Valley, Rattlesnake, Thornton, and Windmill).	priorities) when considering purchasing new lands. WDFW's rigorous review process is designed to determine which properties will

Our comments focus on the value of the Thornton Unit. Page 42—44 Thornton Unit. This area has some of the best shrub steppe habitat in the Sunnyside-Snake River Wildlife Areas. We would like to see even more contiguous quality shrub steppe habitat for this unit. As noted in the plan, the presence of breeding Sagebrush Sparrows indicates the need for large contiguous patches of habitat. In our survey data for the area, we found this species present as well as another PHS species, the Sage Thrasher. The adjacent DNR parcel (10 North 25 East Section 15) would be a valuable habitat. addition. It would form a connection between the northern and central WDFW parcels. Adding section 15 is valuable for habitat connectivity, climate refugia, breeding sites, as well as migration stopover sites. The DNR parcel shares a common 2-mile border with the WDFW Thornton Unit. Audubon is currently working up a Trust Land Transfer parcel nomination with our local chapter, Lower Columbia Audubon Society. We will work with WDFW and any other partners to develop a proposal or method of acquiring this section of land for addition to the Thornton Unit. We suggest immediate acquisition of the Conservation Reserve Plan portion of the Section 15, and acquisition of other two-thirds portion at the end of the current private grazing lease. We suggest talking to the lessee to see if they want out of the lease early. We request that WDFW explore disposition options for Vance-Ferry Unit to include a DNR exchange, as well as using potential disposition funds for additional Thornton acquisition. (Page 101, Table 9, 10.D). Please add language for acquisition to the primary management section for Thornton: We recommend moving Section 15 ownership and management from DNR to WDFW. This section has high wildlife values for birds, and for elk, mule deer, and upland bird hunting. WDFW is the only state agency that manages

best meet the state's conservation goals and recreational priorities. For this reason, the process is employed even in situations when property owners seek to donate their land.

WDFW will discuss with DNR ways of maintaining or improving the DNR land to benefit species in this important shrubsteppe habitat.

WDFW does have an objective to determine the management of the Vance Ferry Road Unit. We will edit Vance-Ferry Unit objective 10.D to include "long-term management".



The PHS species sagebrush sparrow and sage thrasher are listed in the species section (Table
11, p. 115). Appendix A has a table showing the relationship between terrestrial Species of Greatest Conservation Need and the Ecological Systems of Concern.
-Snake Wildlife biologists and wildlife area managers are exploring the most effective places to install platform nests for the ferruginous hawk. This would be in areas where there is little or no natural nesting sites. Beaver can be beneficial or can be a nuisance, depending on the situation and the management objective. The intent of developing the beaver protocol is so staff have a set of rules and guidance to help determine how to manage beavers in various situations.
k that ing). The Vision for the wildlife area does describe types of non-consumptive activities such as birding and hiking because we did not want to use the words "non consumptive". We can put them on equal par by removing the words "as well as" and replacing it with "and". its. We will add Thornton to the list of priority units based on its exceptional habitat and recreational/educational opportunities.



Page 22, Primary management objectives I-82 Ponds: We support coordination with the Yakima Basin Integrated Plan in restoring floodplain functions.

Page 30. Sunnyside Unit. We support the use of all non-toxic shot in Sunnyside and all other Wildlife Area units. We suggest making non-toxic shot a requirement for all units due to lead poison issues for birdlife.

Page 29, Paragraph 4 & page 30, Paragraph 3, Giffin Lake – Rice Ponds The Yakima Valley Audubon Society has been impressed with spring water management of the "rice ponds," south of Giffin Lake. For at least the past two springs, water in this managed, diked area have supported an impressive array of waterfowl, marsh and wading birds. Many of these have attempted to nest. By June, water delivery to this area has been turned off and these ponds have become completely dry by the end of the month. This has resulted in nesting failure by wetland-dependent birds and other wildlife. We request that WDFW review the strategy of early spring of filling the ponds, then allowing ponds to go dry during the nesting season. WDFW has explained lack of water rights is the issue here. If water is the limiting resource in this issue, we urge WDFW to explore opportunities in obtaining additional water to effectively manage the ponds as wetlands throughout the nesting season. These ponds have real potential of becoming wetlands of regional and statewide importance with proper management.

The use of shot is regulated by WAC 220-414-040 and is beyond the scope of this plan. On this wildlife area, it is unlawful to possess shot (either in shotshells or as loose shot for *muzzleloading*) other than nontoxic shot for any purpose on the Sunnyside, Byron, and Windmill Ranch Units. Additionally, It is unlawful to possess shot (either in shotshells or as loose shot for muzzleloading), other than nontoxic shot, when hunting for upland game birds (pheasant, quail, chukar, and gray partridge), mourning doves, and band-tailed pigeons on all WDFW designated pheasant *release sites.* On this wildlife area. that includes Hope Valle as well as the four mentioned above.

WDFW will keep water in some of the cells for migrating shorebirds and nesting waterfowl and has repaired some levies. For vegetation management, it is necessary to have some cells dry out.

Appendix D of the management plan is the "Sunnyside-Snake River Wetland Management Plan". This plan will be updated by mid-2020, and will focus on the Windmill, Bailie, Mesa Lake, Sunnyside, and Byron units. It will include a treatment plan and implementation schedule



Page 74. Grazing permits. We support grazing permits that are limited to existing agricultural fields, and note that currently none are in wildlife areas. We suggest adding language to Policy C-6003 contain specific language prohibiting grazing in wildlife areas. Rationale: Grazing spreads cheat grass and other invasives degrading habitat and making it more wildfire prone. DNR is designated to manage state lands for multiple uses including grazing.	It is beyond the scope of this plan to address Fish and Wildlife Commission Policy C-6008. All state public land used for grazing and agriculture must be managed in keeping with the ecosystem standards developed for this purpose, found in chapter 4. Laws of 1993 sp. sess. ecosystem standards.
Page 79. Table 5 Research and Studies Add Sagebrush Songbird Survey to table. Researcher: WDFW, Audubon Date: 2015-2019 Description: Columbia Plateau shrub steppe avian obligate songbird occurrence study	WDFW will discuss with DNR ways to ensure that grazing is conducted in accordance with the state ecosystem standards for grazing, found in , found in <u>chapter 4</u> , Laws of 1993 <u>sp.</u> <u>sess. ecosystem standards</u> . Thank you, we will add the study to the table.
 Page 79-81. Recreation Audubon supports the Rattlesnake Target Range, recognizing a need for safe practice sites during the wildfire season. For wildfire prevention, we suggest no shooting within Wildlife Areas in the fire season other than at formally designated and maintained target ranges. We base this proposal on our survey sites burned by range fires started by stray bullets. Page 88. Wildlife Area goals, objectives and performance measures Audubon understands that objectives are dependent on funding. Audubon proposes full funding for WDFW's budget as a 2020 legislative priority. Management objectives priorities for Audubon: Statewide goal to maintain or improve the ecological integrity of priority sites and systems. 	The Rattlesnake Mountain Shooting Facility is open year-round for shooting practice. It is the only place where target shooting is allowed on the Sunnyside-Snake River Wildlife Area.



	Page 91. Table 9. 1. H: Objective H Identify upland areas with high potential for	We will add the objective to the climate
	habitat improvement and develop action items by 2021. Rationale: Important	change section on Table 17.
	for climate change refugia.	
	Page 95. Table 9. 4. B: Maintain and improve upland bird habitat and nesting	
	cover.	
	Rationale: The North American bird population has drastically declined in the	
	past half-century. Nesting habitat is a critical component for recovery.	
	Citation: Kenneth Rosenberg, Dokter Adriaan et al. The Decline of North	
	American Avifauna. Science Vol. 366, Issue 6461 October 2019. Pages 120-124.	
	Page 130. Table 14 Fires greater than 50 acres.	
	Audubon supports the use of prescribed fire as pioneered effectively at WDFW	
	Sinlahekin Wildlife Area by former Area Manager. Dale Swedburg.	
	Page 132-5. Species of concern with high vulnerability to Climate Change.	
	Audubon supports WDFW efforts to make the goals and objectives of the	Wildlife area staff are working with the hurn
	Wildlife Area Management Plan climate resilient	team to identify some controlled hurn projects
	Thank you for the opportunity to make comments for this important public process	
	inanity ou for the opportantly to make commonity for and important public processi	
	Christi Norman	
	-Christi Norman, Audubon Washington, Program Manager	
	-Jim Castle, President, Lower Columbia Basin Audubon Society	
	-Andy Stepniewski, Conservation Chair, Yakima Valley Audubon Society	
	-Art Campbell, President, North Central Audubon	
	-Judy Hallisey, President, Kittitas Audubon Society	
	-Marie Dymkoski, President, Palouse Audubon Society	
	-Lindell Haggin, Treasurer, Spokane Audubon Society	
	-Sam Merrini, Conservation Chair, Black Hills Audubon	
	-janet Strong, Freshuent, Grays Harbor Audubon Cindy Eastarson, Prosident, Dilchuck Audubon	
	-Cilluy Basterson, Freshvent, Filchuck Audubon Dan Straiffart Conservation Chair Rainier Audubon	
	-Van Streinert, Conservation Chair, Rainer Audubon -Kurt Kirkland Conservation Chair, Tahoma Audubon	
	-Steve Pudiacombe President Willana Hills Audubon Society	
L	store r addecense, r oblacht, trinapa milis madabon boolety	

